

Ber. nat.-med. Verein Innsbruck	Band 99	S. 49 - 66	Innsbruck, Dez. 2014
---------------------------------	---------	------------	----------------------

Effect of long-term cage fish-farming on the phytoplankton biodiversity in two large Bulgarian reservoirs

by

Kostadin T. DOCHIN & Maya P. STOYNEVA *)

Synopsis: The present paper provides data on recent (2010-2012) biodiversity of planktonic algae in two Bulgarian reservoirs, in which cage fish farming ran for more than 30 years (with different composition of fishes). Comparisons are made between the species composition of algae in both reservoirs, and between the recent phytoplankton structure and its structure from previous periods, for which literature data are available. In spite of different degree of impact of the cage fish farming on both reservoirs, a general trend of increase of number of species (double to triple) and shift in phytoplankton taxonomic structure and composition of dominant complexes was proved. However, it has to be noted that this trend ran in parallel with some losses in biodiversity and mainly with enhancement of cyanoprokaryotes and green algae.

Keywords: cyanoprokaryotes, green algae, diatoms, eutrophication, similarity

1. Introduction:

Bulgaria has long traditions in studies of algal biodiversity and recent estimates show that on its territory ca. 6000 algal taxa have been recognized (STOYNEVA 2014). Attention to water reservoirs and their planktonic algae started to be paid in the end of 50s of 20th century, generally simultaneously with the impoundment of new water bodies in the country. In total, data on 23 reservoirs with ca. 250 species had been published (STOYNEVA & TEMNISKOVA-TOPALOVA 2007; STOYNEVA 2014). However, studies were focused mainly on phytoplankton quantitative structure, seasonal dynamics and dominants. Effects of fish rearing on the abiotic parameters and water biota in Bulgaria have been studied in fish farms and fish ponds, for which a total of ca. 600 algal taxa have been published (STOYNEVA & TEMNISKOVA-TOPALOVA 2007; STOYNEVA 2014). The effects of fish cage farming in Bulgaria on physical and chemical water parameters and fish have been firstly documented and discussed by NAUMOVA & ZHIVKOV (1988). Later, mainly chemical data for reservoirs with cage farming were provided (e.g. ILIEV & HADJINIKOLOVA 2011; HADJINIKOLOVA & ILIEV 2012). The subject of cage fish farming impact has attracted the attention of phyto-

*) Addresses of the authors: K. T. Dochin, Institute of Fisheries and Aquaculture, ul. Vasil Levski 248, Plovdiv - 4000, Bulgaria; doksi11@abv.bg; M. P. Stoyneva, Sofia University "St. Kliment Ohridski", Faculty of Biology, Dept. Botany, bld. Dragan Zankov 8, Sofia-1164, Bulgaria; mstoyneva@abv.bg.

planktonologists worldwide, but the main studies have been aimed on its short-term effects and mainly on phytoplankton abundance, with the general assumption that the impact results and eventual danger are due to changed nutrient load and are generally similar to eutrophication (e.g. ARZUL et al. 1996; NOGUEIRA 2000; DEMIR et al. 2001; DIAZ et al. 2001; KARAKASSIS et al. 2005; THANH HUNG et al. 2005; CAO et al. 2007; DEGEFU et al. 2011; BARTOZEK et al. 2014). Generally, since water renewal time is shorter in reservoirs than in natural lakes, changes in ecosystems such as eutrophication resulting from fish cage culture may be less harmful and because of this reservoirs are accepted as very appropriate water bodies for cage culture (DEMIR et al. 2001). However, yet the evidence for long-term effects of different farm capacities is scarce. According to CROSSETTI et al. (2008) and BORGES et al. (2010) the increased eutrophication decreases the phytoplankton diversity and enhances the growth of cyanoprokaryotes, and this is important for water reservoirs, which can be regarded as biodiversity hot-spots (e.g. WILLIAMS et al. 2004). On the other hand, the results of BARTOZEK et al. (2014) did not confirm the hypothesis of changes in the concentration of nutrients for the sites influenced by net cages and on the attributes of the phytoplankton community, including its composition and richness. The aim of the present paper is to compare the effect of long-term cage fish farming (ca. 30 years) on the algal biodiversity in two reservoirs Dospat and Kurdzhali, which were the first reservoirs used for such purpose in Bulgaria.




Figure 1: Maps of the reservoirs Kurdzhali (A) and Dospat with their position in Bulgaria and location of the studied sites (NN1-6, starting from the dam wall).

	Reservoir Kurdzhali	Reservoir Dospat
Geographic coordinates	41°37' 25°20'	41°41' 54'' 24°05' 10''
Average height (m a.s.l.)	280	1 200
Volume (m³)	497 235 698	449 248 693
Aquatory (m²)	15 991 735	22 099 371
Maximum length (m)	23 400	16 700
Average width (m)	1 323	683
Average depth (m)	31	20
Maximum depth (m)	85	50
Watershed basin (km²)	1882	432.30
Retention time (days)	205	180
Water temperature (range; °C)	3.8-26.4	3.9-23
Conductivity (range; µs/m)	294-346	103-118.6
Transparency (Secchi depth; m)	1.2-5.5	2.1-6.5
pH (range)	7.06-9.26	6.47-8.83
NH₄-N (range; mg l⁻¹)	0.001-0.423	0.001-0.685
NO₃-N (range; mg l⁻¹)	0.1-3.48	0.1-2.03
TN (range; mg l⁻¹)	0.12-3.5	0.1-1.924
PO₄³⁻ (range; mg l⁻¹)	0.001-0.38	0.001-0.92
TP (range; mg l⁻¹)	0.03 - 0.215	0.01-0.418

Table 1: Morphometric characteristics and physical and chemical parameters (the last measured during the investigation period 2010-2012) of the reservoirs Kurdzhali and Dospat.

2. Material and methods:

Studied sites are two reservoirs situated in the low (Kurdzhali) and middle (Dospat) parts of Rhodopi Mts in Bulgaria. Their main morphometric characteristics and physical and chemical parameters (the last measured during the investigation period) are given in Table 1. More details on the abiotic features and biodiversity of both reservoirs are provided in MICHEV & STOYNEVA (2007). Cage fish farming started in both reservoirs 30 years ago, but the reared fishes are different: sturgeons (Acipenseridae), common carp (*Cyprinus carpio*) and wels catfish (*Sylurus glanis*) in Kurdzhali and only rainbow trout (*Oncorhynchus mykiss*) in Dospat. According to our knowledge, there is no other significant impact on these reservoirs.

Phytoplankton samples have been taken by Danish bathometer (at each 5 m from the surface layer till the reservoir bottom) and processed by standard method of fixation with 4% formalin and further sedimentation. In both reservoirs sampling was conducted from April 2010 till March 2012 in 6 sites (Figs. 1a, b), with site numbers starting from the dam (1) and cage farms (2-3) down to the tail waters (4-6). Totally 547 samples have been analyzed (307 from deeper Kurdzhali reservoir and 240 from shallower Dospat reservoir).

Microscope work has been done on Bürker blood-counting chamber on microscope “Carl Zeiss Axioscope 2 plus” (magnification 400x) using standard taxonomic literature with critical use of AlgaeBase (GUIRY & GUIRY 2014). Comparisons have been done with the only available data on species composition, published on Dospat for the period 1972-1975 by NAIDENOV & SAIZ (1977) and on Kurdzhali for the periods 2000-2002 and 2006 by TRAYKOV (2005) and BELKINOVA et al. (2007), respectively. The similarity was calculated by the standard Sørensen's Similarity Index (SSI), according to the formula $(2c/a+b)$, where: a - number of species in site (period) a, b - number of species of site (period) b, and c - number of common species for sites (periods) a and b.




Figure 2: Taxonomic structures of the phytoplankton from the reservoirs Kurdzhali and Dospat during different periods of investigations (1973-1975 – acc. to NAIDENOW & SAIZ (1977); 2006-2012 – acc. to combined data of TRAYKOV (2005) and BELKINOVA et al. (2007); 2010-2012 – data from present study).




Figure 3: Sørensen's Similarity Index (SSI) between different taxonomic groups of algae, recorded in the reservoirs Kurdzhali and Dospat in the period 2010-2012. Abbreviations: Cya – Cyanoprocaryota; Eug – Euglenophyta, Pyr – Pyrrhophyta, Cry – Cryptophyta, Chr – Chrysophyceae, Syn – Synurophyceae, Bac – Bacillariophyceae, Chl – Chlorophyta and Str – Streptophyta.

3. Results and discussion:

During our studies in Kurdzhali 137 taxa from 6 divisions have been identified: Cyanoprokaryota (21), Euglenophyta (7), Pyrrophyta (2), Ochrophyta (51: Bacillariophyceae - 48; Synurophyceae - 3), Chlorophyta (44) and Streptophyta (12) – Table 2, Fig. 2. In Dospat 131 taxa from 7 divisions have been identified: Cyanoprokaryota (18), Euglenophyta (11), Pyrrophyta (6), Cryptophyta (3), Ochrophyta (62: Bacillariophyceae 58; Chrysophyceae – 1, Synurophyceae - 3), Chlorophyta (22), Streptophyta (8) – Table 2, Fig. 2. Common for both reservoirs are 70 taxa and the Sørensen's similarity index (SSI) between them is 0.52. Similarity by different taxonomic groups ranged between 0.28 and 0.44 and together with common taxa in each group is shown on Fig. 3.

Data on phytoplankton of both reservoirs under investigation, published before this study, are quite scarce. According to TRAYKOV (2005), the phytoplankton of Kurdzhali Reservoir in the three studied by him sites during the period 2000-2002 contained 55 taxa from 6 divisions (Table 2) and the number of species during different months of his investigation ranged between 28 and 55. The author did not provide detailed data on seasonal differences in the species composition and dominant species, but mentioned the summer mass development of *Fragilaria crotonensis* and "intensive" autumn bloom of *Microcystis aeruginosa*, *Anabaena* sp. and *Aphanizomenon flos-aquae*. In 2006, based on two summer samplings in July and September in four sites of the same reservoir, BELKINOVA et al. (2007) also published 55 algal taxa from 6 divisions (Table 2). The authors noted that the number of species in July and September 2006 was equal (40) but common for both months were only 24, with a big difference in the species composition of cyanoprokaryotes. According to the two cited above works, the highest number of taxa belonged to green algae and the highest number of species was detected in the tail waters. BELKINOVA et al. (2007) marked 17 algae as abundant in their taxonomic tables and outlined in the text the following dominants: *Synechococcus elongatus*, *Fragilaria crotonensis*, *Synedra ulna*, *Stephanodiscus hantzschii*, *Ankyra judayi*, *Monactinus simplex* and *Eudorina elegans* (Table 2). Obviously, the comparison of data for the periods 2000-2002 and 2006 with our results is difficult due to the big differences in the number of investigated samples and their horizontal and temporal distribution. However, some similarities and dissimilarities could be outlined. Similarity coefficient of Sørensen (SSI) between the periods 2000-2002 and 2009-2012 is only 0.25, 2006 and 2009-2012 – only 0.38, and when data from both previous studies are combined (2000-2006) the similarity between them and 2009-2012 is only 0.38. This low similarity is easily explainable with differences in the sampling design, but since the studies from two different previous independently came to the same general number of 55 algae for each of them, and our investigation proved presence of 137 species (with notable increase of number of cyanoprokaryotes, green algae and diatoms – Fig. 2), it is possible tentatively to suggest a trend of general increase of biodiversity due to augmented nutrient loading as effect of long-term cage fish farming. However, the effect on phytoplankton biomass and quantitative distribution is not so obvious – the average biomass detected by us is on conformity with data, provided by BELKINOVA et al. (2007) and the quantitative phytoplankton

distribution and trophic gradient follow the normal spatial and temporal patterns, known for temperate reservoirs with general increase from dam to tail waters (DOCHIN, in prep.).

According to NAIDENOW & SAIZ (1977), the phytoplankton in Dospat Reservoir comprised of 45 algal taxa, found in a three-year period (1972-1975). The authors explained this low number with the low insolation of this reservoir, peculiar by its narrow shape and deep situation in a mountain gorge. Another feature of the qualitative composition was the similar number of species of diatoms and green algae (19 and 18, respectively). The authors provided a generalized table of species composition by seasons, where 19 taxa were pointed for spring, 35 – for summer and 31 – for autumn period. In the text, the number of algae (without flagellates), detected annually during different seasons was much lower – 12 species in spring (11 diatoms and 1 green), 11 species in summer (1 diatom and 5 greens) and 14 in autumn (9 diatoms and 5 greens). The comparison of these data with our results shows significant (triple) increase of the species number (from 45 to 131 – Table 2), and also a change in the taxonomic structure and enrichment of the phytoplankton with the group of cyanoprokaryotes (Fig. 2). The increase in the number of species belonging to euglenophytes and pyrrhophytes during the 2010-2012 period is also notable (Fig. 2). In contrast with the situation in the 70s, when diatoms and green algae have been almost equally represented in the phytoplankton, during our study, diatoms have clear leading role in the species composition – they comprise 45% of all species found, while green algae occupied second place. Detailed analysis of the species composition also shows significant shifts with “appearance” and “disappearance” of given species (Table 2) during the last 30 years. For some of them, like *Uroglena volvox* and *Synura uvella*, ‘disappearance’ was predicted by former authors since they accepted these species as a part of the “primary biopool” (NNAIDENOW & SAIZ 1977). Other, algae from the same “primary” group, also known as worldwide broadly distributed, like *Trachelomonas volvocina* and *Eudorina elegans*, continue to grow in the reservoir in recent times. In general, most of the “new” taxa found, are algae, pointed in standard manuals and floras as common members of eutrophic waters (e.g. *Dolichospermum spiroides*, *Planktothrix agardhii*, *P. rubescens*) but some are with not well-known ecology (e.g. *Limnococcus limneticus* – KOMÁREK & ANAGNOSTIDIS 1999). Common for both periods of investigations are 22 taxa, which means that their similarity (acc. to Sörensens’ coefficient) is only 0.25.

Differences concern also the composition of dominants and subdominants. In 70s, they were represented mainly by diatoms, one dinoflagellate and two green streptophyte algae, whereas during our studies, in addition to them, chlorophyte green algae and cyanoprokaryotes appeared in dominant complexes (Table 2). The “new” complexes include some species typical for oligotrophic waters (e.g. *Gymnodinium uberrimum* – STARMACH 1974; POPOVSKÝ & PFIESTER 1990), but also some well-known harmful algae, producers of cyanotoxins, related with eu- and even hypertrophic waters (e.g. *Aphanizomenon flos-aquae*, *Planktothrix rubescens*). It is difficult to make strong conclusions, based on dominant composition only, since for some of the species different ecological characteristics are provided from different authors. For example, *Staurostrum gracile* according to COESEL

(1997) is mesotrophic, according to LENZENWEGER (1997) – oligotrophic, while in the opinion of PALAMAR-MORDVINTZEVA (1982) it is broadly distributed and strongly variable in different water bodies. According to the previous studies of one of the authors, this species was found by us in strongly eutrophicated Bulgarian lakes Srebarna and Durankulak (STOYNEVA 1998, 2000 ab). In Dospat, this species was pointed once among July 1973 dominants (NAIDENOV & SAIZ 1977), while in our studies it was relatively equally distributed in the reservoir aquatory, but was not recorded in the dominant complexes. However, the detected recent shift in the taxonomic structure of dominant complexes (with appearance of cyanoprokaryotes, chlorophytes and small centric diatoms) combined with their diversification with twice increased number of species (from 8 in 70s to 17 nowadays – Table 2), indicates, in our opinion, the trend of water eutrophication caused by the long-term cage-fish farming and increased organic matter in the reservoir due to non-utilized (uneaten) food from the cages. The increased trophic state of Dospat waters was proved also by quantitative phytoplankton spatial distribution and double increase of average annual algal biomass (from 0.7 in 70s to 1.35 mg l⁻¹ – DOCHIN & STOYNEVA, in press). In addition, in Dospat, much more abundant phytoplankton was detected near the dam and cage farms than in tail waters (DOCHIN & STOYNEVA, in press).

In conclusion, the results obtained during this study allow us to suggest gradual general increase in algal biodiversity of the phytoplankton communities in two Bulgarian reservoirs, combined with shifts in dominant complexes and in total composition with some species losses and enhancement of harmful species, caused by long-term fish cage farming (ca 30 years). It is possible to suggest also that due to accumulation effect, this process is gradually speeded-up. For example, during the last 30 years in Dospat the number of species showed triple increase, while the recorded increase in Kurdzhali in the period of the last 6-12 years there was double increase in the species number. The comparison between the effects of cage fish farms in both reservoirs shows that in Dospat the impact is strongly expressed than in Kurdzhali. Most probably, the larger volume of Kurdzhali, its deeper character and longer insolation are among the reasons for these differences. There are almost no data concerning the influence of species composition of reared fishes on water quality and therefore for the moment this factor is not discussed here in detail. The cage impact concerns the general more notable change in the biodiversity, in the phytoplankton abundance and distribution, and in the general trophic status in Dospat reservoir. The lower trophic state of Kurdzhali and the “normal” pattern of spatial distribution of phytoplankton abundance indicates lower impact from fish cages, and the main data, which suggest such influence could be found in the recorded double increase of algal biodiversity and dominant shifts during the last 12 years.

Table 2: Species composition of the phytoplankton from the reservoirs Kurdzhali (K) and Dospat (D) during different periods of investigations (1973-1975 – acc. to NAIDENOV & SAIZ 1977; 2000-2002 – acc. to TRAYKOV 2005; 2006- acc. to BELKINOVA et al. 2007; 2010-2012 – data from present study): xxx – dominant, xx – subdominant; x - occurrence.

Taxa	K-2000-2002	K-2006	K-2010-2012	D-1972-1975	D-2010-2012
Cyanoprokaryota					
<i>Anabaena sphaerica</i> Bornet et Flahault			x		
<i>Anabaena</i> sp.	xxx				
<i>Anabaenopsis</i> sp.					x
<i>Anathece clathrata</i> (West et G.S.West) Komárek, Kastovsky et Jezberová			x		
<i>Aphanizomenon flos-aquae</i> Ralfs ex Bornet & Flahault	xxx	xxx/x	x		xxx/x
<i>Aphanocapsa</i> spp.			x		x
<i>Calothrix</i> sp.					x
<i>Chroococcus dispersus</i> (Keissler) Lemmermann			x		
<i>Chroococcus minutus</i> (Kützing) Nägeli			x		
<i>Chroococcus turgidus</i> (Kützing) Nägeli			x		x
<i>Coelosphaerium kutzingianum</i> Nägeli			x		x
<i>Dolichospermum affine</i> (Lemmermann) Wacklin, Hoffmann et Komárek		xxx/x			
<i>Dolichospermum flos-aquae</i> (Brébisson ex Bornet et Flahault) Wacklin, Hoffmann et Komárek			x		x
<i>Dolichospermum solitarium</i> (Klebahn) Wacklin, Hoffmann et Komárek		xxx	x		
<i>Dolichospermum spiroides</i> (Klebahn) Wacklin, Hoffmann et Komárek			x		x
<i>Eucapsis alpina</i>		x			
<i>Limnococcus limneticus</i> (Lemmermann) Komárková, Jezberová, O. Komárek et Zapomélková			x		xx/x
<i>Merismopedia glauca</i> (Ehrenberg) Kützing	x				
<i>Microcystis aeruginosa</i> (Kützing) Kützing	xxx	xxx/x	x		x
<i>Microcystis pulvereae</i> (H.C. Wood) Forti emend. Elenkin		x	x		
<i>Noctoc</i> sp.					x
<i>Oscillatoria limosa</i> C.Agardh ex Gomont			x		xx/x

Taxa	K-2000- 2002	K-2006	K-2010- 2012	D-1972 1975	D-2010- 2012
<i>Oscillatoria</i> spp.	x				
<i>Phormidium willei</i> (Gardner) Anagnostidis et Komárek			x		
<i>Phormidium</i> sp.	x		x		x
<i>Planktothrix agadhii</i> (Gomont) Anagnostidis et Komárek					x
<i>Planktothrix rubescens</i> (De Candolle ex Gomont) Anagnostidis et Komárek					xxx/xx
<i>Scopulonema</i> sp.			x		
<i>Scytonema</i> sp.	x				
<i>Snowella lacustris</i> (Chodat) Komárek et Hindák			xxx/xx/x		
<i>Spirulina major</i> Kützing ex Gomont					x
<i>Synechococcus elongatus</i> (Nägeli) Nägeli		xxx	x		
<i>Woronichinia naegeliana</i> (Unger) Elenkin			x		
Chlorophyta			x		
<i>Actinastrum hantzschii</i> Lagerheim					x
<i>Actinastrum hantzschii</i> var. <i>gracile</i> V. K. Tschernov			x		
<i>Ankyra judai</i> (G.M. Smith) Fott		xxx/x	x		
<i>Ankyra occellata</i> (Korshikov) Fott			x		
<i>Ankyra</i> sp.					
<i>Carteria</i> sp.			x		
<i>Chlamydomonas reinhardtii</i> P.A.Dangeard			x		
<i>Chlamydomonas</i> spp.	x		x		
<i>Coelastrum microporum</i> Nägeli in A. Braun		x	x		
<i>Coelastrum sphaericum</i> Nägeli			x		x
<i>Coelastrum</i> sp.		x			
<i>Coenochloris hindakii</i> Komárek		x			
<i>Coenochloris</i> sp.		x			
<i>Coenococcus planctonicus</i> Korshikov		x			
<i>Coenococcus</i> sp.			x		
<i>Comassiella arcuata</i> var. <i>platydiscus</i> (G. M. Smith) Hegewald et M. Wolf				x	
<i>Crucigenia quadrata</i> Morren			xxx		
<i>Crucigenia tetrapedia</i> W. et G. S. West				x	
<i>Desmodesmus bicaudatus</i> (Dedusenko) P. M. Tsarenko		x			

Taxa	K-2000- 2002	K-2006	K-2010- 2012	D-1972 1975	D-2010- 2012
<i>Dictyosphaerium simplex</i> Korshikov			x		
<i>Dictyosphaerium</i> sp.	x				
<i>Elakatothrix gelatinosa</i> Wille			x		x
<i>Elakatothrix genevensis</i> (Reverdin) Hindak					x
<i>Elakatothrix spirochroma</i> (Reverd.) Hind					x
<i>Ennalax acutiformis</i> (Schöder) Hindák					xx
<i>Eudorina elegans</i> Ehrenberg	x	xxx	x	x	xxx/x
<i>Golenkinia radiata</i> Chodat			x		x
<i>Hariotina reticulata</i> P. A. Dangeard			x		
<i>Koliella</i> sp.					x
<i>Lagerheimia ciliata</i> (Lagerheim) Chodat			xxx		
<i>Lagerheimia longiseta</i> Lemmermann) Printz			x		
<i>Micractinium pusillum</i> Fresenius			x		x
<i>Monactinus simplex</i> (Meyen) Corda	x	xxx/x	x		x
<i>Monoraphidium pusillum</i>		x			
<i>Monoraphidium</i> sp.	x				
<i>Mucidosphaerium pulchellum</i> (H.C. Wood) C. Bock, Proschold et Krienitz		xxx/x	x	x	x
<i>Mychonastes elegans</i> (Bachmann) Krienitz, C. Bock, Dadlech et Proschold		x			
<i>Oocystidium ovale</i> Korshikov			xx		xxx/x
<i>Oocystis borgei</i> J.Snow		x	x		x
<i>Oocystis lacustris</i> Chodat		xxx/x	xx		
<i>Oocystis socialis</i> Ostenfeldt		x			
<i>Oocystis</i> sp.				x	
<i>Oedogonium</i> sp.					x
<i>Pandorina morum</i> (O.F.Müller) Bory de Saint-Vincent		x	xx/x		xxx/x
<i>Pediastrum duplex</i> Meyen	x	xxx/x	x		x
<i>Pseudopediastrum boryanum</i> (Turpin) Hegewald in Buchheim et al.	x		x		
<i>Pseudoschroederia robusta</i> (Korshikov) Hegewald et E.Schnepf		x	x		
<i>Pseudostaurastrum limneticum</i> (Borge) Couté et G. Rousselin			x		
<i>Quadricoccus ellipticus</i> Hortobágyi		x			

Taxa	K-2000- 2002	K-2006	K-2010- 2012	D-1972 1975	D-2010- 2012
<i>Scenedesmus acuminatus</i> (Lagerheim) Chodat incl. var. <i>biseriatus</i> <i>Reinsch sensu</i> Kors.	x			x	x
<i>Scenedesmus acuminatus</i> var. <i>elongatus</i> G. M. Smith			x		x
<i>Scenedesmus communis</i> (Breb.) Hegewald	x		x	x	x
<i>Scenedesmus intermedius</i> var. <i>bicaudata</i> Hortobágyi			x		
<i>Scenedesmus obliquus</i> (Turpin) Kützing			xxx		x
<i>Scenedesmus obtusus</i> Meyen			x		
<i>Scenedesmus</i> sp.	x				
<i>Schroederia setigera</i> (Schröder) Lemmermann		xxx/x	x		
<i>Schroederia spiralis</i> (Printz.) Korshikov		x	x		
<i>Schroederia</i> sp.	x				
<i>Selenastrum bibraianum</i> Reinsch			x		
<i>Tetrachlorella</i> sp.			x		
<i>Tetradesmus cumbricus</i> G. S. West		x			
<i>Tetrastrum glabrum</i> (Y. V. Roll) Ahlstrom et Tiffany			x		x
<i>Tetrastrum</i> sp.			x		
<i>Treubaria planctonica</i> (G.M.Smith) Korshikov			x		
<i>Trochiscia planctonica</i> E. M. Lind et Pearsall		x			
<i>Volvox aureus</i> Ehrenberg		x	x		
<i>Willea irregularis</i> (Wille) Schmidle			x		x
Streptophyta					
<i>Arthrodesmus</i> sp.	x				
<i>Closterium aciculare</i> T. West	x	x	xxx/x		
<i>Closterium acutum</i> Brébisson		x	x		
<i>Closterium baillyanum</i> (Brébisson ex Ralfs) Brébisson				x	x
<i>Closterium moniliferum</i> Ehrenberg ex Ralfs			x		
<i>Closterium pronum</i> Brébisson			x		
<i>Closterium venus</i> Kützing ex Ralfs		x	x		
<i>Closterium</i> spp.	x	x			

Taxa	K-2000- 2002	K-2006	K-2010- 2012	D-1972 1975	D-2010- 2012
<i>Cosmarium depressum</i> Bailey					x
<i>Cosmarium margaritiferum</i> Meneghini ex Ralfs			x		
<i>Cosmarium pseudoholmii</i> Borge					x
<i>Cosmarium</i> spp.	x		x	xxx/x	x
<i>Gonatozygon pilosum</i> Wolle				x	x
<i>Gonatozygon monotaenium</i> De Bary	x	x	x		
<i>Mougeotia</i> sp.				x	
<i>Spirogyra</i> sp.				x	
<i>Staurastrum apiculatum</i> Brébisson				x	
<i>Staurastrum depressum</i> (Nägeli)			x		
<i>Staurastrum dickiei</i> Ralfs				x	
<i>Staurastrum gracile</i> Ralfs.	x	xxx/x	x	xxx/x	x
<i>Staurastrum pinque</i> Teiling			xxx/x		x
<i>Staurastrum planctonicum</i> Teiling			x		x
<i>Staurastrum</i> sp.	x		x		
Euglenophyta					
<i>Euglena clara</i> Skuja		x	x		
<i>Euglena granulata</i> (Klebs) F. Schmitz			x		x
<i>Euglena polymorpha</i> f. <i>minor</i> T. Hortobágyi					x
<i>Euglena</i> spp.	x	x	x		
<i>Lepocynclis oxyuris</i> (Schmarda) Marin et Melkonian in Marin et al.					x
<i>Lepocynclis acus</i> (O. F. Müller) Marin et Melkonian in Marin et al.			x		x
<i>Phacus longicauda</i> (Ehrenberg) Dujardin					x
<i>Phacus orbicularis</i> K. Hübner			x		x
<i>Phacus</i> sp.	x				
<i>Strombomonas</i> sp.					x
<i>Trachelomonas hispida</i> (Perty) F. Stein emend. Deflandre				x	x
<i>Trachelomonas planctonica</i> Svirenko			x		
<i>Trachelomonas rotunda</i> Svirenko		xxx/x	x		
<i>Trachelomona volvocina</i> Ehrenberg				x	
<i>Trachelomonas</i> sp.	x	x			
Pyrrhophyta			x		
<i>Ceratium cornutum</i> (Ehrenberg) Claparède et J. Lachmann					x

Taxa	K-2000- 2002	K-2006	K-2010- 2012	D-1972 1975	D-2010- 2012
<i>Ceratium hirundinella</i> (O.F. Muller) Dujardin	x	x	x	x	
<i>Glenodinium</i> sp.					x
<i>Gymnodinium uberrimum</i> (G.J.Allman) Kofoid et Swezy					xxx/xx/x
<i>Gyrodinium helveticum</i> (Penard) Y. Takano et T. Horiguchi					x
<i>Peridinium</i> spp.	x	x	x	xxx/x	
Cryptophyta					
<i>Cryptomonas caudata</i> J.Schiller					x
<i>Cryptomonas erosa</i> Ehrenberg					x
<i>Cryptomonas ovata</i> Ehrenberg					x
<i>Cryptomonas</i> spp.	x			x	
Ochromophyta			x		
Chrysophyceae					
<i>Dinobryon divergens</i> O.E.Imhof				x	x
<i>Dinobryon</i> sp.	x				
<i>Uroglena volvox</i> Ehrenb.	x				
Synurophyceae			x		
<i>Mallomonas akrokomos</i> Ruttner in Pascher					x
<i>Mallomonas caudata</i> Iwanoff			x		x
<i>Mallomonas elongata</i> Reverdin			x		x
<i>Mallomonas</i> spp.			x	x	
<i>Synura uvella</i> Ehrenberg				x	
Bacillariophyceae					
<i>Achnanthes</i> sp.			x		x
<i>Amphora ovalis</i> (Kützing) Kützing			x		x
<i>Asterionella formosa</i> Hassal		x	xxx/xx/x	xxx/x	xxx/xx/x
<i>Asterionella gracillima</i> (Hantzsch) Heib.				xxx/x	
<i>Asterionella ralfsii</i> W. Smith			x		x
<i>Asterionella</i> sp.	x				
<i>Aulacoseira ambigua</i> (Grunow) Simonsen					x
<i>Aulacoseira granulata</i> (Ehrenberg) Simonsen	x	x	xxx/xx/x	x	xx/x
<i>Aulacoseira granulata</i> var. <i>angustissima</i> (O. F. Müller) Simonsen				xxx	

Taxa	K-2000- 2002	K-2006	K-2010- 2012	D-1972 1975	D-2010- 2012
<i>Aulacoseira islandica</i> O. Müll			x		x
<i>Aulacoseira italica</i> (Ehrenberg) Simonsen			x		x
<i>Aulacoseira italica</i> var. <i>tennuissima</i> (Grunow) Simonsen			x		
<i>Bacillaria</i> sp.					x
<i>Caloneis amphisbaena</i> (Bory de Saint Vincent) Cleve			x		x
<i>Caloneis silicula</i> (Ehrenberg) Cleve					x
<i>Ceratoneis arcus</i> (Ehrenberg) Kützing				x	
<i>Cocconeis pediculus</i> Ehrenberg			x		
<i>Cocconeis placentula</i> Ehrenberg			x		x
<i>Cocconeis</i> spp.			x	x	x
<i>Cyclotella meneghiniana</i> Kützing			xxx/xx/x		xxx/x
<i>Cyclotella</i> spp.	x			x	
<i>Cymatopleura solea</i> (Brébisson) W. Smith			x	x	x
<i>Cymbella cistula</i> (Hemprich & Ehrenberg) O. Kirchner					x
<i>Cymbella cymbiformis</i> C. Agardh			xx/x		x
<i>Cymbella helvetica</i> Kützing			x		
<i>Cymbella lanceolata</i> (C. Agardh) van Heurck			x		
<i>Cymbella tumida</i> (Brébisson) van Heurck			x		
<i>Cymbella ventricosa</i> C. Agardh					x
<i>Cymbella</i> spp.	x	x		x	
<i>Diatoma ehrenbergii</i> Kützing					x
<i>Diatoma elongatum</i> (Lyngb.) Ag.			x		
<i>Diatoma hyemalis</i> (Roth) Heiberg			x		x
<i>Diatoma vulgare</i> Bory			x	x	x
<i>Diploneis ovalis</i> (Hilse) Cleve					x
<i>Diploneis</i> sp.			x		
<i>Epithemia</i> sp.			x		x
<i>Eunotia bilunaris</i> (Ehrenberg) Schaarschmidt					x
<i>Eunotia minor</i> (Kützing) Grunow					x
<i>Eunotia soleirolii</i> (Kützing) Rabenhorst					x
<i>Eunotia</i> sp.	x				

Taxa	K-2000- 2002	K-2006	K-2010- 2012	D-1972 1975	D-2010- 2012
<i>Fragilaria capucina</i> Desmazières		x		x	x
<i>Fragilaria crotonensis</i> Kitton	xxx	xxx	xx/x	xxx/x	xxx/xx/x
<i>Fragilaria</i> sp.	x				
<i>Frustulia rhomboides</i> (Ehrenberg) De Toni					x
<i>Gomphonema acuminatum</i> Ehrenberg					x
<i>Gomphonema acuminatum</i> var. <i>coronatum</i> (Ehrenberg) Ehrenberg					x
<i>Gomphonema constrictum</i> Ehrenberg			x	x	x
<i>Gomphonema gracile</i> Ehrenberg			x		
<i>Gomphonema lagerheimii</i> A.Cleve					x
<i>Gomphonema truncatum</i> Ehrenberg					x
<i>Gomphonema</i> spp.	x		x		
<i>Gyrosigma acuminatum</i> (Kützing)					x
<i>Gyrosigma</i> sp.	x				
<i>Hannaea arcus</i> (Ehrbg.) Patrick	x		xx/x		x
<i>Melosira varians</i> C. A. Agardh		xxx	xxx/xx/x	x	x
<i>Melosira</i> sp.		x			
<i>Meridion circulare</i> (Greville) C.Agardh			x		x
<i>Navicula gracilis</i> Ehrenberg			x		
<i>Navicula lanceolata</i> Ehrenberg			x		x
<i>Navicula radiosa</i> Kützing			x		x
<i>Navicula reinhardtii</i> (Grunow) Grunow					x
<i>Navicula rhynchocephala</i> Kützing			x		x
<i>Navicula vulpina</i> Kützing					x
<i>Navicula</i> spp.	x		x	x	
<i>Nitzschia closterium</i> (Ehrenberg) W. Smith			x		
<i>Nitzschia graciliformis</i> Lange-Bertalot & Simonsen			x		
<i>Nitzschia</i> sp.			x		x
<i>Pinnularia borealis</i> Ehrenberg					x
<i>Pinnularia</i> spp.	x		x	x	
<i>Pleurosigma elongatum</i> W. Smith			x		
<i>Rhopalodia gibba</i> (Ehrenberg) O. Müller					x
<i>Stephanodiscus astraea</i> (Ehrenberg) Grunow			x		xx/x
<i>Stephanodiscus hantzschii</i> Grunow		x	x		x
<i>Stephanodiscus</i> sp.	x				

Taxa	K-2000-2002	K-2006	K-2010-2012	D-1972-1975	D-2010-2012
<i>Surirella</i> sp.					x
<i>Synedra acus</i> Kützing	x		x	xxx/x	x
<i>Synedra acus</i> var. <i>radians</i> Kützing				xxx/x	
<i>Synedra famelica</i> Kützing			x		
<i>Synedra pulchella</i> (Ralfs) Kützing					x
<i>Synedra rumpens</i> Kützing					x
<i>Synedra tabulata</i> (C. Agardh) Kützing			x		
<i>Synedra ulna</i> (Nitzsch.) Ehrenberg	x	x	xxx/xx/x	x	x
<i>Synedra ulna</i> var. <i>biceps</i> (Kützing) Schönfeldt					x
<i>Synedra vaucheriae</i> (Kützing) Kützing			x		
<i>Tabellaria fenestrata</i> (Lyngbye) Kützing					xxx/xx/x
<i>Tabellaria fenestrata</i> var. <i>asterionelloides</i> Grunow in Van Heurck				x	x
<i>Tabellaria fenestrata</i> var. <i>intermedia</i> Grunow				x	
<i>Tabellaria flocculosa</i> (Roth.) Kützing				x	xxx/x

Acknowledgements: The study was carried out in a framework of the project DO 02-307/19.12.2008 “Complex investigation and assessment of the impact of fish cage culture on the hydro-ecosystem in some Bulgarian dam-lakes” funded by the National Science Fund, Ministry of Education, Youth and Science of Bulgaria.

4. References:

- ARZUL G., CLÉMENT A., PINIER A., 1996: Effects on phytoplankton growth of dissolved substances produced by fish farming. *Aquat. Living Resour.* 9: 95-102.
- BARTOZEK E. C. R., BUENO N. C., RODRIGUES L. C., 2014: Influence of fish farming in net cages on phytoplankton structure: a case study in a subtropical Brazilian reservoir. *Braz. J. Biol.* 74 (1). DOI: <http://dx.doi.org/10.1590/1519-6984.21912>
- BELKINOVA D., MLADENOV R., DIMITROVA-DYULGEROVA I., CHESHMEDJIEV S., ANGELOVA I., 2007: Kurdjali Reservoir phytoplankton research. *Phytologia Balcanica* 13: 47-52.
- BORGES P. A. F., TRAIN S., RODRIGUES L. C., 2008: Estrutura do fitoplâncton, em um curto período de tempo, em um braço do reservatório de Rosana (ribeirão do Corvo, Paraná, Brasil). *Acta Scientiarum Biological Science* 30 (1): 57-65.
- CAO L., WANG W., YANG Y., YANG C., YUAN Z., XIONG S., DIANA J., 2007: Environmental Impact of Aquaculture and Countermeasures to Aquaculture Pollution in China. *Env. Sci. Pollut. Res.* 14 (7): 452–462.
- COESEL P. F. M., 1997: De Desmidiaceen van Nederland - Sieralgen - Deel 6 Fam. Desmidiaceae (4). *Wetenschappelijke Mededelingen van de KNNV* 220: 1-93.

- CROSSETTI L.O., BICUDO D.C., BICUDO C. E .M., BINI, L. M., 2008: Phytoplankton biodiversity changes in a shallow tropical reservoir during the hypereutrophication process. *BRAZ. J. BIOL.* 68 (4): 1061-1067. DOI: <http://dx.doi.org/10.1590/S1519-69842008000500013>.
- DEGEFU F., MENGISTU S., SCHAGERL M., 2011: Influence of fish cage farming on water quality and plankton in fish ponds: A case study in the Rift Valley and North Shoa reservoirs, Ethiopia. *Aquaculture* 316: 129–135.
- DEMIR N., KIRKAGAC M. U., PULATSÜ S., BEKCAN S., 2001: Influence of trout cage culture on water quality, plankton and benthos in an Anatolian Dam lake. *The Israeli Journal of Aquaculture – Bamidgeh* 53 (3-4): 115-127.
- DIAZ M. M., TEMPORETTI P. F., PEDROSO F. L., 2001: Response of phytoplankton to enrichment from Cage fish farm waste in Alicura reservoir (Patagônia, Argentina). *Lakes and Reservoirs: Research and Management* 6: 151-158.
- DOCHIN T., STOYNEVA M. P.: Phytoplankton of the reservoir “Dospat” (Rodopi Mts, Bulgaria) as indicator of negative trend in reservoir development due to long-term cage fish farming. *Ann. Univ. Sof., Fac. Biol., Book 2-Botany*, in press.
- GUIRY M. D., GUIRY G. M., 2014. *AlgaeBase*. World-wide electronic publication, National University of Ireland, Galway. <http://www.algaebase.org>.
- HADJINIKOLOVA L., ILIEV I., 2012: Seasonal and vertical variations of water temperature and oxygen content in the Dospat reservoir, Bulgaria, Conf. Proceedings V Internat. Conf. „Aquaculture and Fishery“ 01-03 June, 2011, Belgrad: 457-463.
- ILIEV I., HADJINIKOLOVA L., 2011: Seasonal and vertical dynamics of the water temperature and oxygen content in Kurjali reservoir, Bulgaria. *Agricultural Science and Technology* 5 (2): 212-215.
- KARAKASSIS J., PITTA P., KROM M. D., 2005: Contribution of fish farming to the nutrient load of the Mediterranean. *Scientia Marina* 69 (2): 313-321.
- KOMÁREK J., ANAGNOSTIDIS K., 1999. Cyanoprokaryota. 1. Teil: Chroococcales. In: ETTL H., GÄRTNER G., HEYNIG H., MOLLENHAUER D. (Eds.): *Süßwasserflora von Mitteleuropa*, 19 (1), G. Fischer, Jena, Stuttgart, Lübeck, Ulm, 549 pp.
- LENZENWEGER R., 1997: Desmidiaceenflora von Österreich. Teil 2. *Bibl. Phycol.* 102: 1-216.
- MICHEV T. M., STOYNEVA M. P. (Eds.), 2007: *Inventory of Bulgarian Wetlands and their Biodiversity. Part 1: Non-Lotic Wetlands*. Publ. House Elsi-M, Sofia, 364 pp. + CD supplement.
- NAIDENOW W., SAIZ D., 1977: Das Plankton im Stauraum “Dospat” in den ersten Jahren nach der Aufspeicherung. *Hydrobiologia* (Sofia) 5: 24- 37 (In Bulgarian, Russian and German summ.).
- NAUMOVA S., ZHIVKOV M., 1988: The influence of the intensive trout cage farming and hydrochemical and hydromicrobiological quality of the water in the reservoir “Dospat”. *Hydrobiologia* (Sofia) 33: 45-57 (In Bulgarian).
- NOGUEIRA M. G., 2000: Phytoplankton composition, dominance and abundance as indicators of environmental compartmentalization in Jurumirim Reservoir (Paranapanema River), São Paulo, Brazil. *Hydrobiologia* 431: 115-128.
- PALAMAR-MORDVINTZEVA G. M., 1982: Chlorophyta: Conjugatophyceae, Desmidiales (2). In: GOLLERBAK, M. M., (Ed.): *Manual of freshwater algae of USSR*, Vol. 11 (2), Leningrad, Nauka, 620 pp. (In Russian).
- POPOVSKÝ J., PFIESTER A. L., 1990: Dinophyceae Dinoflagellida. In: ETTL H., GERLOFF J., HEYNIG H., MOLLENHAUER D. (Hrgs.): *Süßwasserflora von Mitteleuropa*. Band 6. Stuttgart, Jena, Gustav Fisher, 272 pp.

- STARMACH K., 1974: Cryptophyceae – Kryptofity. Dinophyceae – Dinofity. Raphidophyceae – Rafidofity. In: STARMACH K., SIEMIŃSKA J. (Eds): Flora Ślōdkowodna Polski. Tom 4. Warszawa – Kraków, PWN, 520 pp.
- STOYNEVA M., 1998. Algae. In: MICHEV T. M., GEORGIE B. B., PETROVA A. V., STOYNEVA M. P., (Eds.): 1998: Biodiversity of the Srebarna Biosphere Reserve. Checklist and bibliography, Sofia, Co-publ. Context & Pensoft: 10-37.
- STOYNEVA M. P., 2000a: Algological studies of Bulgarian coastal wetlands. I. Species composition of the phytoplankton of Durankulak and Shabla-Ezeretz lakes. Ann. Univ. Sof. 91 (2): 27-48.
- STOYNEVA M. P., 2000b: Planktic green algae of Bulgarian coastal wetlands. Hydrobiologia 438: 25-41.
- STOYNEVA M. P., 2014: Contribution to the studies on biodiversity of aero- and hydrobiontic pro- and eukaryotic algae in Bulgaria. DrSc Thesis, Sofia University “St Kliment Ohridski”, Faculty of Biology, Department of Botany, 825 pp. (In Bulgarian).
- STOYNEVA M. P., TEMNISKOVA-TOPALOVA D. N., 2007: Cyanoprokaryotes and of Bulgarian Non-Lotic Wetlands and their Biodiversity. In: MICHEV T. M., STOYNEVA M. P. (Eds.): Inventory of Bulgarian Wetlands and their Biodiversity. Part 1: Non-Lotic Wetlands. Publ. House Elsi-M, Sofia: 155-167.
- THANH HUNG L., THE NHAN D., YI Y., DIANA J. S., KWEI LIN C., 2005: Mitigating environmental impact of cage culture through integrated cage-cum-cove culture system in Tri An reservoir of Vietnam. In: BURRIGHT J., FLEMMING C., EGNA H. (Eds.): Twenty-Second Annual Technical Report. Aquaculture CRSP, Oregon State University, Corvallis, Oregon: 132-145.
- TRAYKOV I., 2005: Factors influencing the trophic state of the reservoir Kurdzhali. PhD Thesis, Sofia University “St Kliment Ohridski”, Faculty of Biology, 188 pp. (In Bulgarian).
- WILLIAMS P., WHITFIELD M., BIGGS J., BRAY S., FOX G., NICOLET P., SEAR D., 2004: Comparative biodiversity of rivers, streams, ditches and ponds in an agricultural landscape in Southern England. Biol. Cons. 115: 329–341.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte des naturwissenschaftlichen-medizinischen Verein Innsbruck](#)

Jahr/Year: 2014

Band/Volume: [99](#)

Autor(en)/Author(s): Dochin Kostadin T., Stoyneva Maya P.

Artikel/Article: [Effect of long-term cage fish-farming on the phytoplankton biodiversity in two large Bulgarian reservoirs 49-66](#)