

BRAUERIA (Lunz am See, Austria) 20: 7-9 (1993)
VERTICAL DISTRIBUTION OF CADDIS LARVAE IN
VARIOUS TYPES OF LAKE LITTORAL

Stanisław Czachorowski

This contribution is supplementary to papers: Czachorowski 1989 and Czachorowski in prep. General model of habitat distribution of Trichoptera in lakes was presented during the 7th International Symposium on Trichoptera. Detailed data of this problem will be published shortly. Investigations are now being continued in the Lobelian lakes (The Pomerania Lakeland, North-West Poland).

Caddis larvae were collected during three years (1988-1990) at monthly intervals in three lakes: Narckie, Warchafdzkie and Brajnckie (Northern Poland). Vertical distribution of caddis larvae were investigated at seven stations (various types of littoral, see Bernatowicz & Zachwieja 1966).

Fig.1. Narckie Lake (mesotrophic), littoral with great-lake helophytes, elodeids distant from helophytes. Species typical for submergent plant (Magnocaracetum, Ceratophyllum, Myriophyllum) do not occur in helophytes (group C2: *Cyrnus flavidus*, *C. crenaticornis*, *Oxyethira* sp.) or occur in small numbers (group C1: *Leptocerus tineiformis*, *Mystacides longicornis*, *Athripsodes aterrimus*). Limnephilidae occur only in helophytes.




FIG. 1

Abbreviations in the figures:

- A (A1, A2) - species occurring only in the most shallow littoral (helophytes zone).
 - B (B1, B2) - species occurring throughout the vertical range.
 - C (C1, C2, C3) - species occurring only or mainly in elodeids zone.
- Mean density: mean density of caddis larvae to 0,3 - 0,5 m² of bottom.

Fig.2. Narckie Lake, small-lake phytolittoral, elodeids directly adjoin helophytes. Special typical for large under-water meadows occur in helophyte zone in great numbers and species typical for helophytes (rushes and bulrush zone) in deep water.




FIG. 2

Fig.3. Narckie Lake, atrophic phytolittoral, muddy bottom, elodeids represented mainly by *Stratiotes aloides*, *Characeae*, *Elodea canadensis*. *Limnephilus griseus* occur in temporary pools. Species of elodeids occur very near-by marsh helophytes.




FIG. 3

Fig.4. Warchafdzkie Lake (eutrophic), small-lake phytolittoral.
Fig.5. Warchafdzkie Lake, pond phytolittoral.




FIG. 4




FIG. 5

Fig.6 and 7. Brajnicksie Lake (hypertrophic), artificially changed phytolittoral. Submergent plants occur very deeply (depth 0,2 - 0,6 m) and represented mainly by Potamogeton. Species typical for perennial elodeids do not occur.

The presented data show that trophy of lake and type of habitat are of importance for vertical distribution of Trichoptera. However, spatial and vertical configuration of littoral and connected with them migrations of larvae between adjoining habitat modify this vertical distribution. The intensity of migrations is directly proportional to the distance between habitats.




FIG. 6




FIG. 7

Marks of species used in fig. 1 - 7:

Aa - *Athripsodes aterrimus* (Steph.), Ac - *A.cinereus* (Curt.), Ag - *Agraylea multipunctata* Curt., Al - *Anabolia laevis* (Zett.), Ca - *Ceraclea annulicornis* (Steph.), Cc - *Cyrnus crenaticornis* (Kol.), Cf - *C.flavidus* McL., Ci - *C.insolutus* McL., Cs - *C.sp.juv.*, Eb - *Erotesis baltica* McL., Et - *Ecnomus tenellus* (Ramb.), Hd - *Holocentropus dubius* (Ramb.), Hp - *H.picicornis* (Steph.), Hy - *Hydroptila* sp., La - *Limnephilus auricula* Curt., Lb - *L.borealis* (Zett.), Lc - *L.fuscicornis* Ramb., Ld - *L.decepiens* (Kol.), Le - *Leptocerus tineiformis* (Curt.), Lf - *Limnephilus flavicornis* (Fabr.), Lg - *L.griseus* (L.), Li - *L.lunatus* Curt., Lm - *L.marmoratus* Curt., Ln - *L.nigriceps* (Zett.), Lp - *L.politus* McL., Lr - *L.rombicus* (L.), Ls - *L.sp.juv.*, Lu - *L.sparsus* (Curt.), Ly - *Lype* sp., Ml - *Mystacides longicornis* (L.), Mn - *M.nigra* (L.), Mo - *Molanna angustata* Curt., Ms - *Mystacides* sp.juv., Ne - *Nemotaulius punctatolineatus* (Retz.), Of - *Oecetis furva* (Ramb.), Ol - *O.lacustris* (Pict.), Oo - *O.ochracea* (Curt.), Or - *Orthotrichia* sp., Os - *Oecetis* sp.juv., Ox - *Oxyethira* sp., Pb - *Phryganea bipunctata* Retz., Pg - *P.grandis* L., Ps - *P.sp.juv.*, Tb - *Triaenodes bicolor* (Curt.), Ts - *Tinodes* sp., Tw - *T.waeneri* (L.).

REFERENCES

Bernatowicz, S., Zachwieja, J., 1966, Types of littoral found in the lakes of the Masurian and Suwalki Lakelands. - *Ekol.pol.* 14:519-545.

Czachorowski, S., 1989, Vertical distribution of Trichoptera in three Masurian lakes - results of preliminary studies. - *Pol.Arch.Hydrobiol.* 36:351-358.

Czachorowski, S. (in prep.), Vertical distribution of caddis larvae in lakes. - *Pol.Arch.Hydrobiol.*

Author's address:
Stanisław Czachorowski
Teacher's Training College
Dept. of Ecology, Żołnierska 14
PL - 10-561 Olsztyn, Poland



MEETINGS

5th European Congress of Entomology,
28.8.-2.9.1994:
Institute of Applied Biology, University of
York, York YO1 5DD, U.K.

14. Internationales Symposium über
Entomofaunistik in Mitteleuropa, München
(BRD), 5.-10.9.1994
Information: Dr. Gerhard SCHERER, Zoologische
Staatsammlung, Münchenhausenstraße 21,
D - 8000 München 60, BRD.

8th International Symposium on Trichoptera
Minneapolis/St. Paul, Minnesota, USA,
9 - 15 July 1995
Information: Ralph Holzenthal,
University of Minnesota, Department of
Entomology, 219 Hodson Hall,
1980 Folwell Ave., St. Paul MN 55108-6125

20th International Congress of Entomology
Florence, Italy, August 25 - 31, 1996
Information: Organizing Secretariat,
Via A. La Marmora 24, I - 50121 Firenze

BRAUERIA (Lunz am See, Austria) 20: 9 (1993)

A.G.McFARLANE: Obituary

A direct link with the 19th century was broken on 24 August 1992 when Alexander Grant McFarlane, who spent most of his life working on the systematics and life histories of the New Zealand Trichoptera, died after a short illness. Alex was born in Takapau, North Island of New Zealand on 9 December 1899. He graduated as a teacher from Wellington Training College in 1923; whilst teaching, he studied at Canterbury University part-time. His chosen thesis topic was the biology of the Rhyacophilidae.

Alex completed his M.A. degree in Zoology with First Class Honours at Canterbury University in 1937; he continued to work on Trichoptera part-time during his career as a school teacher. After retirement he came to Canterbury Museum, where he was appointed Technical Assistant in 1964 and Research Associate in 1965. He retired from the Museum in 1987. By this time he had named and described about (1/3) of the known New Zealand caddisfly fauna; he also studied and described the immature stages of many of the species. His 1951 paper on the larval Rhyacophilidae (now Hydrobiosidae) remains the standard work on the group.

Alex was a keen fisherman and lover of the outdoors all of his life. He is remembered with special affection by all who worked with him as a quiet, unassuming man, with a great interest in people.

A.G.McFarlane: Bibliography

- 1939. Additions to the New Zealand Rhyacophilidae. Part 1. - *Trans.R.Soc.N.Z.* 69:330-340.
- 1951a. A note on the genus *Neurochorema* Till. - *Rec.Canterbury Mus.* 5:253-254.
- 1951b. Additions to the New Zealand Rhyacophilidae. Part 2. - *Rec.Canterbury Mus.* 5:255-265.
- 1951c. Caddis Fly Larvae (Trichoptera) of the family Rhyacophilidae. - *Rec.Canterbury Museum* 5:267-289.
- 1956. Additions to the New Zealand Trichoptera (Part 3). - *Rec.Canterbury Mus.* 7:29-41.
- 1960. Additions to the New Zealand Trichoptera (Part 4). - *Rec.Canterbury Museum* 7:203-218.
- 1964. A new endemic subfamily, and other additions and emendations to the Trichoptera of New Zealand (Part 5). - *Rec.Canterbury Mus.* 8:55-79.
- 1966. New Zealand Trichoptera (Part 6). - *Rec.Canterbury Mus.* 8:137-161.
- 1969. Trichoptera (Caddisflies). In: G.A.Knox (ed.), *The Natural History of Canterbury*, A.H. & A.W.Reed, pp.481-483.
- 1973. Five new species of Trichoptera from New Zealand. - *J.Roy.Soc.New Zealand* 3:23-34.
- 1976. A generic revision of the New Zealand Hydropsychinae (Trichoptera). - *J.Roy.Soc.New Zealand* 6:23-35.
- 1977. Light trapping of caddisflies at Winchmore Irrigation Research Station, Mid-Canterbury, New Zealand (May 1974 - September 1975) (Note). - *Mauri Ora* 5:123-124.
- & B.Cowie, 1981. Descriptions of new species and notes on some genera of New Zealand Trichoptera. - *Rec.Canterbury Mus.* 9:353-385.
- 1990. A generic key to late instar larvae of the New Zealand Trichoptera (Caddis flies). - *Rec.Canterbury Mus.* 10:25-38.
- & J.B.Ward, 1990. *Triplectidina moselyi* n.sp., a previously misidentified New Zealand caddis-fly (Trichoptera: Leptoceridae). - *NZ Entomologist* 13:55-59.

John B.Ward, Research Associate, Canterbury Museum.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Braueria](#)

Jahr/Year: 1993

Band/Volume: [20](#)

Autor(en)/Author(s): Czachorowski Stanislaw

Artikel/Article: [Vertival Distribution of Caddis Larvae in various types of Lake Littoral 7-9](#)