BRAUERIA (Lunz am See, Austria) 20:26 (1993)

International Symposium on Trichoptera, Umeå, Sweden, 3-8 August 1992.

Zoology, Uppsala University, Box 561, S-751 22 Uppsala, Sweden.

On the excursion, Wednesday 5 August 1992, along the Umeälven River and the Vindelälven River many caddisflies and larvae were collected. Five participants sent me the lists of their captures. The recorded captures of adults, including my own, are noted below.

In total 1669 specimens comprising 48 species were recorded (Tab. 1). The Leptoceridae dominated with 16 species followed by Limnephilidae with 10. As expected, the smallest number of species and specimens were captured along the regulated Umeälven river, viz. 382 specimens representing 29 species. However, the most interesting finding was 13 specimens of Oecetis notata, which were collected on an UV-lightened window of a flower shop at the center of Umea. Two males were also recorded at the Stornorrfors Power Station. O. notata is a new record of the province of Västerbotten. It has earlier been found in the province of Dalarna 400 km south of Umeå (Forsslund and Tjeder 1942), but also at Rovaniemi in Finland 400 km north of Umeå (Nybom 1960). Other new records from the province of Västerbotten include Limnephilusmarmoratus,

The caddisflies captured during the 7th Molanna albicans and Occetis testacea. All these were expected and also the new record, Athripsodes albifrons, for Lycksele Lappmark.

Along the unregulated Vindelälven River more Bo Gullefors, Section of Entomology, Department of caddisflies were collected from four different places. In total 1287 specimens belonging to 37 species were recorded.

> Athripsodes commutatus (753 inds.) was the most common species along both rivers, and most of them were netted along the bank close to the landing place of the rafts.

> The recorded number of the caddisfly species in the province of Västerbotten has now increased from 145 to 148, and in the province of Lycksele Lappmark to 123 (Gullefors 1988).

## References:

Forsslund, K.-H., and Tjeder, B. 1942. Catalogus Insectorum Sueciae. II. Trichoptera. - Opusc. Ent. 7: 92 - 107.

Gullefors, B. 1988. The Swedish Trichoptera, a check-list with records for the northern provinces. - Ent. Tidskr. 109: 71 - 80.

Nybom, O. 1960. List of Finnish Trichoptera. -Fauna fenn. 6: 1 - 56.

## Acknowledgements:

I thank Hans Malicky, Oliver Flint, Wolfram Mey, Wolfram Graf and Bronislaw Szczesny for sending me their lists of captures. I also thank Christian Otto for comments on the text.

Tab 1. No. of adult caddisflies (males and females) collected by 6 participants during the excursion along the Umeälven and Vindelälven rivers.

	Umeä River		äl	indel- ven iver		Umeälven River		Vindel- älven River	
Family/Species	Mal	Fem	Ma	Fem	Family/Species	Mal	Fem	Ma	Fem
Rhyacophilidae					Limnephilus decipiens (Kol.)	-	-	-	1
Rhyacophila nubila (Zett)	1	-	6	-	Limnephilus femoratus (Zett.)	-	-	91	101
Glossosomatidae					Limnephilus fuscinervis (Zett		-	11	19
Glossosoma nylanderi McL.	-	•	-	1	Limnephilus marmoratus Cur		-	-	1
Agapetus ochripes Curt.	-	-	-	1	Limnephilus subcentralis Bra		-	-	2
Hydroptilidae					Phacopteryx brevipennis (Cur	;.) -	-	5	4
Hydroptila angulata Mos.	1	1	-	-	Halesus radiatus (Curt.)	1	-	-	-
Hydroptila forcipata (Eat.)	-	•	1	1	Potamophylax latipennis (Cur	t.) 4	-	-	-
Hydroptila simulans Mos.	1	-	-	1	Molannidae				
Philopotamidae					Molanna albicans (Zett.)	6	1	5	2
Wormaldia subnigra McL.	-	-	14	8	Molanna angustata Cur.	3	1	1	2
Psychomyiidae					Molannodes tinctus (Zett.)	-	-	11	7
Lype phaeopa (Steph.)	1	-	3	1	Leptoceridae				
Psychomyia pusilla (F.)	-	-	1	2	Athripsodes albifrons (L.)	5	1	6	3
Tinodes waeneri (L.)	1	-	-	1	Athripsodes aterrimus (Steph.)	3	-	-	2
Polycentropodidae					Athripsodes cinereus (Curt.)	9	3	18	17
Cyrnus trimaculatus (Curt.)	9	2	118	20	Athripsodes commutatus	47	33	471	186
Holocentropus picicornis (Ste	ph.) -	-	4	1	Ceraclea alboguttata (Hag.)	-	1	-	-
Neureclipsis bimaculata (L.)	55	58	-	-	Ceraclea annulicornis (Steph.	2	3	3	2
Polycentropus flavomaculatu	s 2	1	6	2	Ceraclea dissimilis (Steph.)	8	2	-	-
Hydropsychidae					Ceraclea senilis (Burm.)	-	-	2	4
Cheumatopsyche lepida (Pict.	) -	-	1	9	Mystacides azurea (L.)	2	1	42	33
Ceratopsyche nevae (Kol.)	1	-	-	-	Mystacides nigra (L.)	-	-	4	2
Hydropsyche contubernalis	40	23	-	-	Oecetis lacustris (Pict.)	5	5	9	5
Phryganeidae					Oecetis notata (Ramb.)	11	5	-	-
Agrypnia obsoleta (Hag.)	4	1	6	-	Oecetis ochracea (Curt.)	-	-	1	-
Lepidostomatidae					Oecetis testacea (Curt.)	1	2	1	-
Lepidostoma hirtum (F.)	7	4	1	1	Triaenodes bicolor (Curt.)	-	-	_	1
Limnephilidae					Ylodes simulans (Tjed.)	1	1	-	-
Limnephilus binotatus Curt.	1	1		-	Number of specimens:	232	150	842	445
Limnephilus borealis (Zett.)	-	-	-	2	Number of species		29		37
•					-				

## **ZOBODAT - www.zobodat.at**

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Braueria

Jahr/Year: 1993

Band/Volume: 20

Autor(en)/Author(s): Gullefors Bo

Artikel/Article: The caddisflies captured during the 7th International Symposium on

Trichoptera, Umeâ, Sweden, 3 - 8 August 1992. 26