

BRAUERIA (Lunz am See, Austria) 28:15-18 (2001)

Preliminary results on the biometry of Italian Trichoptera fauna

Maria Vittoria DI GIOVANNI, Enzo GORETTI, Umberto CHIAPPAREDDO & Donatella CECCAGNOLI

Abstract A biometric study was carried out on the winged Trichoptera fauna of the entire Italian peninsula. It covered 191 taxa from five families: Rhyacophilidae (33 taxa), Glossosomatidae (11), Hydrophilidae (36), Hydropsychidae (19) and Limnephilidae (92). The specimens examined are in the "GP Moretti Collection", which, at present, represents 90% of the Italian taxa.

Key words: Trichoptera, Italy, biometric data base

A biometric investigation was carried out on the winged Trichoptera fauna of the entire Italian peninsula with the aim of providing an analytical inventory of measurements of the 365 species and 30 subspecies. The specimens studied were those of the "GP Moretti Collection" which includes 90% of known Italian taxa (MORETTI et al. 1997). In this study 191 taxa from five families of the three main taxonomic groups were studied (WIGGINS & WICHARD 1989; WICHARD 1991; WICHARD et al. 1997): *ANNULIPALPIA*-Hydropsychidae (19 taxa); *SPICIPALPIA*-Glossosomatidae (11), Hydroptilidae (36) and Rhyacophilidae (33); *INTEGRIPALPIA*-Limnephilidae (92).

A maximum of six individuals, male and female, from each taxon were examined for their northern, central and southern distribution and the following morphological parameters recorded: head width and lengths of forewing, head-forewing, head-abdomen, metafemur, maxillary palp, labial palp (Fig. 1).

Table 1 reports the mean values for these parameters for the specimens of each taxon examined and, in addition, the more significant basic statistical data for the five families considered. At times, because of the limited number of individuals for each species studied, only data on biometric indications, and not on population variability, could be obtained. Forewing length is the parameter most often used for identifying the dimensional interval of species (MALICKY 1983; NEBOISS 1986). On this basis, all taxa with a forewing measurements of less than about 4 mm belong to the Italian microtrichoptera group, which, with the exception of *Agraylea* and *Allotrichia*, are represented only by the Hydroptilidae genera (Fig. 2).

The dimensional differences between male and female forewings are given for each taxon in Fig. 3, but, naturally, only the taxa where couples of examples originating from the same geographic area were available.

Of the seven parameters examined, the dimensional relationship between ♂/♀ is a measure of sexual dimorphism. Comparison of the five families examined reveals that the mean ratio for the morphometric parameters is 0.96.

On the basis of the information from the five families examined, it is our intention to continue the investigation and extend it to all Italian Trichoptera. We consider the biometric data base of the species an useful body of knowledge for characterizing both populations and their interactions.

Acknowledgement

We should like to express gratitude to our teacher Prof. Giampaolo Moretti.

References

MALICKY, H., 1983, Atlas of European Trichoptera. Junk, The Hague.

MORETTI, G.P., DI GIOVANNI, M.V., GIANOTTI, F.S., GORETTI, E., CHIAPPAREDDO, U., 1997, I Tricotteri italiani della collezione Giampaolo Moretti. Catalogo - Aggiornamento al 31/12/1996. Riv. Idrobiol. 36:1-437.

NEBOISS, A., 1986, Atlas of the Trichoptera of the SW Pacific-Australian Region. Junk, Dordrecht.

WICHARD, W., 1991, The evolutionary effect of overcoming osmosis in Trichoptera. Proceedings of the 6th International Symposium on Trichoptera: 337-342. Adam Mickiewicz University Press, Poznań.

WICHARD, W., KLEIN, H.P., HERTER P., 1997, Pupal cocoon of Amphimesnoptera (Lepidoptera + Trichoptera) with evolutionary considerations of the Trichoptera. Proceedings of the 8th International Symposium on Trichoptera: 475-480. Ohio Biological Survey, Columbus, Ohio.

WIGGINS, G.B., WICHARD, W., 1989, Phylogeny of pupation in Trichoptera, with proposals on the origin and higher classification of the order. - J. North Amer. Benthol. Soc. 8:260-276.

Author's address: Dipartimento di Biologia Animale ed Ecologia, Università di Perugia, I - 06123 Perugia, Italy.




Fig. 1. Drawing of the parameters measured (accurate to 0.025 mm): (1) length of anterior wing, (2) head + anterior wing, (3) head + abdomen, (4) metafemur, (5) maxillary palp, (6) labial palp, (7) width of head.

Table 1 (pages 16-17): Mean values of morphological parameters of 191 taxa investigated. Mean, standard deviation, minimum and maximum values of five Families.

N. TAXA		FOREWING (mm)	HEAD (mm)	FOREWING (mm)	HEAD (mm)	WIDTH (mm)	MAJILLARY PAIP (mm)	LABIAL PAIP (mm)	META-FEMUR (mm)	HEAD-ABOUMEN (mm)	MALES	FEMALES
RHYACOPHILIDAE												
1	<i>Rhyacophila eppamirina</i>	14.75	18.45	2.21	2.94	1.33	3.35	12.55	2	0		
2	<i>Rhyacophila equitana</i>	7.35	8.30	1.13	1.31	0.68	1.64	5.95	1	1		
3	<i>Rhyacophila arcangelina</i>	14.80	16.30	1.98	2.63	1.18	3.15	13.80	1	0		
4	<i>Rhyacophila aurata</i>	12.05	14.00	1.83	2.19	1.01	2.68	11.00	1	1		
5	<i>Rhyacophila dorsalis acutidens</i>	10.60	12.00	1.63	2.03	1.00	2.30	9.40	1	0		
6	<i>Rhyacophila dorsalis dorsalis</i>	11.10	13.10	1.79	2.21	0.93	2.43	11.35	1	1		
7	<i>Rhyacophila dorsalis subacutidens</i>	10.57	12.25	1.70	2.03	0.97	2.40	8.90	2	1		
8	<i>Rhyacophila fasciata</i>	9.78	11.30	1.73	1.90	0.83	2.30	8.80	1	0		
9	<i>Rhyacophila foliacea</i>	13.45	15.35	2.10	2.84	1.15	2.73	10.95	1	1		
10	<i>Rhyacophila hartigi</i>	12.05	14.15	1.90	2.26	1.00	2.68	11.00	1	1		
11	<i>Rhyacophila hirticornis orobica</i>	9.80	11.70	1.50	2.05	—	2.38	8.90	1	0		
12	<i>Rhyacophila intermedia</i>	13.90	15.85	1.85	2.44	1.28	2.75	11.40	1	1		
13	<i>Rhyacophila italice itivana</i>	11.90	14.40	2.15	2.35	1.05	2.80	13.80	1	0		
14	<i>Rhyacophila italice italice</i>	13.53	15.83	2.04	2.41	1.13	2.93	13.05	2	2		
15	<i>Rhyacophila kelnerae</i>	9.45	10.85	1.33	2.09	0.90	2.13	8.95	2	2		
16	<i>Rhyacophila laevis</i>	6.70	7.40	1.06	1.16	0.49	1.55	—	2	0		
17	<i>Rhyacophila meyeri</i>	8.73	9.98	1.24	1.69	0.74	1.88	7.75	2	2		
18	<i>Rhyacophila pallida</i>	11.70	13.20	1.80	2.44	1.08	2.70	12.55	1	1		
19	<i>Rhyacophila nubila</i>	10.50	12.20	1.80	1.95	0.93	2.50	9.30	1	0		
20	<i>Rhyacophila palmeri</i>	10.80	12.60	1.75	1.98	1.03	2.40	—	1	0		
21	<i>Rhyacophila pascoei</i>	11.05	12.50	1.91	1.94	0.86	2.45	10.20	1	1		
22	<i>Rhyacophila praemorsa</i>	13.45	15.05	1.88	2.49	1.03	2.90	—	1	1		
23	<i>Rhyacophila pubescens</i>	8.50	9.82	1.38	1.61	0.70	1.78	7.35	3	2		
24	<i>Rhyacophila ravizani</i>	14.10	16.15	1.96	2.61	1.11	2.85	12.00	1	1		
25	<i>Rhyacophila rectispina</i>	12.60	14.20	2.00	2.70	1.15	3.15	14.30	1	0		
26	<i>Rhyacophila rougemonti</i>	12.90	14.60	1.91	2.37	1.11	2.75	12.53	2	2		
27	<i>Rhyacophila simulatrix</i>	13.38	15.50	2.04	2.52	1.14	2.75	11.53	2	2		
28	<i>Rhyacophila stigmatica</i>	10.70	11.60	1.34	2.01	0.89	2.10	8.50	1	1		
29	<i>Rhyacophila tormentum</i>	15.45	17.85	2.04	2.58	1.18	3.10	12.50	1	1		
30	<i>Rhyacophila trifasciata</i>	13.55	16.20	2.03	2.13	1.00	2.80	16.00	1	1		
31	<i>Rhyacophila tristis</i>	7.46	8.60	1.15	1.59	0.79	1.63	6.98	2	3		
32	<i>Rhyacophila vellei</i>	10.20	11.40	1.78	1.70	0.74	2.33	—	1	1		
33	<i>Rhyacophila vulgaris</i>	13.18	14.90	1.96	2.46	0.97	2.73	10.93	2	2		
		MEAN	11.51	13.20	1.76	2.16	0.98	2.51	10.80			
		MAX	15.45	17.95	2.21	2.94	3.35	16.00				
		MIN	6.70	7.40	1.06	1.16	0.49	1.55	5.95			
		S.D.	2.26	2.61	0.32	0.41	0.19	0.48	2.38			
GLOSSOSOMATIDAE												
34	<i>Glossosoma bifidum</i>	7.40	8.50	1.34	1.38	0.53	1.56	6.95	1	1		
35	<i>Glossosoma bolloni</i>	6.80	7.40	1.18	1.20	0.48	1.35	5.00	1	0		
36	<i>Glossosoma conformis</i>	8.03	9.03	1.31	1.38	0.58	1.64	6.34	3	3		
37	<i>Catagapetus nigrans</i>	5.35	6.22	0.92	0.95	0.43	1.11	4.77	3	3		
38	<i>Syngapetus dubians</i>	5.10	5.65	0.83	0.81	0.36	1.05	4.65	1	1		
39	<i>Agapetus cyrensis</i>	4.00	4.75	0.75	0.61	0.29	0.98	3.65	1	1		
40	<i>Agapetus delicatulus</i>	5.20	6.15	0.85	0.80	0.41	1.08	4.50	1	1		
41	<i>Agapetus fuscipes</i>	4.05	4.80	0.69	0.51	0.25	0.76	3.70	1	1		
42	<i>Agapetus incertulus</i>	4.40	5.60	0.83	0.65	0.30	0.90	—	1	0		
43	<i>Agapetus leniger</i>	5.37	6.17	0.89	0.74	0.32	1.05	4.40	3	3		
44	<i>Agapetus nimbulus</i>	4.45	5.18	0.75	0.63	0.30	0.85	3.94	3	3		
		MEAN	5.47	6.31	0.95	0.86	0.38	1.12	4.78			
		MAX	8.03	9.03	1.34	1.38	0.56	1.64	6.95			
		MIN	4.00	4.75	0.69	0.51	0.25	0.76	3.65			
		S.D.	1.37	1.43	0.23	0.31	0.10	0.28	1.08			
HYDROPTILIDAE												
45	<i>Philocolepus granulatus</i>	3.68	4.52	0.66	0.65	0.30	0.79	3.50	3	3		
46	<i>Stactobia caspersi</i>	1.93	1.93	0.35	—	—	0.45	2.38	1	0		
47	<i>Stactobia eatonella</i>	1.63	2.10	0.39	0.40	0.21	0.40	2.33	1	1		
48	<i>Stactobia ericeae</i>	2.10	2.55	0.45	0.48	0.23	0.43	2.10	1	0		
49	<i>Stactobia furcata</i>	2.25	2.65	0.39	0.45	0.19	0.50	1.93	1	1		
50	<i>Stactobia fuscicornis</i>	1.85	2.33	0.40	0.43	0.19	0.51	2.11	2	2		
51	<i>Stactobia moseyoti</i>	2.38	2.78	0.39	0.40	0.23	0.48	2.17	3	2		
52	<i>Orthotrichia angustella</i>	2.58	3.00	0.36	0.44	0.25	0.56	2.03	1	1		
53	<i>Orthotrichia costalis</i>	2.38	2.80	0.40	0.41	0.18	0.51	2.10	1	1		
54	<i>Orthotrichia fregatii</i>	2.75	3.18	0.46	0.48	0.23	0.55	2.28	1	1		
55	<i>Oxyethira falcata</i>	3.37	3.76	0.64	0.59	0.29	0.72	3.05	3	2		
56	<i>Oxyethira flavicornis</i>	2.98	3.38	0.48	0.46	0.20	0.56	2.53	1	1		
57	<i>Oxyethira pirisnui</i>	2.40	2.80	0.58	0.53	0.23	0.53	2.40	1	0		
58	<i>Oxyethira simplex</i>	—	—	0.45	—	—	0.53	—	—	—	1	0
59	<i>Oxyethira unidentata</i>	3.05	3.60	0.48	0.53	0.25	0.63	—	—	—	1	0
60	<i>Hydroptilia aegyptia</i>	2.67	3.03	0.49	0.43	0.22	0.59	2.18	2	1		
61	<i>Hydroptilia angulata</i>	3.00	3.37	0.60	0.54	0.27	0.63	2.60	3	3		
62	<i>Hydroptilia bifurcata</i>	2.45	2.80	0.48	0.35	0.20	0.55	—	—	—	1	0
63	<i>Hydroptilia cornuta</i>	2.45	2.85	0.58	0.55	0.23	0.58	2.45	1	0		
64	<i>Hydroptilia flori</i>	3.65	4.33	0.68	0.69	0.25	0.81	3.50	1	1		
65	<i>Hydroptilia forcipata</i>	2.68	3.35	0.54	0.54	0.25	0.63	2.55	1	1		
66	<i>Hydroptilia giudicellorum</i>	2.40	2.80	0.50	0.35	0.20	0.55	—	—	—	1	0
67	<i>Hydroptilia insularica</i>	2.55	3.10	0.50	0.49	0.23	0.55	2.60	1	1		
68	<i>Hydroptilia invisa</i>	3.30	3.70	0.63	0.48	—	0.68	—	—	—	1	0
69	<i>Hydroptilia marini</i>	3.43	4.00	0.65	0.61	0.32	0.72	3.61	2	2		
70	<i>Hydroptilia occulta</i>	2.80	3.23	0.60	0.58	0.28	0.59	2.73	1	1		
71	<i>Hydroptilia rufi</i>	2.75	3.20	0.51	0.48	0.25	0.55	2.93	2	2		
72	<i>Hydroptilia similans</i>	2.75	3.20	0.55	0.49	0.24	0.55	2.20	3	2		
73	<i>Hydroptilia sparsa</i>	3.20	3.78	0.62	0.62	0.29	0.70	2.80	3	2		
74	<i>Hydroptilia stellifera</i>	3.18	3.77	0.61	0.62	0.26	0.69	2.70	2	1		
75	<i>Hydroptilia threoides</i>	3.22	3.60	0.62	0.53	0.24	0.64	2.43	3	2		
76	<i>Hydroptilia uncinata</i>	2.76	3.23	0.57	0.54	0.27	0.61	2.55	2	2		
77	<i>Hydroptilia vectis</i>	2.48	2.86	0.50	0.48	0.20	0.55	2.46	3	3		
78	<i>Agryptea sexmaculata</i>	4.05	4.55	0.78	0.53	0.25	0.83	3.20	1	0		
79	<i>Allotrichia pallicornis</i>	4.51	5.08	0.76	0.75	0.33	0.89	4.28	2	2		
80	<i>Microptilia minutissima</i>	2.20	2.53	0.43	0.38	0.20	0.48	—	—	—	1	1
		MEAN	2.79	3.24	0.53	0.51	0.24	0.60	2.62			
		MAX	4.51	5.08	0.78	0.75	0.33	0.93	4.28			
		MIN	1.63	1.93	0.35	0.35	0.18	0.40	1.93			
		S.D.	0.62	0.70	0.11	0.09	0.04	0.12	0.54			
HYDROPSYCHIDAE												
81	<i>Diplectrona atra</i>	5.95	7.15	1.13	2.06	0.60	1.38	5.05	1	1		
82	<i>Diplectrona fella</i>	7.30	8.60	1.13	2.36	0.76	1.50	5.40	1	1		
83	<i>Diplectrona magna</i>	8.65	9.85	1.43	2.39	0.76	1.96	7.25	2	2		
84	<i>Hydropsyche angustipennis</i>	9.10	10.55	1.59	2.53	0.82	1.91	7.37	3	1		
85	<i>Hydropsyche bulgermanorum</i>	8.10	9.50	1.53	2.40	0.80	1.58	—	—	—	1	0
86	<i>Hydropsyche dinarica</i>	13.50	15.80	2.03	3.75	1.20	2.25	—	—			

N. TAXA	FEMALES										MALES										
	FOREWING (mm)	HEAD (mm)	HEAD WIDTH (mm)	MAXILLARY (mm)	PLAB (mm)	LABIAL (mm)	PALP (mm)	META-FEMUR (mm)	HEAD-ABDOMEN (mm)	HEAD (mm)	FOREWING (mm)	HEAD (mm)	HEAD WIDTH (mm)	MAXILLARY (mm)	PLAB (mm)	LABIAL (mm)	PALP (mm)	META-FEMUR (mm)	HEAD-ABDOMEN (mm)	HEAD (mm)	
93 <i>Hydropsyche ometilia</i>	9.30	11.00	1.68	2.73	0.88	1.85	—	—	—	0	1	—	—	—	—	—	—	—	—	—	—
94 <i>Hydropsyche pellucidula</i>	12.80	15.00	2.08	3.78	1.20	2.32	9.92	3	3	3	3	9.92	3	3	3	3	9.92	3	3	3	3
95 <i>Hydropsyche saffleri</i>	11.45	13.45	1.84	2.96	1.05	2.23	9.85	1	1	1	1	9.85	1	1	1	1	9.85	1	1	1	1
96 <i>Hydropsyche saxonicana</i>	13.30	15.60	2.05	3.70	1.13	2.53	11.70	0	1	1	1	11.70	0	1	1	1	11.70	0	1	1	1
97 <i>Hydropsyche spiritalis</i>	9.93	11.57	1.74	3.18	1.06	1.95	8.27	3	3	3	3	8.27	3	3	3	3	8.27	3	3	3	3
98 <i>Hydropsyche tenuis</i>	11.48	13.40	1.78	3.55	1.14	2.09	9.50	3	2	2	2	9.50	3	2	2	2	9.50	3	2	2	2
99 <i>Cheumatopsyche lepida</i>	6.37	7.48	1.09	1.86	0.63	1.23	5.33	3	3	3	3	5.33	3	3	3	3	5.33	3	3	3	3
	6.81	7.52	1.66	2.90	0.95	1.97	8.24					8.24					8.24				
	MEAN	11.52	1.66	2.90	0.95	1.97	8.24					8.24					8.24				
	MAX	13.50	15.80	2.08	3.78	1.20	2.55	11.70				11.70					11.70				
	MIN	5.85	7.15	1.09	1.86	0.60	1.23	5.05				5.05					5.05				
	S.D.	2.18	2.48	0.30	0.58	0.19	0.36	1.98				1.98					1.98				
LIMNephILIDAE																					
100 <i>Apelania fimbriata</i>	6.00	7.20	1.08	0.78	0.40	1.23	5.10	1	0	0	0	5.10	1	0	0	0	5.10	1	0	0	0
101 <i>Apelania fimbriata</i>	7.45	8.90	1.21	1.10	0.45	1.51	6.45	1	1	1	1	6.45	1	1	1	1	6.45	1	1	1	1
102 <i>Drusus volscorum</i>	11.35	13.15	1.56	2.26	0.88	2.56	12.00	1	1	1	1	12.00	1	1	1	1	12.00	1	1	1	1
103 <i>Drusus apriliensis</i>	10.15	11.35	1.50	1.53	0.66	2.04	—	1	1	1	1	—	1	1	1	1	—	1	1	1	1
104 <i>Drusus biguttatus</i>	12.20	13.55	1.55	1.81	0.76	2.44	—	1	1	1	1	—	1	1	1	1	—	1	1	1	1
105 <i>Drusus cerninus</i>	9.70	11.10	1.56	1.54	0.66	2.08	8.05	1	1	1	1	8.05	1	1	1	1	8.05	1	1	1	1
106 <i>Drusus discolor</i>	11.85	13.15	1.69	1.86	0.84	2.46	9.10	1	1	1	1	9.10	1	1	1	1	9.10	1	1	1	1
107 <i>Drusus improvisus</i>	11.40	13.30	1.59	1.79	0.80	2.41	9.30	1	1	1	1	9.30	1	1	1	1	9.30	1	1	1	1
108 <i>Drusus melanchaetes</i>	10.30	11.85	1.45	1.85	0.78	2.18	8.10	1	1	1	1	8.10	1	1	1	1	8.10	1	1	1	1
109 <i>Drusus monticola</i>	12.90	14.90	1.68	1.65	0.78	2.50	—	1	0	0	0	—	1	0	0	0	—	1	0	0	0
110 <i>Drusus mtelleri</i>	13.00	15.70	1.80	2.00	1.05	2.63	12.20	1	1	1	1	12.20	1	1	1	1	12.20	1	1	1	1
111 <i>Ecdiopsyche guttulata</i>	11.50	13.05	1.77	1.80	0.93	2.45	9.43	3	3	3	3	9.43	3	3	3	3	9.43	3	3	3	3
112 <i>Ecdiopsyche malickyi</i>	13.50	15.35	1.76	2.34	1.05	2.74	—	1	1	1	1	—	1	1	1	1	—	1	1	1	1
113 <i>Cryptotritax nebulicola</i>	9.45	10.85	1.35	1.78	0.78	1.95	7.80	1	1	1	1	7.80	1	1	1	1	7.80	1	1	1	1
114 <i>Melanoea flavipennis</i>	9.85	11.25	1.36	1.75	0.74	2.06	8.20	1	1	1	1	8.20	1	1	1	1	8.20	1	1	1	1
115 <i>Melanoea rhaetica</i>	8.95	10.20	1.26	1.55	0.71	1.80	7.45	1	1	1	1	7.45	1	1	1	1	7.45	1	1	1	1
116 <i>Leptodrusus budtzi</i>	10.15	12.15	1.75	1.96	0.93	2.75	12.15	1	1	1	1	12.15	1	1	1	1	12.15	1	1	1	1
117 <i>Monocentra lepidoptera</i>	10.70	12.45	1.61	1.90	0.88	2.31	9.25	1	1	1	1	9.25	1	1	1	1	9.25	1	1	1	1
118 <i>Limnephilus auricola</i>	9.10	10.90	1.34	1.46	0.63	1.75	8.75	1	1	1	1	8.75	1	1	1	1	8.75	1	1	1	1
119 <i>Limnephilus bipunctatus</i>	13.88	16.35	1.91	2.34	1.08	2.96	13.60	2	2	2	2	13.60	2	2	2	2	13.60	2	2	2	2
120 <i>Limnephilus clavicornis</i>	10.75	12.60	1.50	1.70	0.90	2.23	10.00	2	0	0	0	10.00	2	0	0	0	10.00	2	0	0	0
121 <i>Limnephilus coenosus</i>	12.70	14.55	1.73	1.98	1.03	2.58	10.40	1	1	1	1	10.40	1	1	1	1	10.40	1	1	1	1
122 <i>Limnephilus extricatus</i>	13.05	14.65	1.68	2.09	0.94	2.59	—	1	1	1	1	—	1	1	1	1	—	1	1	1	1
123 <i>Limnephilus flavicornis</i>	16.23	18.50	2.13	2.79	1.25	3.19	14.58	3	3	3	3	14.58	3	3	3	3	14.58	3	3	3	3
124 <i>Limnephilus hirsutus</i>	14.88	17.35	2.11	2.66	1.28	2.99	13.75	2	2	2	2	13.75	2	2	2	2	13.75	2	2	2	2
125 <i>Limnephilus hevelicus</i>	12.30	14.70	1.83	2.34	1.08	2.47	11.33	3	2	2	2	11.33	3	2	2	2	11.33	3	2	2	2
126 <i>Limnephilus hirsutus</i>	11.65	13.63	1.69	2.07	0.82	2.45	11.50	2	2	2	2	11.50	2	2	2	2	11.50	2	2	2	2
127 <i>Limnephilus ignavus</i>	11.42	13.35	1.52	1.92	0.83	2.26	10.68	3	3	3	3	10.68	3	3	3	3	10.68	3	3	3	3
128 <i>Limnephilus italicus</i>	11.38	13.25	1.53	1.98	0.88	2.26	10.25	2	2	2	2	10.25	2	2	2	2	10.25	2	2	2	2
129 <i>Limnephilus lunatus</i>	12.84	14.70	1.77	2.30	1.04	2.46	11.76	3	2	2	2	11.76	3	2	2	2	11.76	3	2	2	2
130 <i>Limnephilus mammatulus</i>	14.70	17.20	2.03	2.75	1.18	2.95	13.00	0	1	1	1	13.00	0	1	1	1	13.00	0	1	1	1
131 <i>Limnephilus rhombicus reseri</i>	17.08	19.40	2.23	2.78	1.23	3.22	15.40	2	2	2	2	15.40	2	2	2	2	15.40	2	2	2	2
132 <i>Limnephilus rhombicus rhombicus</i>	18.00	21.00	2.18	2.93	1.33	3.40	17.50	0	1	1	1	17.50	0	1	1	1	17.50	0	1	1	1
133 <i>Limnephilus sericeus</i>	11.60	13.80	1.55	2.08	0.93	2.35	11.70	0	1	1	1	11.70	0	1	1	1	11.70	0	1	1	1
134 <i>Limnephilus sparsus</i>	12.40	14.23	1.60	2.05	0.95	2.54	10.86	3	3	3	3	10.86	3	3	3	3	10.86	3	3	3	3
135 <i>Limnephilus stigma</i>	16.80	19.00	2.05	2.76	1.39	3.23	13.35	1	1	1	1	13.35	1	1	1	1	13.35	1	1	1	1
136 <i>Limnephilus vittatus</i>	10.90	12.43	1.48	1.59	0.79	2.01	9.20	2	2	2	2	9.20	2	2	2	2	9.20	2	2	2	2
137 <i>Colpoclauius incisus</i>	8.20	10.05	1.30	1.45	0.65	2.10	10.50	1	1	1	1	10.50	1	1	1	1	10.50	1	1	1	1
138 <i>Grammotaleius nigropunctatus</i>	17.35	19.90	2.04	2.60	1.23	3.28	14.25	1	1	1	1	14.25	1	1	1	1	14.25	1	1	1	1
139 <i>Glyptotaleius pellucidus</i>	17.20	20.20	2.19	3.04	1.38	3.40	16.25	1	1	1	1	16.25	1	1	1	1	16.25	1	1	1	1
140 <i>Anabolia lombarda</i>	15.50	17.70	2.20	2.45	1.38	3.25	12.50	1	0	0	0	12.50	1	0	0	0	12.50	1	0	0	0
141 <i>Rhadicleptus alpestris</i>	13.40	15.05	1.68	1.99	0.89	2.40	10.35	1	1	1	1	10.35	1	1	1	1	10.35	1	1	1	1
	MEAN	14.49	16.58	1.66	2.37	1.13	2.56	12.56				12.56					12.56				
	MAX	23.50	26.68	2.52	3.80	1.76	4.68	19.20				19.20					19.20				
	MIN	6.00	7.20	1.00	0.78	0.40	1.23	5.10				5.10					5.10				
	S.D.	4.01	4.46	0.36	0.68	0.34	0.74	3.28				3.28					3.28				

N. TAXA

N. TAXA

142 *Potamophylax cingulatus alpinus*
 143 *Potamophylax gembaricus gembaricus*
 144 *Potamophylax gembaricus spinulifer*
 145 *Potamophylax inermis*
 146 *Potamophylax nigricornis*
 147 *Acrophylax zerberus*
 148 *Leptotaleius gracilis*
 149 *Halesus appenninus*
 150 *Halesus calabrus*
 151 *Halesus digitatus*
 152 *Halesus nureg*
 153 *Halesus radiatus*
 154 *Halesus rubricollis*
 155 *Halesus tesellatus*
 156 *Melanophylax melampus*
 157 *Melanophylax vesinorum*
 158 *Anisogamus difformis*
 159 *Parachiona picicornis*





Fig. 2 (above): Mean values for forewing length of the species.

Fig. 3 (below): Forewing length: sex ratio.



ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Braueria](#)

Jahr/Year: 2001

Band/Volume: [28](#)

Autor(en)/Author(s): Di Giovanni Maria Vittoria, Goretti Enzo, Chiappafreddo Umberto, Ceccagnoli Donatella

Artikel/Article: [Preliminary results on the biometry of Italian Trichoptera fauna 15-18](#)