

2. *Ptilocolepus*.

In my Atlas of European Trichoptera (MALICKY 1983) I had placed the genus *Ptilocolepus* among Glossosomatidae. I am often asked whether I had transferred the genus from Hydroptilidae where it is placed by most recent authors. No, I have not transferred it, but left it where it was.

MCLACHLAN (1874-80) arranged *Ptilocolepus* in his section III of Rhyacophilidae together with *Glossosoma* and *Agapetus*, i.e. the family Glossosomatidae in our modern sense, and some authors including MARTYNOV and myself have maintained this position. THIENEMANN (1904) described early stages and biology of *Ptilocolepus granulatus* with the conclusion that it fits neither in Glossosomatinae (of Rhyacophilidae) nor in Hydroptilidae in a satisfying manner. Based on this conclusion, ULMER (1907) transferred the genus into Hydroptilidae. At the beginning of the 20th century, one of the dominant hypotheses in biology was the Biogenetic fundamental law (Biogenetisches Grundgesetz) which indicated, that an organism during its ontogeny repeats its phylogeny in the development of morphological characters. Therefore the larval characters were considered more important than the adult characters. The larva of *Ptilocolepus* shares more characters with hydroptilid larvae, but the adults had more in common with glossosomatids. So THIENEMANN and ULMER thought that it must be placed into Hydroptilidae despite the former author finding that it does not fit into either. This ancient hypothesis is now almost forgotten, and in our modern sense all characters have to be considered equally. I have never studied *Ptilocolepus* in detail myself, but it appears now appropriate to draw a clear conclusion from THIENEMANN's results after one century, and to raise the Ptilocolepinae (genera *Ptilocolepus* and *Palaeagapetus*) to family rank Ptilocolepidae. According to the International Code of Zoological Nomenclature, Art. 34.1, the author is MARTYNOV 1913. For detailed information see THIENEMANN (1904).

2. *Pseudoneureclipsis*

In a recent paper, LI & al. (2001) concluded, based on a cladistic analysis, that *Pseudoneureclipsis* may belong to Dipseudopsidae. The following objections may be made to this decision.

1. The larvae of *Pseudoneureclipsis* and *Dipseudopsis* are very different in many characters, as one can easily see by comparing the descriptions and figures by e.g. GIBBS (1968) and TACHET & al. (2001), although they have a similar biology, living in tubes burrowed in the substrate.

2. The genus *Kambaitipsyche* which may have a key position in the analysis was not included in the study.

3. In the character state matrix given by LI & al (l.c.) on which the cladistic analysis is based, the inferior appendages of both Polycentropodidae and *Pseudoneureclipsis* are said to be one-segmented (character no. 10). However, in *Kambaitipsyche* which is considered to belong to Polycentropodidae, the inferior appendages are clearly two-segmented. In *Pseudoneureclipsis*, the inferior appendages have a dorso-basal process which is spine-shaped in most species (but see also *P.philemon* MALICKY & PROMMI 2000), and which is distinctly separated by an intersegmental membrane. The primary condition of inferior appendages in Trichoptera is two-segmented; so, if a caddis adult has a structure, separated by an intersegmental membrane from the basal segment, what else could it be than the second segment? Instead, LI & al.(l.c.) give as character no. 11 "with articulated basodorsal process..." Certainly, the dorso-basal position of this structure is unusual, but also in other caddis groups the second segment may be inserted in another place than distally, e.g. in some odontocerids where it links to the inner surface of the basal segment. Similar dorso-basal processes in many Lepidostomatidae or Leptoceridae cannot be second segments because they are not segmented.

In my opinion, the placement of *Pseudoneureclipsis* in the family Dipseudopsidae can therefore not be accepted. Instead, the relations between *Pseudoneureclipsis* and *Kambaitipsyche* should be studied. I have a larva from the high elevations of Doi Inthanon in Thailand which could be *Kambaitipsyche*, but this is not proved. I hope that one of our doctoral students at the University of Chiangmai will clear up this problem.

References

- GIBBS, D.G., 1968, The larva, dwelling-tube and feeding of a species of *Protodipseudopsis* (Trichoptera: Dipseudopsidae). – Proc.R.ent.Soc.Lond.(A) 43:73-79.
- LI, Y.J., MORSE, J.C., TACHET, H., 2001, Pseudoneureclipsinae in Dipseudopsidae (Trichoptera: Hydropsychoidea), with descriptions of two new species of *Pseudoneureclipsis* from East Asia. – Aquat.Insects 23:107-117.
- MCLACHLAN, R., 1874-80, A monographic revision and synopsis of the Trichoptera of the European fauna. Reprint 1968, Classey: Hampton.
- MALICKY, H., 1983, Atlas of European Trichoptera. Junk: The Hague, 1-300.
- SCHMID, F., 1955, Contribution à l'étude des Limnophilidae (Trichoptera). – Mitt. Schweiz. Ent. Ges. (Beiheft) 28:1-245.
- TACHET, H., MORSE, J.C., BERLY, A., 2001, The larva and pupa of *Pseudoneureclipsis lusitanicus* Malicky, 1980 (Trichoptera: Hydropsychoidea). Description, ecological data and taxonomic considerations. – Aquatic Insects 23:93-106.
- THIENEMANN, A., 1904, *Ptilocolepus granulatus* Pt., eine Übergangsform von den Rhyacophiliden zu den Hydroptiliden. – Allg. Z. Ent. 9:418-424, 437-441.
- ULMER, G., 1907, Trichopteren, in: Genera Insectorum 60:1-259.



If you find a red mark here  in your copy, please read this:

Braueria / Trichoptera Newsletter is produced by private initiative and gets no support from institutions, governments or other official bodies. Its purpose is to communicate among trichopterists, and needs their substantial support. One possibility is the exchange of trichopterological publications. Reprints and books sent to the editor are used to make them known to the community in the Literature List of each number. The other possibility is to send money. The official price of 10 Euro or 10 US\$ covers only one third of the production and mailing cost. Voluntary donations are therefore most welcome. It is expected that reprints or donations (or at least letters or other information) will be sent regularly. From my notes I see that there was no response from you for more than three years. If I have no news from you next time, your address will be deleted from the mailing list.



Moretti collection in Bergamo

In addition to the large "official" Trichoptera collection of the late Professor G.P. Moretti, he had a smaller private collection in his home near Miláno which included much of the material which he had collected when he was young and which was published in some of his early publications. It includes also other caddisflies which he got by exchange or purchase from other workers. Dr. Marco Valle tells me that this collection is now located in the Museo Civico di Scienze Naturali "E. Caffi" in Bergamo where it will be restored and catalogued. Ma.

News from Mara Marinković

For many years we had no information about Dr. Mara Marinković-Gospodnetić who was with us at the symposia in Lunz and Reading, who contributed important papers to the knowledge of the Trichoptera of former Yugoslavia, and described several remarkable species, e.g. *Hydropsyche dinarica*. Dr. Gerhard Tarman tells me now that during his recent visit in Sarajevo he was informed that the collections and notes of Dr. Marinković were in her institute which burnt down during the war, and that she has ceased her scientific activity. Ma.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Braueria](#)

Jahr/Year: 2001

Band/Volume: [28](#)

Autor(en)/Author(s): Malicky Hans

Artikel/Article: [Moretti Collection in Bergamo 20](#)