

Quaternary large mammals from the Apidima Caves (Lakonia, S Peloponnese, Greece)

by

Evangelia TSOUKALA*

TSOUKALA, E., 1999. Quaternary large mammals from the Apidima Caves (Lakonia, S Peloponnese, Greece).—
Beitr. Paläont., 24:207–229, 14 Figs., 12 Tabs., 2 Plates, Wien.

Contents

| | |
|--------------------------------|-----|
| Summary, Introduction | 207 |
| Systematic Palaeontology | 208 |
| Taxonomy | 208 |
| Discussion, Conclusion | 222 |
| References | 221 |

Key words Large mammal fauna (Quaternary Palaeolithic), Apidima caves, (S Peloponnese, Greece)

Summary

The large mammalian remains from the most important Quaternary site in Apidima (South Peloponnese, Greece), have been studied in the present paper. Two Middle Pleistocene hominid skulls, associated with many fossilized animal bones, as well as bone and lithic artefacts, underline the importance of this prehistoric site. The study of the most representative specimens of the fauna showed the presence of carnivores, mainly felids, mustelids and foxes; herbivores, mainly small bovids and cervids; very large mammals, such as hippopotamids and elephants; as well as micro-mammals (leporids and rodents) and birds, which lived in the area from the Middle Pleistocene to Holocene. From the study of this fauna the paleoecology and the paleoenvironment of the area are discussed, in conjunction with the Palaeolithic activities of the Man during this period.

Introduction

Historical overview – Geological background of the region

The Apidima complex of six maritime caves is located in Mesa Mani (Lakonia), on a steep coast of the western Mani peninsula, which is the southernmost continental point of the Balkan peninsula, very near to the active Hellenic subduction zone. Tectonic events, in conjunction with the eustatic sea level fluctuations

of the Quaternary, developed a series of interactive speleoenvironments such as Apidima (BASSIAKOS, 1993). The Neogene is represented by Late Pliocene marine deposits, which are yellowish marls (RONDOYANNI et al., 1995, VERGINIS et al., 1975). Quaternary terrestrial deposits are represented, in some places, by very cohesive cemented breccias, sometimes with fossils; and scree and talus cones compacted during the Mindel, Riss and Würm. North of Apidima, the site of Kalamakia, where excavations are still in progress, has yielded few elephant and rhino remains, as well as remains of caprids and cervids (considered as hominid food), dating to $80-40 \times 10^3$ years BP (LUMEY de & DARLAS, 1994). In the south, the presence of *Dama* and *Elephas* and the absence of carnivore and dwarf forms (BARTSIOKAS, 1998) as well as of many other large mammalian fossil remains, found mainly in cohesive cemented breccia, indicates the "bridges" joining the Cythera island with the continent, so the partition of the island from the Peloponnese mainland happened during the Late Pleistocene (MANOLESSOS, 1955).

The Apidima caves are karstic formations within the Middle Triassic-Late Eocene limestone (Plattenkalk) of depth 500m, from 4 to 24m above sea level, in a vertical zone of depth 20m (Fig.1). The development of the caves is due to the vertical strikes of the limestone, while the horizontal opening is made by the sea according to the geological observations made by RONDOYANNI et al. (1995). In the very thick and cohesive breccia up to 23m above sea level, two Middle Pleistocene hominid skulls were found in cave A, 4m above sea level. These finds give major importance to the Apidima site (PITSIOS, 1979, 1996, COUTSELINIS et al., 1991).

About 20.000 bones, bone fragments and teeth have been collected by Prof. Dr. Theodore PITSIOS and his team since 1978, during four field seasons of systematic excavations from the four caves (A, B, C and D) and they are now stored in the Museum of Anthropology (Athens University). Among the rich macro- and micro- mammal and bird material, a skeleton of a buried woman (+18m), numerous bone and lithic artefacts, fire traces and food remains were found (PITSIOS, 1979, 1983, 1985a,b, PITSIOS & LIEBHABER, 1995). The ESR dating of the travertine calcite and the fossilized

* Aristotle University, School of Geology, 54 006 Thessaloniki, Macedonia, Greece




Figure 1: Map of Greece with the locality (left) and the Apidima maritime caves (above): a) Ground plan 5 m above sea level b, c) vertical sections of the caves (drawings from PITSIOS, 1985b, PITSIOS & LIEBHABER, 1995) (o.s.l.: old sea level)

skeletal remains gave ages for two samples $25-45 \times 10^3$ and $20-30 \times 10^3$ years BP (LIRITZIS & MANIATIS, 1995, BASSIAKOS, 1993).

Systematic Palaeontology

Introduction

The paleontological study is mainly based on the determinable, well preserved and most representative large mammal material, as the statistical study of the bone fragments is given by LAX (1995). The study took place in the Museum of Anthropology of Athens University, where this material is stored, during the years 1991–97.

Taxonomy

Order CARNIVORA Bowdich, 1821
 Superfamily FELOIDEA SIMPSON, 1931
 Family Felidae GRAY, 1821
 Subfamily Felinae PIVETEAU, 1961
 Genus *Panthera* OKEN, 1816

Panthera pardus (LINNAEUS, 1758)

(Plate 1, Figs. 1–8, Text-figs. 2, 5, Table 1)

M a t e r i a l Maxilla fr. dex, I^3 LAO-B-346 sin, C^s LAO-B-091 dex, C^s fr. LAO-B-409 dex, I_3 LAO-B-383a dex, dC LAO-B-175c dex, P_4 LAO-B-159e sin, 2 femur prox. LAO-C-729 dex, 697+700d sin, radius LAO-B-388c +389 sin, radius prox. LAO-C-649c, tibia LAO-C-722 sin, 2 patellae LAO-C-814 sin, LAO-B-390 dex, astragalus LAO-C-641 dex, navicularis LAO-B-389b sin, McV LAO-B-389a dex, PhI LAO-B-410a, PhII LAO-B-348e. **D e s c r i p t i o n** Anterior part of a maxilla, with the alveoli of C^s , P^2 , P^3 and the P^4 antero-external root are preserved. The dimensions are: $L^* P^2$ alveoli=5.2, BP^2 alveoli=4.5, LP^3 alveoli=16.6, BP^3 alveoli=8.5 mm. A slightly worn I^3 is robust, with a well developed labial tubercle and longitudinal crest. A C^s is slightly worn, with well distinguished longitudinal flutes. Half a fragment of a C^s , preserved in cohesive breccia, is very robust. A slightly worn milk canine has dimensions very close to those of recent leopard (dC^s : $L \times B$ =6.5x5.4mm, I.P.H* coll. Regalia, Tab. 1). A P_4 lacks only a small posterior part of the protoconid; the anterior accessory cuspid is well developed and the talonid is wide. Of two femurs, the proximal part with the caput are preserved; and of two radii, one is complete, while the other preserves the proximal part and most of the diaphysis. A tibia is almost complete and robust. Two patellae, an astragalus, a navicular, a fifth metacarpal, a first and a second phalanx of a medial metapodial, are very well preserved and covered with calcite. The Ph2 was found in Cave B in association with caprid and fox remains (Fig. 2).

Table 1. *Panthera pardus*-Apidima: Measurements of teeth and postcranial skeleton, in mm

| *see abbreviations | C* | I ³ | P ₄ | I ₃ | dC _i |
|--------------------|-------------|----------------|----------------|----------------|-----------------|
| L* | 14.0 (16.5) | 8.0 | 16.5 | 6.2 | 6.6 |
| B | 11.0 13.0 | 6.2 | 8.7 | 4.5 | 5.7 |
| H | 63.0 - | 26.5 | 26.5 | | |
| Hcrown | (25.0) | | 11.0 | | |

| | RADIUS | TIBIA | Mc V | PhI | Ph II |
|-----------|-------------|---------|------|------|--------|
| L | 179.0 | (230.0) | 52.2 | 44.2 | 28.0 |
| DTprox. | 23.0 25.4 | (49.5) | 13.7 | 16.5 | 13.5 |
| DAPprox. | 16.2 18.2 | (40.5) | 12.9 | 12.8 | 13.2 |
| DTdia. | 17.0 (18.2) | 19.0 | 7.8 | 11.2 | min8.1 |
| DAPdia. | 10.5 | 22.2 | 8.2 | 8.9 | 8.1 |
| DTdistal | 35.5 | 36.4 | 12.3 | 13.0 | 12.8 |
| DAPdistal | 21.2 | 24.8 | 12.5 | 9.7 | 10.4 |
| DTd.art. | 24.5 | 26.7 | 12.0 | | |
| DAPd.art. | 17.2 | 21.5 | 12.5 | | |

| ASTRAGALUS | PATELLAE | | NAVICULARIS | | FEMUR | |
|------------|----------|--------|-------------|------|-------|------|
| L | 38.4 | L | 34.5 | 33.0 | L | 16.5 |
| DT | 32.5 | DT | 27.0 | 24.5 | DT | 23.0 |
| DAP | 24.5 | DAP | 14.6 | 14.0 | DAP | 18.4 |
| DTcaput | 21.8 | Hart. | 24.0 | 22.0 | | |
| DTcollum | 17.4 | DTart. | 26.7 | 24.2 | | |
| DTtrochlea | 22.5 | | | | | |

***Abbreviations:** (In material the letters A, B, C and D after LAO indicate the cave, where the specimen has been found).

I: Incisor, dC: deciduous canine, C: Canine, P: Premolar, M: Molar, Cr: Cranium, Mx: Maxilla, Md, md: Mandible, diast.:diastema, pr: protocone, hy: hypocone, pa: paracone, me: metacone, pr^d: protoconid, hy^d: hypoconid, pa^d: paraconid, me^d: metaconid, en^d: endoconid, cg: cingulum, a.c.: accessory cuspid, pas: parastyle, mes: metastyle, mcl: metaconulus, tr^d: trigonid, tl^d: talonid, tl: talon, cons.: constriction, , int.: internal, ext.: external, prox., pr.: proximal, dis., d.: distal, a.: anterior, p: posterior, s: superior, i: inferior, L: Length, B: Breadth, H: Height, D: Diameter, DT: Diameter transversal, DAP: Diameter anteroposterior, Id: Infradental, a.s.: articular surface, dia.:diaphysis, Mc: Metacarpal, Mt: Metatarsal, Mp: Metapodial, Ph: Phalanx, art.: articulation, cond.: condylus, inc. sem.: incisura semilunaris, pro. ang.: processus angularis, sym.: symphysis. IPH: Institut Paleontologie Humaine, Paris

Discussion The Apidima leopard is robust like the one from Arago Cave (France), more so than the modern specimens at the collections in Vienna University and in Institut Paleontologie Humaine, Paris and less so than those from Vraona Cave (Attiki, Greece) (NAGEL, 1995, 1999) and Jaurens (France) (Fig. 5) of Würm III age (BALLESIO, 1980), while the protoconid angle of the P₄ Apidima specimen is 75° and 80° for the Jaurens specimen. Leopard specimens from Lezetxiki Cave (Spain) is also much more robust (ALTUNA, 1972), but not those from Stranska Skala (Brno) (HEMMER, 1971, THENIUS, 1972).

The leopard is particularly ubiquitous, widely distributed and eurythermic animal, which flourished in Europa, in the coldest phases of the Würm. Its prey consists of medium-sized herbivores, such as deer (KURTEN, 1968, ARGANT A. & J., 1990, ARGANT, 1991), which are most abundant in the Apidima palaeofauna. In Greece, Pleistocene leopards have been identified at very few sites, such as Vraona cave (NAGEL, 1995), where Latest Pleistocene *Felis (Lynx) lynx* and *Felis silvestris* coexist with the leopard (SYMEONIDIS et al., 1980). Leopard is also found in Dyros associated with hippopotamus of age 25–30x10³ years BP (BASSIAKOS,

1993) and recently Loutraki Almopia caves of about the same age. From the Middle Pleistocene of Arago (Tautavel) felid coexistence (leopard, lynx and wild cat) is also recorded by CREGUT (1976).

Genus *Felis* LINNAEUS, 1758

Lynx KERR, 1792

Felis (Lynx) lynx (LINNAEUS, 1758)

(Plate 1, fig. 9, Text-figs. 3, 5, table 2)

M a t e r i a l : 2C^sLAO-D-087 dex, LAO-B-412b sin, humerus LAO-C-713c dex, calcaneus LAO-C-646 sin

D e s c r i p t i o n Of the upper canines one is an unworn, germ, with hollow root, of a rather small juvenile (L^{*}C^s=6.5, BC^s=5.1, HC^scrown=13.8 mm) (Fig. 3, 2). The dimensions are similar with those of the Stranska Skala lynx (THENIUS, 1972) and that of L' Escale (BONIFAY, 1971) (Fig. 5). Of the post-cranial skeleton a calcite-covered humerus and calcaneus are rather small (of *L. pardina* size, maybe female). Of the former, the distal diaphysis with the upper part of the foramen supracondylicum is preserved. The oldest known *L. lynx* are of Eemian age and their widest distribution was reached during the Würm (KURTEN, 1965,




Figure 2: *Panthera pardus*: Apidima. 1) Maxilla fr. LAO-B-371a sin, a) palatinal, b) basal, 2) I³ LAO-B-346 sin, a) lateral, b) lingual, 3) C¹ LAO-B-091 dex. a) labial, b) lingual, c) posterior, 4) P₄ LAO-B-159e sin, lingual, femur prox. 5) LAO-C-729 dex and 6) 697+700d sin, anterior, 7) radius LAO-B-388c+389 sin a) anterior, b) posterior, 8) tibia LAO-C-722 sin, a) anterior, b) distal, patellae 9) LAO-C-814 sin and 10) LAO-B-390 dex, plantar, 11) astragalus LAO-C-641 dex, a) dorsal, b) plantar, 12) McV LAO-B-389a dex, a) dorsal, b) plantar, 13) PhI LAO-B-410a, 14) PhII LAO-B-348e, dorsal.




Figure 3: *Felis (Lynx) lynx*-Apidima: C^s 1) LAO-D-087 dex, 2) LAO-B-412b sin, a) lingual, b) labial, 3) humerus distal LAO-C-713c dex, a) anterior, b) posterior, 4) calcaneus LAO-C-646 sin, a) anterior, b) lateral

1968). The fossil specimens are similar to Recent ones, which shows that there was no reduction in size after the Pleistocene. The paleoenvironment seem to be forest (preferably coniferous) (KURTEN, 1965, 1968) and its food mainly hares and wild goats, both abundant in the Apidima site.

Table 2. *Felis (Lynx) lynx*-Apidima: Measurements of tooth and post-cranial skeleton, in mm.

| | C ^s | Calcaneus | Humerus | |
|--------|----------------|-----------|---------|-----------------|
| L | 9.7 | L | 40.2 | DT distal |
| B | 8.0 | DAP | 14.0 | DTd.art. |
| H | 43.0 | DT | 18.0 | DT trochlea max |
| Hcrown | 16.0 | DAPd.art. | 9.1 | min 8.0 |
| | | DTd.art. | 10.6 | |

Genus *Felis* LINNAEUS, 1758

Felis silvestris SCHREBER, 1777 (Text-figs. 4, 5, Table 3)

M a t e r i a l Mandible fr. with C_i and P₃ LAO-C-558b sin, M₁ LAO-C-589 sin.

D e s c r i p t i o n The anterior part of a calcite covered mandible includes the crown of an unworn and not well-risen canine, high and slender with 2 buccal longitudinal flutes. The premolar is half-risen and robust and there are also the P₄ alveoli. The dimensions show a more robust wild cat than that from Portel (France, Late Pleistocene) and Arago (Middle Pleistocene) (Fig. 5). The long and large lower carnassial is unworn, with a posterior longitudinal crest (Fig. 4). Its dimensions are close to the maximum for the fossil specimens given by KURTEN (1965). In cave C, a humerus, radius and ulna LAO-C-765b dex, all of the same, rather small individual, and a radius prox. LAO-C-797a dex, are surface findings and seem to belong to a recent cat.

The Pleistocene wild cat is very large (KURTEN, 1965a), although of a remarkable size, is difficult to identify, especially on this fragmentary material but size alone may be sufficient to distinguish wild cat from the domestic form, which are both present in the Apidima




Figure 4: *Felis silvestris*-Apidima: 1) Mandible fr. with C_i and P₃ LAO-C-558b sin, lateral, 2) M₁ LAO-C-589 sin, a)lingual, b)labial

cave C. The teeth of the wild cat, especially the lower carnassial, are much larger than that of the Europe fossil wild cat (KURTEN, 1965). The stable diet consists mainly of small rodents, abundant in the Apidima site. It is mainly a woodland animal preferring mixed forest (KURTEN, 1968).

Table 3. *Felis silvestris*-Apidima: Measurements of mandible, teeth and post-cranial skeleton, in mm

| Mandible | C _i | P ₃ | M ₁ |
|---------------------------------------|----------------|----------------|----------------|
| Hmddiast. | 10.7 | L | 5.9 (6.0) |
| Hmd(P ₃ , P ₄) | 10.2 | B | 4.3 (3.2) |
| Dmax.sym. | 15.4 | Hcrown | 10.5 |
| Dmin.sym. | 6.0 | | 13.0 |

Superfamily CANOIDEA SIMPSON, 1931

Family Canidae GRAY, 1821

Genus *Vulpes* (LINNAEUS, 1758)

Vulpes vulpes (LINNAEUS, 1758)

(Plate 1, figs. 10, 11, Text-figs. 6, 7, Table 4)

M a t e r i a l: Maxilla fr. LAO-C-715b dex, 2C^s LAO-C-610 dex and 641b sin, P⁴ LAO-C-727a dex, P⁴ fr. 571b dex, 2M¹ LAO-C-746c (maxilla fr. with the C^s, P¹ alveoli and P²) and LAO-C-787d dex, M¹ fr. LAO-D-681a dex,




Figure 5: Ratio-diagram comparing the dimensions of various teeth and elements of the post-cranial skeleton of felids: 1. LC^s, 2. BC^s, 3. LI³, 4. BI³, 5. LC_p, 6. BC_p, femur: 7. DTprox., radius: 8. L, 9. DTprox., 10. DTdia, 11. DTdistal, tibia: 12. DTdistal, astragalus: 13. L, 14. DT, 15 DAP, 16. DTcollum, 17. DTcaput, calcaneus: 18. L, 19. DT, 20. DAP, navicularis: 21. L, 22. DT, 23. DAP, patella: 24. L, 25. B, McV: 26. L, 27. DTprox., 28. DTdia, 29. DTdistal, PhiI: 30. L, 31. DTprox., 32. DTdia, 33. DTdistal. Standard *F. (L.) lynx* -recent, IPH coll. Regalia. (LAO: Apidima, IPH: Institut Paleontologie Humaine-Paris, Jaurens (BALLESIO, 1980), L'Escale (BONIFAY, 1971), Stranska Skala (THENIUS, 1972), Vraona, Attiki (NAGEL, 1995).

| * see abbreviations | C ^s | P ⁴ | M ¹ | M ² |
|---------------------|----------------|----------------|----------------|----------------|
| L* | 7.4 | 15.6 | 9.8 10.3 | - |
| B | 4.9 | 7.0 | 13.5 2.5 | 11.3 |
| H | 31.0 | | | |
| Hcrown | 17.0 | | | |

| | P ₁ | P ₂ | P ₃ | P ₄ | M ₁ | M ₂ | M ₃ | dC _i | D ₄ |
|--------------------------|----------------|--------------------------------------|----------------|----------------|----------------|------------------------|----------------|-----------------|----------------|
| L | 4.4 | 4.4 | 9.0 | 9.5 10.3 | 11.0 | 16.0 15.7 | 16.2 14.4 | 7.2 | 4.0 3.5 |
| B | 2.7 | 3.1 | 3.4 | 3.8 3.8 | 4.3 | 6.0 6.5 | 6.5 5.5 | 5.2 | 2.5 4.0 |
| H | 9.8 | 9.7 | | | | 14.2 | 17.7 | | 16.8 |
| Hcrown | | | | 7.0 | | | 9.0 9.0 | | |
| MANDIBLE | | | | | | | | | |
| LI ₁ -cond. | 109.0 | LP ₁ alv. | 35.2 | DTcond. | 14.7 | HmdP ₃ a. | | | 13.8 |
| LC _i a.-cond. | 107.5 | LM _i alv. | 26.2 | Hcond. | 7.3 | HmdM ₁ a. | | | 16.2 |
| LC _p -cond. | 100.5 | LP ₁ -M ₃ alv. | 61.2 | Hmddiast. | 11.6 | HmdM ₁ p. | 16.5 | 17.0 | |
| Dmax sym. | 28.0 | Dmin sym. | 10.0 | | | Hramus(pro.ang.-cond.) | | | 18.0 |

| | RADIUS | TIBIA | CALCANEUS | |
|-----------|--------|-------|-----------|------|
| L | | 130.0 | L | 30.5 |
| DTprox. | | 20.8 | DT | 11.4 |
| DAPprox. | | 22.3 | DAP | 12.0 |
| DTdia. | | 8.5 | | |
| DAPdia. | | 8.3 | | |
| DTdistal | 14.5 | 15.0 | | |
| DAPdistal | 8.2 | 10.2 | | |
| DTd.art. | 12.0 | 11.0 | | |
| DAPd.art. | 7.2 | 9.5 | | |

Table 4. *Vulpes vulpes*-Apidima: Measurements of teeth, mandible and post-cranial skeleton, in mm




Figure 6: *Vulpes vulpes*-Apidima: 1) C⁵ LAO-C-610 dex, a)lingual, b)labial, 2) P⁴ LAO-C-727a dex, a) labial, b) lingual, c)occlusal, 3) M¹ LAO-C-746c a) labial, b) occlusal, mandible frs. 4) with P₃, M₁ LAO-C-543 dex, lateral and 5) with M₁, M₂ LAO-B-352c-353 dex, a) lateral, b) M₁ lingual, c)occlusal, 6) P₂ LAO-C-559c sin, lingual, 7) P₃ LAO-B-358c dex, labial, 8) P₄ LAO-C-589 sin, a) labial, b) lingual, 2M₁, 9) LAO-A-033 dex, lingual and 10) LAO-C-641b sin, a) labial, b) lingual, 11) dC₁ LAO-D-82z sin, a) lingual, b) anterior, 12) D₄ 88d sin, a) lingual, b) occlusal, 13) radius distal LAO-B-393b sin and 14) tibia LAO-B-402a+404 sin, anterior, 15) calcaneus LAO-B-348e dex, medial.

M² LAO-C-813d dex, 2mandible frs. with P₃, M₁ LAO-C-543 dex and with M₁-M₂ LAO-B-352c-353 dex, C₁ fr. LAO-C-539b dex, P₁ and M₃ LAO-C-786b dex, P₁ LAO-C-669c dex, P₂ LAO-C-559c sin, P₃ LAO-B-358c dex, 2P₃ fr. LAO-D-239f and LAO-C-584a dex, P₄ LAO-C-589 sin, P₄ fr. LAO-C-501b dex, 2M₁ LAO-A-033 dex and LAO-C-641b sin, dC₁ LAO-D-82z sin, 2D₄ LAO-D-86h and 88d sin, femur fr. LAO-C-813d, radius distal LAO-B-393b sin, tibia LAO-B-402a+404 sin, calcaneus LAO-B-348e dex.

Description A maxilla fragment with molar alveoli is preserved. Of two C⁵ one is robust, while the other is completely worn. An upper carnassial, of height 3.5mm, has anteriorly a weak crest and cingulum, while the protocone is pronounced anteriorly. The M¹ are relatively robust, and an external cingulum is well developed. The molar M¹ LAO-C-746c is associ-

ated with a maxilla fragment. Of two mandible fragments, one lacks the processus coronoideus, and contains the P₃ and M₁, while the other is small fragment with the first two, almost unworn, molars (Fig.6). Two small isolated and slightly worn teeth, P₁ and M₃, are each other, probably of the same individual. An anterior half of P₃ and posterior half of a P₃ and of a P₄ are preserved, with breadths 3.2, 3.4 and 4.5mm respectively. A lower carnassial lacks the exterior half of the trigonid, while the talonid has developed tubercles. Of a long, slightly worn P₄ there is a short but distinguishable cingulum, on the hiddenmost part. Long premolars characterize the Apidima fox (Fig. 7). The slightly worn lower carnassials lack an accessory cuspid between the well-developed metaconid and hypoconid, which is present on those of the recent species. A lower milk canine is slender, with a well-distinguished cuspid at

| | C ^s | | | | C _i | MANDIBLE | | | M ₁ | | P ₂ | HUMERUS | | | ULNA |
|--------------------|----------------|-------|------|------|----------------|----------|--------------------------------------|------|---------------------|------|----------------|------------|-------------------|--|-------|
| | n | x | min | max | | | | | L | B | | | | | n=2-3 |
| L | 4 | 4.80 | 4.5 | 5.0 | 5.5 | 5.0 | Lmdcond.-C _i alv. | 53.0 | L | 11.0 | 4.5 | L | 73.8 | | |
| B | 4 | 3.80 | 3.6 | 3.9 | 3.0 | 3.5 | Lmdcond.-I _i alv. | 54.0 | B | 4.5 | 2.9 | DTprox. | 14.0 | | |
| H | 3 | 20.23 | 19.2 | 21.0 | 17.6 | 18.0 | LI _i -M ₂ alv. | 38.0 | H _{crown} | 5.3 | | DAPprox. | 15.0 | | |
| H _{crown} | 3 | 9.87 | 9.4 | 10.2 | 8.0 | 9.8 | LP ₁ -M ₂ alv. | 30.8 | Ltr ^d | 7.2 | | DTdia. | 5.5 | | |
| | | | | | | | LC _i -M ₂ alv. | 35.0 | Ltl ^d | 3.8 | | DAPdia. | 6.0 | | |
| | | | | | | | LP _i alv. | 16.5 | Btl ^d | 3.3 | | DTdistal | 15.97 (15.6-16.3) | | |
| | | | | | | | LM _i alv. | 14.8 | | | | DAPdistal | 9.33 (8.7-10.3) | | |
| | | | | | | | DTcond. | 9.4 | Bc _i alv | 4.5 | 4.6 | DTtrochlea | 11.1 (11.0-11.2) | | |
| | | | | | | | Hcond. | 3.7 | | | | DTcaput | 10.0 | | |
| | | | | | | | Dsym. | 14.5 | | | | | | | |
| | | | | | | | max | 8.2 | | | | | | | |
| | | | | | | | min | | | | | | | | |


Table 5. *Martes foina*-Apidima: Measurements of mandible, teeth and post-cranial skeleton, in mm

Figure 7: Ratio-diagram comparing the dimensions of various teeth and elements of the post-cranial skeleton of foxes. 1. LC^s, 2. BC^s, 3. LP⁴, 4. BP⁴, 5. LM¹, 6. BM¹, 7. LI_i-cond., 8. LC_i-a.-cond., 9. LC_p-cond., 10. DTcond., 11. Hcond., 12. Hmddiast., 13. HmdP₃a, 14. HmdM₁a, 15. Hramus (pro.ang.-cond.), 16. LP₂, 17. LP₃, 18. LM₁, 19. BM₁, 20. LM₂, 21. LD₄, Radius: 22. DTdistal, 23. DAPdistal, Tibia: 24. L, 25. DTprox., 26. DAPprox., 27. DTdia. 28. DTdistal, Calcaneus: 29. L, 30. DT, 31. DAP. Standard: *Vulpes vulpes* -recent, IPH coll. Regalia (Jaurens after BALLESIO, 1979)

the base of the crown. A milk carnassial has a very developed metaconid and hypoconid, while the talonid is tricuspid. Only distal part of a slender radius is preserved. A complete, well-fossilized, tibia and calcaneus were found in cave B, in association with the leopard.

The Apidima fox has similar dimensions with that from Jaurens (BALLESIO, 1979). The comparison of the dimensions with those from Portel (Late Pleistocene), Arago and Lazaret (Middle Pleistocene) (France) is shown on the ratio-diagram (Fig. 7). In Greece, Middle Pleistocene *V. vulpes* has been identified at Megalopolis (SICKENBERG, 1976) and Petralona cave (TSOUKALA, 1989), while Late Pleistocene species have been recorded in Vraona cave, Attiki (SYMEONIDIS et al., 1980) and Agios Georgios cave (Kilkis, TSOU-

KALA, 1992a). In Apidima, its presence is consistent with the presence of abundant rodent, leporid and bird remains (KURTEN, 1968).

Family Mustelidae FISCHER, 1817
 Genus *Meles* LINNAEUS, 1758
Meles meles LINNAEUS, 1758
 (Text-fig. 8)

Material C^s LAO-C-278b dex, C_i LAO-C-443c dex, M₁ LAO-C-628d sin, ulna distal LAO-B-367b sin. Description: AC^s is very worn, with robust root (LC^s=7.0, BC^s=5.4 mm). Of a slender, short-crowned C_i, the base of the intensely curved crown is of oval transversal section (LC_i=5.9, BC_i=4.9 mm). A little-worn lower carnassial lacks small postero-lingual part (LM₁=16.3, BM₁=7.3, Ltr^d=8.6, Btr^d=6.4 mm). Of the




Figure 8: *Meles meles*-Apidima: 1) C^s LAO-C-278b dex, labial, 2) C^s LAO-C-443c dex, a) labial, b) lingual, 3) M₁ LAO-C-628d sin, a) labial, b) occlusal, 4) ulna distal fr. LAO-B-367b sin, a) anterior, b) lateral

post-cranial skeleton, an ulna preserves the robust distal part (Fig. 8).

The badgers although inhabited in the Early Middle Pleistocene of Europe, they are similar to modern species (KURTEN, 1965). Apidima badger seem to be less robust (sexual dimorphism is also considered), as this from Portel (Late Pleistocene), Arago and Lazaret (Middle Pleistocene).

Genus *Martes* PINEL, 1792

Martes foina (ERXLEBEN) (Text-fig. 9, Table 5)

M a t e r i a l 4C^s LAO-C-755b dex, 811b, 765 and

568c sin, 2 mandible frs. with M₁ LAO-C-483 dex and with P₂ LAO-C-779b dex, C₁ LAO-C-735a dex and 254i sin, humerus LAO-C-808a sin, 2 humerus distal LAO-C-483b sin and LAO-C-772 dex, 2 ulnae LAO-C-778a sin and LAO-C-483b

D e s c r i p t i o n : Four C^s are slightly worn or unworn, of circular transversal section at the base of the crown, with well developed root. Of two mandible fragments, one preserves condylus, all alveoli and lower carnassial, while the other preserves an anterior part of corpus, with premolar alveoli and P₂ (Fig. 9). The anterior and posterior mental foramina are rather close each other, while in pine marten they are farther apart, and this character, according to Janossy (ANDERSON, 1970), is useful in separating the two species. The unworn carnassial has a rather long trigonid, but not broad talonid, with well-developed hypoconid, hypoconulid and metaconid. The height of the mandible corpus, in front of the P₂, is 9.0mm. Of three humeri, bearing foramen supracondyllicum, one is complete, while the other two lack the proximal epiphysis. Of two ulnae, one lacks only the tuber olecrani, while the other lacks the distal part, therefore seem to be of a juvenile (DTprox.art.=6.8mm). These post-cranial bones seem to belong to a recent marten.

Beach or stone marten, occurring in mixed forests and on rocky hill sides, has a Middle East or south-west Asia origin and a more southern and eastern distribution than *Martes martes*. Although it is absent from their fossil record, it is found on Crete and Rhodes islands (may be due to man), but not on other Mediter-




Figure 9: *Martes foina*-Apidima: C^s 1) LAO-C-755b dex, 2) 811b, 3) 765 lingual and 4) 568c sin, a) lingual, b) labial, mandible frs. with M₁, 5) LAO-C-483 dex a, d) lingual, b) labial, c, e) occlusal, with P₂, 6) LAO-C-779b dex, labial, 7) C₁ LAO-C-735a dex, 8) 254i sin, lingual, 9) humerus LAO-C-808a sin, posterior, humerus distal 10) LAO-C-483b sin, a) posterior, b) anterior, 11) LAO-C-772dex, 12) ulna LAO-C-483b sin, lateral.

ranean islands, nor in N. Africa and Arabia (ANDERSON, 1970). In Greece, Late Pleistocene mustelids have been identified at Vraona cave, Attiki (*Meles meles*, *Martes* sp., *Mustela cf. putorius* (SYMEONIDIS, et al., 1980), and *Mustela putorius robusta* at Agios Georgios cave, Kilkis (TSOUKALA, 1992a).

Order ARTIODACTYLA OWEN, 1848

Suborder RUMINANTIA SCOPOLI, 1777

Family Bovidae GRAY, 1821

Subfamily Caprinae GILL, 1872

Genus *Capra* LINNAEUS, 1758

***Capra ibex* LINNAEUS, 1758**

(Plate 1, Figs. 12, 13, Plate 2, Figs. 1–4,
Text-figs. 10, 11, Table 6)

M a t e r i a l : Skull fr. with horn LAO-B-163a, horn frs LAO-B-78, 78d, 80, 137, 6maxilla frs with P²-M³ LAO-C-285a, with P³-M³ LAO-D-26a sin, with P³-M² LAO-B-174a/b, with P⁴-M³ LAO-D-24 dex, with P²-P⁴ LAO-B-190, with M²,M³ LAO-74, P³ LAO-B-414b, P⁴ LAO-70a, 3M¹ LAO-C-269b and 622c, LAO-B-158e, 9M² LAO-B-148, 358d, 362c, 426c, LAO-C-251, 653a, 656, 814a dex, LAO-D-821a, M^{1,2} and M_{1,2} LAO-A-19, 34, 44a, LAO-B-70, 77, 166a, 166c, 408b, 413b, 414b, 415a, 417c, 418a, 11M³ LAO-A-07, 08a, 15 dex, LAO-B-139, 364, 393a sin, 410c, 416c, 426, LAO-C-655, 666 dex, 12mandible frs with P₂-M₃ LAO-C-728 sin, 2with P₃-M₃ LAO-B-74, 179 sin, with P₄-M₃ LAO-D-18a sin and 79e dex, , with M₁,M₂ LAO-B-30, with M₂,M₃ LAO-B-382 sin, with M₃ LAO-C-497, with D₃-M₃ LAO-B-167c, with D₂-M₁ LAO-C-509, 2with D₄-M₁ LAO-B-151, 417, with D₃-D₄ LAO-A-41, with D₄ LAO-B-375,377, mandible fr. juv. LAO-B-361, 361a, I₁ LAO-A-105, LAO-B-80, 165e, 357c,d, 416c, LAO-C-579a, 585a, 640c, 653a, 786b, 2P₂ LAO-B-375b, LAO-C-267, P₃ LAO-A-120 sin, 6P₄ LAO-A-34c, 101c, LAO-B-70, 156n, 168, 176b, 5M₁ LAO-B-28a, 159f, 171c, 362c, LAO-C-622c, 2 M₁,M₂ LAO-B-161e, 360, 3M₂ LAO-C-695b, 699c, 703b, 5M₃ LAO-B-189b, 347a, 392, 426e, LAO-C-703b, 5D₄ LAO-B-77, 78, 80, 347a, 419a, pelvis fr. LAO-B-65, humerus distal LAO-C-64 3b dex, 2radius frs LAO-B-387a (and ulna) dex, 92, 4astragali LAO-D-681b, LAO-B-75, 348e, 363 dex, 2cuboscaphoidea frs LAO-B-74a, 139a, McIII+IV LAO-B-385 dex, MtIII+IV fr. juv. LAO-B-425, Mp distal LAO-A-55a, 2Phi LAO-B-156g and LAO-D-232b, 3PhII LAO-B-344b 346a, 416, PhIII LAO-B-153d. **D e s c r i p t i o n** A juvenile skull fragment, with part of a horn (its dimensions at the base, are: Dmaxxmin = 25x21 mm), as well as few horn fragments are preserved (Fig.10.1, 2, pl.1.12). A maxilla fragment includes all cheek-teeth, which are slender and thin enamelled (P² and M³ especially), indicating a not robust, but young individual. Molars vary little to deeply worn. A M³ has intense metastyle; some morphological differences with that from Petralona are indicated on Fig.10. Some mandibles are robust, and may belong to the same individual. Apidima ibex is

less robust than the Petralona ibex. Concerning P², P⁴ and M¹ of the former are notably smaller than the latter, the large M₃ of which, has the talonid angle more open. Of the post-cranial skeleton, a cuboscaphoideum is robust (DT=37mm); a distal, calcite-covered humerus bears trace of carnivore gnawing; and a radio-ulna lacks the olecranon. A complete metacarpal is calcite-covered, the dimensions of which indicate a female (Fig.10, pl.1.13). Of two PhiI, one is well fossilized and robust, while the other seems recent. A PhiII bears a big hole posteriorly, probably trace of carnivore gnawing (86h). Of two astragali, one is rather small, while the other is robust. Caprid bones from Cave B, have been found in breccia, highly fossilized, while there are recent bones of *Capra hircus*. This species is descended from the wild goat *C. aegagrus* ERXLEBEN, which is distributed from Greek islands and Asia Minor eastward to India (KURTEN, 1965).

Family Cervidae GOLDFUSS, 1820

Genus *Megaloceros* BROOKES, 1828

***Megaloceros* sp.**

(Text-fig. 12, Table 7)

M a t e r i a l : Maxilla frs. with P³-M¹ LAO-C-629 sin, M² LAO-C-798d, 800d sin, mandible fr. LAO-C-647a with P₃-M₃ 647 sin, M₂ LAO-C-308e sin, P₄ LAO-C-803a sin.

D e s c r i p t i o n : A maxilla fragment is calcite-covered, with thick-enamelled worn teeth. A large M² has very thick (1.4mm) enamel and an intense cingulum between lobes, but less strong than of this from the Petralona *Premegaceros* sp. (Fig. 12). The mandible is very fragmentary, while the teeth are very robust and well preserved. The P₃ is very robust, the M₁ is worn, the M₂ has a small cingulum between lobes and the M₃ is middle worn (crown-height 23mm), with very intense cingulums between lobes and postero-lingual of talonid.

The giant deer is widely distributed in Europe and northern Asia, occurred in the Holsteinian and Eemian interglacials and also in "cold stages", therefore it doesn't indicate particular stage(s) (LISTER, 1986). In Macedonia (N. Greece), among other sites (TSOUKALA, 1992b), the giant deer has been recorded in Agios Georgios cave, Kilkis (BASSIAKOS & TSOUKALA, 1996) and in Drama (KOUFOS, 1981), of Late to Latest Pleistocene age.

Genus *Cervus* LINNAEUS, 1758

***Cervus elaphus* LINNAEUS, 1758**

(Plate 2, Fig. 1, Text-fig. 12, Table 7)

M a t e r i a l 4 Antler frs. LAO-C-311, 529, 518 and 782, 6maxilla frs. with P²-M³ LAO-D-27 dex, with M²-M³ LAO-C-585 dex, with D²-M¹ LAO-D-242 dex, LAO-C-286b dex and 309d sin and with D² LAO-C-327d, 2C^s LAO-C-254, 290a, 2P² LAO-B-159c sin, LAO-D-90 dex,




Figure 10: Artiodactyla-Apidima: *Capra ibex*, 1)horn frs, a) LAO- B- 80, b)LAO-B-79, 2) M³ a) LAO-B-364sin, palatinal, b) LAO -C-703b, c) Petralona PEC 326 dex, labial (for comparison), 3)astragalus LAO-B-363 dex, a) anterior, b) posterior 4) McIII+IV LAO-B-385 dex, *Dama dama* 5) Mc distal fr. LAO-B-087a.

| TEETH | n | x | min | max | s _{n-1} | v | | n | x | min | max | s _{n-1} | v |
|---------------------------------|------|-------|---------------------------------|------------|------------------|-------|---------------------------------|-----------------|-------|------|--------|------------------|-------------|
| LP ² | 1 | 7.8 | | | | | LP ₂ | 3 | 6.33 | 6.0 | 7.0 | | |
| BP ² | 1 | 4.8 | | | | | BP ₂ | 3 | 4.70 | 4.0 | 5.5 | | |
| LP ³ | 5 | 9.24 | 8.2 | 10.0 | 0.81 | 9.10 | LP ₃ | 4 | 8.65 | 8.4 | 9.0 | | |
| BP ³ | 5 | 8.84 | 7.7 | 9.0 | 0.69 | 7.78 | BP ₃ | 4 | 6.73 | 6.1 | 7.7 | | |
| LP ⁴ | 6 | 9.26 | 7.0 | 11.0 | 1.65 | 17.88 | LP ₄ | 9 | 10.15 | 9.0 | 11.3 | 0.72 | 7.05 |
| BP ⁴ | 6 | 10.53 | 9.0 | 11.5 | 0.86 | 8.20 | BP ₄ | 9 | 7.28 | 6.2 | 8.4 | 0.64 | 8.96 |
| LM ¹ | 10 | 14.40 | 12.0 | 16.5 | 1.88 | 13.10 | LM ₁ | 19 | 14.27 | 10.5 | 16.5 | 1.78 | 12.49 |
| BM ¹ | 10 | 12.84 | 10.5 | 14.5 | 1.26 | 9.74 | BM ₁ | 19 | 8.55 | 7.5 | 9.6 | 0.52 | 6.13 |
| LM ² | 24 | 18.45 | 17.0 | 22.0 | 1.30 | 7.08 | LM ₂ | 22 | 17.98 | 15.0 | 20.5 | 1.34 | 7.45 |
| BM ² | 24 | 14.61 | 12.0 | 17.2 | 1.31 | 8.95 | BM ₂ | 22 | 10.66 | 9.0 | 14.5 | 1.44 | 13.52 |
| LM ³ | 13 | 23.52 | 20.0 | 26.0 | 1.64 | 7.00 | LM ₃ | 12 | 26.97 | 24.2 | 30.0 | 1.59 | 5.91 |
| BM ³ | 17 | 13.70 | 11.5 | 15.5 | 0.98 | 7.18 | BM ₃ | 12 | 10.29 | 9.5 | 11.8 | 0.62 | 6.08 |
| LD ² | 1 | 8.5 | | | | | LD ₂ | 1 | 5.00 | | | | |
| BD ² | 1 | 6.0 | | | | | BD ₂ | 1 | 3.80 | | | | |
| LD ³ | 1 | 12.3 | | | | | LD ₃ | 4 | 9.20 | | | | |
| BD ³ | 1 | 9.0 | | | | | BD ₃ | 3 | 5.87 | | | | |
| LD ⁴ | 1 | 14.0 | | | | | LD ₄ | 11 | 17.45 | | | | |
| BD ⁴ | 1 | 9.5 | | | | | BD ₄ | 11 | 7.22 | | | | |
| LM ¹ | 1 | 14.0 | | | | | | | | | | | |
| BM ¹ | 1 | 11.0 | | | | | | | | | | | |
| MANDIBLE | | | | | | | MAXILLA | | | | | | |
| LP ₂ -M ₃ | 76.2 | | LP ₄ -M ₃ | 56.0 | 53.0 | | LM ⁶ | 49.0 | | | | | (40.0-55.5) |
| Hmd(diast.) | 14.5 | | LM ₁ | 66.0 | 64.0 | | LP ² -M ³ | 61.0 | | | | | |
| Hmd(M ₁) | 28.2 | | | | | | | | | | | | |
| HUMERUS | | | | RADIUS | | | | CUBOSCAPHOIDEUM | | | | | |
| DTdistal | | 37.5 | | DTprox. | 40.0 | 35.5 | | L | | | | | |
| DTd.art. | | 36.5 | | DAPprox. | 18.0 | 17.3 | | DT | | | | | |
| DAPd.art. | | 21.5 | | | | | | DAP | | | | | |
| ASTRAGALUS | | | | McIII+IV | | | | PhI | | | | | |
| Llat. | 28.2 | 37.2 | 34.0 | L | 129.0 | 46.0 | 36.3 | 28.0 | 28.0 | 30.5 | H | | |
| Lmed. | 25.8 | | | DTprox. | 30.5 | 16.2 | 12.5 | 16.0 | 14.3 | 17.0 | Hart. | | |
| DTprox. | 16.8 | | | DAPprox. | 21.0 | 18.5 | 14.3 | 15.5 | 14.5 | 17.2 | DTart. | | |
| DTdistal | 18.3 | 25.4 | 22.2 | DTdia. | 21.2 | 12.6 | 9.8 | 11.2 | 10.0 | 12.0 | | | |
| DAPlat. | 14.7 | | | DAPdia. | 15.0 | 15.6 | 11.7 | 13.0 | 11.2 | 14.0 | | | |
| DAPmed. | 16.0 | 21.2 | 19.2 | DTdistal | 34.0 | | | | | | | | |
| | | | | DAPdistal | 20.0 | | | | | | | | |
| | | | | DTtrochlea | 15.8 | | | | | | | | |

Table 6. *Capra ibex*-Apidima: Measurements of teeth and post-cranial skeleton in mm




Figure 11: Ratio diagram comparing the dimensions of teeth and metacarpal of the Pleistocene *Capra*

P³ LAO-D-90a dex, 2M¹ LAO-C-708a and 636 dex, M^{1,2} LAO-D-239f sin, 5M² LAO-C-280i, 627, 677a, 707 dex, 655 sin, M³ fr. LAO-B-143a, D² LAO-C-809d, D³ LAO-C-305h dex, 2mandible frs with D₂-M₁ LAO-C-309c sin, mandible fr. and P₄, M₁ LAO-C-286c, I₁ LAO-B-78, 159d, 165e, 367b, LAO-C-308g, 553, 633, 644d, 779a, 813c, P₄ LAO-C-801b, M₁ LAO-C-264, 2M_{1,2} LAO-D-227d dex, LAO-C-727b, M₃ LAO-D-86d dex, M₃ fr. LAO-D-197b dex, D₃ LAO-C-811b, 2D₄ LAO-C-691a, 785, atlas LAO-C-244, epistropheus LAO-C-780, cervical LAO-C-602, 2pelvis fr. LAO-D-80, 242b sin, humerus distal LAO-C-546, patella LAO-C-445a, scaphoid and pyramidal LAO-C-321c dex, semilunar LAO-C-325d sin, 6astragali LAO-C-306b, 445a, 531, 590a dex, LAO-D-231, 237c, calcaneus LAO-242a dex, 2cuboscaphoidea LAO-C-282d and LAO-C-451a, 4Mp frs LAO-C-307, 309f, 311a, 588, 4PhI LAO-C-445a, 498, 582a,

LAO-D-237, 3PhII LAO-C-763f, 813e, LAO-D-86h, 2PhIII LAO-C-533a, 757.

Description: Of four small antler fragments one is tine, tine-tip and two almost circular at the base (D = 13.0 mm, 23.5 x 26.5 and 15.0 x 16.0mm). Among the upper teeth, the molars are more or less worn, with intense cingulum, in some cases antero-palatal, or/and between the two lobes. Of very worn milk teeth, there is a strong cingulum on D² and a very intense and high anteriorly on D³ and a weak one between the two lobes of D⁴. A mandible with milk teeth is probably of the same individual with the maxilla LAO-D-309d. Lower milk teeth are very thin enamelled. A M₂ has a small external cingulum between the lobes, a M_{1,2} has no cingulum, with wrinkled enamel and well developed metastylid. Its occlusal length is 32.0 mm.

| | <i>Megaloceros</i> sp. | | | | <i>Cervus elaphus</i> | | | | | | <i>Dama dama</i> | | | | | |
|---------------------------------|------------------------|--------|------|------|-----------------------|--------|------|------|------------------|------|------------------|-------|------|------|------------------|-------|
| | n | x | min | max | n | x | min | max | s _{n-1} | v | n | x | min | max | s _{n-1} | v |
| LP ² | | | | | 4 | 16.20 | 15.5 | 16.5 | | | 8 | 13.81 | 12.2 | 15.8 | 11.35 | 8.22 |
| BP ² | | | | | 4 | 16.72 | 15.0 | 19.2 | | | 8 | 12.59 | 12.0 | 13.8 | 0.54 | 4.27 |
| LP ³ | 1 | 20.8 | | | 2 | 17.30 | 17.0 | 17.6 | | | 5 | 13.04 | 11.9 | 14.1 | 0.83 | 6.36 |
| BP ³ | 1 | 21.6 | | | 3 | 19.50 | 18.5 | 20.0 | | | 5 | 15.40 | 13.9 | 16.5 | 1.07 | 7.00 |
| LP ⁴ | 1 | 18.0 | | | 1 | 18.20 | | | | | 6 | 12.57 | 11.5 | 13.3 | 0.71 | 5.66 |
| BP ⁴ | 1 | 21.7 | | | 1 | 21.30 | | | | | 6 | 16.90 | 15.8 | 18.0 | 0.88 | 5.21 |
| LM ¹ | 1 | 25.5 | | | 1 | 24.20 | | | | | 13 | 18.90 | 16.5 | 21.6 | 1.76 | 9.18 |
| BM ¹ | 1 | 25.0 | | | 1 | 24.80 | | | | | 13 | 18.42 | 16.5 | 20.8 | 1.29 | 6.96 |
| LM ² | 1 | 22.5 | | | 5 | 26.76 | 26.0 | 27.5 | 0.60 | 2.25 | 6 | 21.65 | 19.8 | 23.0 | 0.46 | 2.20 |
| BM ² | 1 | 28.5 | | | 5 | 24.82 | 22.1 | 26.0 | 1.56 | 6.29 | 6 | 21.26 | 19.8 | 22.0 | 1.04 | 4.90 |
| LM ³ | 1 | 27.5 | | | 1 | 26.40 | | | | | 7 | 20.45 | 19.2 | 22.5 | 1.10 | 5.39 |
| BM ³ | 1 | (27.3) | | | 1 | 25.00 | | | | | 6 | 19.65 | 18.5 | 20.8 | 0.91 | 4.66 |
| LD ² | | | | | 6 | 17.85 | 17.0 | 19.2 | 0.83 | 4.64 | 7 | 12.74 | 12.4 | 13.6 | 0.46 | 3.62 |
| BD ² | | | | | 6 | 11.17 | 10.6 | 12.7 | 0.50 | 4.48 | 7 | 8.74 | 8.20 | 9.0 | 0.29 | 3.36 |
| LD ³ | | | | | 4 | 21.83 | 20.7 | 22.5 | 0.78 | 3.57 | 8 | 14.62 | 13.0 | 15.0 | 0.71 | 4.88 |
| BD ³ | | | | | 4 | 16.87 | 16.0 | 17.5 | 0.71 | 3.20 | 8 | 12.37 | 10.4 | 13.4 | 0.92 | 7.44 |
| LD ⁴ | | | | | 3 | 21.70 | 20.6 | 22.4 | | | 8 | 15.08 | 15.2 | 16.4 | 0.38 | 2.42 |
| BD ⁴ | | | | | 3 | 18.87 | 18.5 | 19.1 | | | 8 | 15.68 | 14.2 | 17.4 | 1.08 | 6.92 |
| LP ² -M ³ | | | | | 1 | 126.50 | | | | | 1 | 90.0 | | | | |
| LD ² -D ⁴ | | | | | | | | | | | 2 | 43.15 | 42.3 | 44.0 | | |
| LP ₂ | | | | | | | | | | | 2 | 9.35 | 8.7 | 10.0 | | |
| BP ₂ | | | | | | | | | | | 2 | 6.20 | 5.6 | 6.8 | | |
| LP ₃ | 1 | 19.50 | | | | | | | | | 5 | 12.86 | 12.5 | 13.5 | 0.49 | 3.87 |
| BP ₃ | 1 | 12.20 | | | | | | | | | 5 | 7.58 | 6.5 | 8.6 | 0.90 | 11.88 |
| LP ₄ | 2 | 21.65 | 21.2 | 22.1 | 2 | 18.20 | 17.0 | 19.4 | | | 5 | 13.64 | 12.5 | 15.2 | 1.03 | 7.59 |
| BP ₄ | 2 | 12.55 | 12.4 | 12.7 | 2 | 9.75 | 9.5 | 10.0 | | | 5 | 8.86 | 8.5 | 9.3 | 0.41 | 4.63 |
| LM ₁ | 1 | 24.50 | | | 4 | 22.55 | 19.2 | 25.5 | | | 5 | 16.94 | 14.7 | 18.3 | 1.55 | 9.20 |
| BM ₁ | 1 | 15.50 | | | 4 | 14.17 | 12.8 | 15.7 | | | 5 | 11.48 | 11.0 | 12.2 | 0.57 | 5.02 |
| LM ₂ | 2 | 29.75 | 28.8 | 30.7 | | | | | | | 3 | 18.90 | 18.0 | 19.5 | | |
| BM ₂ | 2 | 16.90 | 16.0 | 17.8 | 1 | 34.0 | | | | | 3 | 12.10 | 11.8 | 12.5 | | |
| LM ₃ | 1 | 38.70 | | | 1 | 14.0 | | | | | 3 | 24.17 | 23.0 | 25.5 | | |
| BM ₃ | 1 | 16.50 | | | 1 | 13.0 | | | | | 3 | 11.73 | 11.0 | 12.5 | | |
| LD ₂ | | | | | 1 | 6.6 | | | | | 1 | 8.2 | | | | |
| BD ₂ | | | | | 2 | 18.1 | 18.0 | 18.2 | | | 1 | 5.0 | | | | |
| LD ₃ | | | | | 1 | 9.0 | | | | | 2 | 12.80 | 12.3 | | 13.3 | |
| BD ₃ | | | | | 3 | 29.67 | 29.0 | 30.0 | | | 2 | 8.25 | 6.1 | | 8.4 | |
| LD ₄ | | | | | 3 | 12.67 | 12.0 | 12.7 | | | 4 | 20.15 | 19.0 | | 21.4 | |
| BD ₄ | | | | | | | | | | | 4 | 9.00 | 7.0 | | 10.0 | |
| LP ₂ -M ₃ | | | | | | | | | | | 1 | 95.0 | | | | |
| LP ₁ | | | | | | | | | | | 1 | 33.2 | | | | |
| LM ₁ | | | | | | | | | | | 1 | 60.0 | | | | |
| LD ₂ -D ₄ | | | | | | | | | | | 1 | 41.8 | | | | |

| HUMERUS | | | D. dama | | | | | | C. elaphus | |
|------------|------|---------|-----------|------|--------|--------|------|-----------|------------|--------|
| C. elaphus | | D. dama | Ph I | | | Ph III | | | McIII+IV | |
| DTdistal | 68.0 | 41.0 | L | 41.0 | | L | 27.7 | 37.0 | DTdia. | (22.7) |
| DAPdistal | 62.0 | 37.8 | DTprox. | 15.5 | H | 20.2 | 20.5 | DAPdia. | (23.3) | |
| DTdistal | 64.0 | 36.5 | DAPprox. | 18.0 | DT | 12.0 | 13.5 | DTdistal | (39.6) | |
| DTtr.max. | 45.0 | 28.0 | DTdia. | 12.5 | Hart. | 14.0 | 17.0 | DAPdistal | (25.1) | |
| DTtr.min. | 32.0 | 17.8 | DTdistal | 14.0 | DTart. | 10.5 | | | | 49.5 |
| DTtr.crest | 35.2 | 23.7 | DAPdistal | 14.0 | | | | | | 32.5 |
| D cond. | 31.0 | | | | | | | | | |
| PATELLA | | | | | | | | | | |
| Lart. | 37.5 | | | | | | | | | |
| DTart. | 25.0 | | | | | | | | | |

| ASTRAGALUS | | | | | |
|--------------|----------|------------|---------------|--------|-----------|
| C. elaphus | | | D. dama (n=2) | | |
| L | 4 | 58.07 | 55.5-61.0 | 35.55 | 32.7-38.4 |
| DT | 4 | 36.00 | 34.5-37.0 | 24.00 | 23.0-20.0 |
| DAP | 4 | 31.62 | 30.0-34.0 | 21.20 | 20.0-22.2 |
| Llat. | 4 | 58.07 | 55.5-61.0 | 38.0 | |
| Lmed. | 2 | 53.30 | 50.6-56.0 | 36.5 | |
| DTprox. | 2 | 32.30 | 32.0-35.4 | 23.0 | |
| DTdistal | 4 | 36.00 | 34.5-37.0 | 24.4 | |
| DAPlat. | 3 | 31.27 | 29.3-34.0 | 21.5 | |
| DAPmed | 4 | 31.37 | 30.0-33.0 | 22.0 | |
| CALCANEUS | | | | | |
| L | 111.0 | L | 29.0 | 32.0 | 22.0 |
| DT | 35.0 | DT | 44.4 | 42.5 | 27.2 |
| DAP | (36.5) | DAP | 39.8 | (34.2) | 24.5 |
| DTtuber | 25.5 | DTar.(as.) | 37.3 | | |
| DAPtuber | 32.0 | | | | |
| DTart.(as.) | 20.0 | | | | |
| DAPart.(as.) | 25.0 | | | | |
| Hart.(cub.) | 23.0 | | | | |
| Bart.(cub.) | 10.0 | | | | |
| | SCAPHOID | PYRAMIDAL | SEMI LUNAR | | |
| L | 26.6 | 30.0 | 24.5 | | |
| DT | 18.7 | 18.5 | 23.5 | | |
| DAP | 33.0 | 27.5 | 30.5 | | |

Table 7. Cervidae-Apidima: Measurements of teeth and postcranial skeleton, in mm.




Figure 12: Ratio-diagram comparing dimensions of teeth of cervids from LAO (Apidima), PEC (Petalona, TSOUKALA, 1989), Megalopolis (MELENTIS, 1965), Camiac, Combe Grenal (GUADELLI, 1987), coll. Reilhac, Museum Nationale Histoire Naturelle-Paris. 1. LP², 2. BP², 3. LP³, 4. BP³, 5. LP⁴, 6. BP⁴, 7. LM¹, 8. BM¹, 9. LM², 10. BM², 11. LM³, 12. BM³, 13. LP₂, 14. BP₂, 15. LP₃, 16. BP₃, 17. LP₄, 18. BP₄, 19. LM₁, 20. BM₁, 21. LM₂, 22. BM₂, 23. LM₃, 24. BM₃. Standard: *D. dama*, recent, Rome (LEONARDI & PETRONIO, 1976).

A robust *D₄* has a very intense and high cingulum between lobes, as well as lingual, anterior and posterior well developed cingulums. Of atlas, the breadth of the anterior articulation is 45 mm; of the posterior one 39.5mm, while its maximum height is 51.5mm and the caudal height of corpus is 23.5 mm. Most of the post cranial bones are calcite covered. Of pelvis, the acetabulum has dimensions: Dmax = 49 and Dmin = 43 mm. Only a distal part of humerus is well preserved, which indicates a robust individual. Of six astragali, one was found on the surface, outside of cave D, while among the others, found in cave C, one is poorly preserved, but robust (L = 52 mm). A calcaneus is slender but high, and lacks the anterior articulation for malleolus. The sustentaculum tali is long but not high, with a posterior projection. The body is +/- constant antero-posteriorly, the articulation for astragalus +/- rectangular and there is a quite deep sulcus posterior of tuber. The cuboscaphoidea, patella and PhI (one of a juvenile) are well preserved, while the distal part of a metacarpal is poorly preserved, with cutting and rolling marks. In addition to recent specimens, many more tooth and bone fragments, mainly from Cave C (LAO 309) have been identified as cervid remains (the minimum number of individuals of which is 5).

Genus *Dama* FRISCH, 1775

Dama dama (LINNAEUS, 1758)

(Plate 1, Text figs. 10–12, Table 7)

M a t e r i a l 14 Maxilla frs. with P²-M³ LAO-C-520b sin, with P²-M³ (M³ 705b) and 2M^{1,2} 706d, with P²-M¹ 771, with P³-M³ 527, with P⁴-M³ 520c sin, with P⁴-M² 634 dex, with D²-M² 576a dex, with D²-D⁴, with D²,D³ 579a dex, with D³-M² 644d, with D⁴-M² and P⁴ dex, LAO-C-700b, with D²-D⁴ and 2M¹ LAO-C-726a and 708a, with D³, D⁴ 643d, 505 sin, P² and P³ LAO-C-529b, 4P² LAO-C-573c and two 588b dex & sin, LAO-B-139d, P³ LAO-B-184d, P⁴ LAO-A-17c sin, P⁴ and M¹ LAO-C-653a, M¹ LAO-C-558h, 2M² LAO-C-655 and 709 sin, M^{1,2} LAO-C-700b, 2M³ LAO-A-34, LAO-C-638a sin, 2D² LAO-C-544c, 701a, 13mandible frs with P₂-M₃ LAO-C-565 dex, with P₃-M₃ LAO-B-171a, with P₄-M₃ LAO-C-593 sin, with P₃-M₁ 672, with M₁-M₃ 579, 2with D₄-M₂ 577 and 639 dex, 2with M₃ 568, 572a dex, 2with D₂-D₄ 549, 306 sin, with D₃,D₄ 594a dex, with D₂ LAO-C-557c, I₁ LAO-C-280f, 554a, 571b, 622c, 633, 644d, 4P₂ LAO-C-614b, 744c, 810d, 814a, dex, 3P₃ LAO-C-698b, 725e, 746c, P₄ LAO-A-101c dex, 2M₁ LAO-C-547b, 701a, M₂ LAO-C-649c dex, 2D₄ LAO-B-162d, LAO-C-531b, 2 pelvis frs LAO-C-580, 618, 2 humeri distal LAO-C-703c sin, 2 astragali LAO-C-562, 717 dex, 3cuboscaphoidea LAO-

C-633, 700d, 810c sin, astragalus-calcaneus-cuboscaphoid. juv.LAO-C-644, 644a, Mc dis.LAO-B-87a, Mp fr. LAO- A-05, 4PhI LAO-C-488b, 531a, 626, 805d, 2PhII LAO-C-605, 643f, 6PhIII LAO-C-533a, 603, 610a, 622, 622c, 633a.

Description: The teeth vary in worn stage and have very plicated and thin enamel and, in some cases, a very high and strong cingulum between lobes, as well as closed fossets. On some molars (M^1 LAO-B-22, M^3 LAO-C-638a etc.) there is a well developed and high (7.4 mm) cingulum between lobes. Mandible fragments are preserved in consolidated breccia. The lower molars have small external cingulums, the M_2 have anterior and between lobes intense cingulums, while the M_3 have small or no cingulums between lobes. At the base of a D_4 there are two external cingulums between the lobes. Of the post-cranial skeleton, only pelvis acetabulums ($D = 33$ and 37 mm), two distal humeri (one is of a juvenile bearing trace of carnivore gnawing), astragali, calcaneus and cuboscaphoideum, mainly of juveniles, a distal metapodial ($DT_{distal} = 29$ mm) (Fig. 10.5, Pl. 1.14) and phalanges are well preserved and most representative. It is rather smaller in size than the fallow deer from Petralona cave (Fig. 12).

The fallow deer comes from Eemian (KURTEN, 1968), it is known during historical times, among other Mediterranean sites, in Crete island, while today very few representatives are living in Rhodes island. Late Pleistocene representatives are referred in Kythera island (MANOLESSOS, 1955) and Vraona cave, Attiki (RABEDER, 1995).

Suborder SUIFORMES JAECKEL, 1911
Family Hippopotamidae GRAY, 1821
Genus *Hippopotamus* LINNAEUS, 1758

Hippopotamus amphibius antiquus
DESMAREST, 1822
(Text-fig. 13, Table 8)

Material C^s LAO -A-117 sin and Ph II LAO-B-175.

Description A fragment of upper canine is partly calcite-covered and poorly preserved, while the very well preserved second phalanx, corresponds to the second or fifth anterior metapodial, according to the index $Ia = (DAP_{pr.art.}/DT_{pr.art.} \times 100) = 66.87$.




Figure 13: *Hippopotamus amphibius antiquus*-Apidima. Above: C^s LAO -A-117 sin, a. labial, b. distal, c. mesial. (AA; cross section) Below: Ph II LAO 175, a. anterior, b. posterior.

Discussion: The dimensions of the hippopotamid remains found in caves B and A (where the two hominid skulls have been found), are relatively small, therefore may belong to a smaller individual, such as a young female, but not to a dwarf type, although dwarf endemic races of hippopotamids evolved in several Mediterranean islands. The hippopotamids are widely distributed, either in mainland or in the Greek islands (SYMEONIDIS & THEODORU, 1985/86), while only few fossil localities have been recorded in the south-east European and transcaucasian area (KAHLKE, 1987, 1989). On the other hand *Hippopotamus amphibius antiquus* DESMAREST, 1822, or *H. a. incognitus* FAURE, 1984, has been identified at many (~350) fossiliferous sites in western Europe (FAURE, 1985). In Peloponnese (S. Greece) this species has been identified at several sites, such as Dyros cave (SYMEONIDIS & THEODOROU, 1985/86); Kleitoria (MELENTIS, 1969); Megalopolis, of Holsteinian age (MELENTIS, 1964, 1965b), in a paleoenvironment with rich vegetation; Marathousa (Megalopolis basin) of Early Biharian (SICKENBERG, 1976), with main associated fauna *Mammuthus meridionalis* and *Praemegaceros verticornis*. The paleoenvironment was open landscape with many lakes, low moors, marshy forests in a humid, but not very warm climate. *H. ex. aff. amphibius major* has been recorded in Elis (THENIUS, 1955) of Astian (Middle Villafranchian) age and in central Macedonia of Villaynian age (KOUFOS et al., 1989). Hippopotamid remains have been identified also at Thessaly (Pinios river, BOESSNECK, 1965); Aliakmon river (MELENTIS, 1966); Euboia; Kos; Kephallonia; Agios Demetrios-Kato Salmeniko (SYMEONIDIS & THEODOROU, 1985/86); Neapolis (STEENSMA, 1988); Ptolemaida etc.

Table 8. *Hippopotamus amphibius antiquus*-Apidima: Measurements of the upper canine and the second phalanx, in mm.

| C ^s | PhII |
|----------------|------------------|
| L 8.4 | L 33.5 |
| B 25.3 | DT prox. 34.3 |
| | DAP prox. 21.8 |
| | DT pr.art. 31.7 |
| | DAP pr.art. 21.2 |
| | DAP d.art. 20.0 |

Order Proboscidea ILLIGER, 1811
 Family Elephantidae GRAY, 1821
 (Text-fig. 14, Table 9)

M a t e r i a l: Ph II LAO-C-282e

D e s c r i p t i o n A rather short but robust second phalanx is the unique elephant remain. It is much higher posteriorly than anteriorly, while the proximal articular surface is relatively very wide and concave, without medial crest and may belong to *Elephas (Paleoloxodon) antiquus* FALCONER & CAUTLEY (1847).

Discussion

The rich but very fragmentary material from the Apidima site is not generally preserved in good condition, and is therefore mostly indeterminable. On the other hand the determinations were based on the few but very well preserved specimens. There are well-fossilized specimens, mainly in breccia, as well as recent bones e.g. goats, leporids, birds. There are many burnt and calcite- or salt- covered bones. The action of water is obviously indicated on some specimens, either by the erosion or by the rounding and abrasion of bones and the scattered skeletons.

- Cave C is the most representative in faunal composition, as only hippopotamid remains have not been found up to now. This may have to do with the altitude of the cave (Fig. 1), as the high sea-waves have moved most material (fissure fillings or cone of debris) of the other caves away. The presence of many cervid remains is notable. The presence of many micromammals (rodents), birds (mostly




Figure 14: Elephantidae-Apidima, second phalanx, Ph II, LAO-C-282e, anterior.

Ph II

| | | |
|----|---------------|------|
| 1. | Lmax | 38.0 |
| 2. | Lant. | 30.5 |
| 3. | DT max. | 51.0 |
| 4. | DT prox. | 50.7 |
| 5. | DAP prox. | 32.0 |
| 6. | DT prox.art. | 48.5 |
| 7. | DAP prox.art. | 29.3 |
| 8. | DT d.art. | 41.6 |
| 9. | DAP d.art. | 20.4 |

Table 9.
 Measurements of
 the Apidima
 elephant second
 phalanx, in mm

| Taxon / Cave | A | B | C | D |
|--|---|---|---|---|
| <i>Panthera pardus</i> | - | + | + | - |
| <i>Felis(Lynx) lynx</i> | - | + | + | + |
| <i>Felis silvestris</i> | - | - | + | - |
| <i>Vulpes vulpes</i> | + | + | + | + |
| <i>Meles meles</i> | - | ? | + | - |
| <i>Martes foina</i> | - | - | + | - |
| <i>Capra ibex</i> | + | + | + | + |
| <i>Megaloceros</i> sp. | - | - | + | - |
| <i>Cervus elaphus</i> | - | + | + | + |
| <i>Dama dama</i> | + | + | + | ? |
| <i>Hippopotamus amphibius antiquus</i> | + | + | - | - |
| Elephantidae | - | - | + | - |

Table 10. Representation of Quaternary large mammal remains in the four Apidima caves

beaks), leporids and burnt bones is also notable. There are also many turtle bones. Some bones are covered by a brownish-reddish calcite layer. There are a few specimens with probable traces of butchering.

The cave B faunal remains do not include wild cat, marten, giant deer or elephant, while the badger is not well represented. Of the very fragmentary material, mostly indeterminable, some bones are eroded and some are well fossilized, preserved in breccia. The majority of the material is represented by a mixture of caprid and cervid bones and teeth, especially the latter (incisors included). The presence of leopard remains is notable. Surface finds include small felid, fox, goat and deer remains. The cave D faunal remains include lynx, red fox, wild goat and red deer, while fallow deer is poorly represented. There are bones with carnivore tooth traces, carnivore milk teeth, many birds of galliform size, burnt bones, bones which are rounded or covered with a calcite and/or salty layer, and recent bones, mainly of Caprinae.

The cave A faunal remains include very few representatives, mainly associated with the two hominid skulls, such as the red fox, wild goat, fallow deer and hippopotamus. Some specimens found in breccia on the floor, are eroded, rounded or burnt. The hippopotamus is found in association with rodents, birds and some indeterminable bones in breccia. The material from depths of about 0.70-0.94m consists mainly of artiodactyls, a few burnt bones, and rodents.

Conclusions

A study of the determinable and the most representative large mammal material, among 20.000 teeth, bones and bone fragments from the Apidima caves, has shown the presence of the following species:

- Panthera pardus* (LINNAEUS, 1758)
Felis (Lynx) lynx (LINNAEUS, 1758)

Felis silvestris SCHREBER, 1777

Vulpes vulpes (LINNAEUS, 1758)

Meles meles LINNAEUS, 1758

Capra ibex LINNAEUS, 1758

Megaloceros sp.

Cervus elaphus LINNAEUS, 1758

Dama dama (LINNAEUS, 1758)

Hippopotamus amphibius antiquus DESMAREST, 1822

Elephantidae

In addition to these, there are recent specimens which belong to *Felis*, *Martes*, *Meles*, *Capra*, cervids, leporids, rodents and birds.

The presence of fossil and recent bones in this mixed fauna shows that the caves were used during different periods, from Middle Pleistocene to present, and this has to do with the eustatic sea-level fluctuations before today's transgression, which eroded almost all the cave A fillings.

The majority of the finds belongs to herbivores - mainly artiodactyls-which, except for their presence by chance, were the main food of the carnivores - mainly felids- that sometimes inhabited the caves. Traces of carnivore gnawing on herbivore bones, especially on juvenile caprid and cervid bones, indicate the food remains of the carnivores.

The juvenile carnivore remains, as well as the food remains, establish them as inhabitants.

The absence of perissodactyl, suid, ursid etc. remains is notable, although a richer fauna could be expected at this, southernmost continental site of Europe, which might have worked as a natural coastal barrier.

The presence of many micromammalian remains, mainly murid and cricetid rodents (mice and voles), as well as insectivores (mostly soricids), birds, reptiles (mostly turtles), and fish (represented by vertebrae) is notable.

The presence of many leporid, rodent and bird remains is consistent with the presence of the foxes and felids.

Some specimens show traces of butchering, and the habitation of man is also established by skeletal remains, artefacts, burials, fire traces etc..

The age of the fossil fauna, according to the stratigraphical distribution of the large mammals (the carnivore distribution was also based on WOLSAN (1993) and cervid distribution on LISTER (1986)), is Middle to upper Middle Pleistocene, and Late Pleistocene.

The climate was temperate, rather mild (not "cold") although in other parts of Europe there were glacial episodes with cold climate. The favourable climatic conditions in combination with the increased land area, because of the lowering of sea-level, helped the immigration and settling of animals and humans. It is an important feature of the Greek

Quaternary faunas, that species typical of cold climate (except very few and isolated specimens), are missing up to now. The few extensive studies of Quaternary sites (TSOUKALA, 1992b) do not show faunas typical of cold episodes, such as in rest Europe. Further studies would bring to light valuable conclusions.

The paleoenvironment of the broader area of Mani peninsula, during Pleistocene, was mixed and variable steppe like during sea regression, interrupted by forests (mainly of coniferous trees, wild olive-trees, pistachio and other Mediterranean flora). The mixed forests and rocky hillsides are favourable to certain species such as *Martes foina*. As the Apidima fauna is stratigraphically mixed, careful interpretation is required.

Acknowledgements

Deepest thanks to Prof. Theodore PITSIOS (Anthropological Museum of Athens University) for trusting me the study of this important Apidima material. Deep thanks to Prof. S. Verginis and Prof. G. Rabeder (Vienna University) as well as to Prof. Henry de Lumley, Director of the French National Museum of Natural History, for their invitation, so I had the chance to see, study and compare various Quaternary faunas (IPH-Institut Paleontologie Humaine-Paris, Lazaret, Vallonet, Arago, Portel etc). Also many thanks to Dr. Anne-Marie Moigne, Dr. F. Spitzerberger and Dr. G. Hoeck-Daxner. To Dr. A. Lister for the linguistic corrections my sincerest thanks. Special deep thanks to Prof. W.v. Koenigswald and Dr. D. Nagel for the critical reading of the manuscript.

References

- ALTUNA, J., 1972. Fauna de mamíferos de los yacimientos prehistóricos de Guinuzcoa. — Thes. Doct. Munibe (1971), **XIV**(1-4):1-464, 72 fig., 28 pl., San Sebastian.
- ANDERSON, E., 1970. Quaternary evolution of the genus *Martes* (Carnivora, Mustelidae). — Acta zoologica fennica, **130**:1-132, 151 figs., 38 tabl., Helsinki.
- ARGANT, A. & J., 1990. Blanot (Sâone-et-Loire): Gisement paléontologique de pléistocène supérieur ancien (Riss-Würm) en Mâconnaisse. — "La Physiophile", **113**:72-96.
- ARGANT, A., 1991. Carnivores Quaternaires de Bourgogne. — Docum. Lab. Géol. Lyon, **115**:1-301, 39 fig., 89 tabl., 9 pl., Lyon.
- BALLESIO, R., 1979. Le gisement pléistocène supérieur de la grotte de Jaurens à Nespouls, Corrèze, France: Les carnivores (Mammalia, Carnivora). I. Canidae et Hyaenidae. — Nouv. Arch. Mus. Hist. Nat. Lyon, **17**:22-55, 17f., 5 pl., Lyon.
- BALLESIO, R., 1980. Le gisement pléistocène supérieur de la grotte de Jaurens à Nespouls, Corrèze, France: Les carnivores (Mammalia, Carnivora). II. Felidae. —

- Nouv. Arch. Mus. Hist. Nat. Lyon, **17**:61–102, 15f., 4 pl., Lyon.
- BASSIAKOS, J.E., 1993. Dating of fossils from caves and speleothems: evidence from electron spin resonance (E.S.R.) technique, the study of underground karst morphology and the relevant radiometric and geological conditions in speleoenvironments of Dyros, Mani. — Doctorate Thesis submitted in School of Geology, Athens University: 1–380, 82 figs., 23 tabls., 35 pls., Athens (in Greek with English summary).
- BASSIAKOS, J.E. & TSOUKALA, E., 1996. ESR dating suitability of Quaternary fossil remains; a hyaenid tooth example and new data on the fauna from Agios Georgios Cave (Kilkis, Macedonia). — Proceedings 2nd Symposium of the Hellenic Archaeometrical Society (26–28 March 1993), Thessaloniki.
- BARTSIOKAS, A., 1998. The Paleontology of Cythera island. — Society of Cytherian Studies, 1–96, 73 pls., Athens (in Greek).
- BOESSNECK, J., 1965. Die Jungpleistozänen Tierknochenfunde aus dem Peneiostal bei Larissa in Thessalien. — [in:] MILOJZIC, V.: Paläolithicum um Larissa in Thessalien. — Beitr. ur-frühgesch. Archaeol. Mitteleuropa-Kulturraumes, (R.H. Verlag).
- BONIFAY, M.-F., 1971. Carnivores quaternaires du Sud-Est de la France. — Mémoires de Musée National Histoire naturelle de Paris, Series C, **21** (2):43–377, Paris.
- COUTSELINIS, A., DRITSAS, C. & PITSIOS, Th., 1991. Expertise medico-legale sur crâne pléistocène LAO1/S2 (Apidima II), Apidima, Laconie, Grèce. — L' Anthropologie, **95** (2/3):401–408, Paris.
- CREGUT, E., 1976. La faune de mammifères du pléistocène moyen de la Caune de l'Arago à Tautavel (Pyrénées Orientales). — Trav. Lab. Paléont. Humaine et Préhist., **3**:1–381. Marseille.
- FAURE, M., 1985. Les hippopotames quaternaires non-insulaires d'Europe occidentale. — Nouv. Arch. Mus. Hist. nat. Lyon, **23**:13–79, Lyon.
- GERBER, J.P., 1973. La faune de grands mammifères du Würm ancien dans le sud-est de la France. — Thèse Univers. Provence. Travaux du Laboratoire de Géologie historique et Paléontologie **5**:1–310, Marseille.
- GUADELLI, J.-L., 1987. Contribution à l'étude des zoocénoses préhistoriques en Aquitaine (Würm ancien et interstade Würmien). — Thèse doctor. a l' Univ. Bordeaux **3**:1–568, 1–450, 1–427, Bordeaux.
- HEMMER, H., 1971. Zur Kenntnis pleistozäner mittel-europäischer Leoparden (*Panthera pardus*). — Neues Jb. Geol. Paläont. Abh., **138**(1):15–36, Stuttgart.
- KAHLKE, R.-D., 1987. On the occurrence of *Hippopotamus* (Mammalia, Artiodactyla) in the Pleistocene of Achalkalaki (Gruzinian SSR, Soviet Union) and on the distribution of the genus in Southeast Europe. — Z. geol. Wiss., **15**: 407–414, Berlin.
- KAHLKE, R.-D., 1989. Die unterpleistozänen *Hippopotamus*-Reste von Würzburg – Schalksberg. — Quartär, **39/40**: 67–94 Bonn.
- KOUFOS, G., 1981. A new Late Pleistocene (Würmian) mammal locality from the basin of Drama (N. Greece). — Scientific Annals, Faculty Physics & Maths, Aristotle University, **21**:129–148 Thessaloniki.
- KOUFOS, G., SYRIDES, G. & KOLIADIMOU, K., 1989. A new Pleistocene mammal locality from Macedonia (Greece). Contribution to the study of Villafranchian (Villangian) in central Macedonia. — Bull. Geol. Soc. Greece, **XXIII/2**: 113–124, Athens.
- KURTEN, B., 1965. On the evolution of the European Wild Cat, *Felis silvestris* SCHREBER. — Acta zoologica fennica, **111**:1–29, Helsinki.
- KURTEN, B., 1965a. The Carnivora of the Palestine Caves. — Acta zoologica fennica, **107**:1–74, Helsinki.
- KURTEN, B., 1968. Pleistocene Mammals of Europe. — Weidenfeld & Nicolson, 1–316, London.
- LAX, E., 1995. Quaternary faunal remains from the Cave Site of Apidima (Lakonia, Greece). — Acta Anthropologica, **1**:127–156. Proceedings Meeting Research Programme of Mani peninsula, 1978–1988, Athens.
- LIRITZIS, Y. & MANIATIS, Y., 1989. ESR experiments on Quaternary calcites and bones for dating purposes. — J. Radioanalytical and Nuclear Chemistry, Articles, **129**(1):3–21, Akad. Kiadó, Budapest (Elsevier Sequoia S.A. Lausanne).
- LIRITZIS, Y. & MANIATIS, Y., 1995. Acta Anthropologica, **1**: 65–84 (in Greek) and 85–92 (in English). — Proceedings of the Meeting on the Research Programme of Mani peninsula, 1978–1988, Athens.
- LISTER, A., 1986. New results on deer from Swanscombe, and the stratigraphical significance of deer in the Middle and Upper Pleistocene of Europe. — Journal Archaeological Science, **13**:319–388.
- LEONARDI, G. & PETRONIO, C., 1976. The fallow deer of European Pleistocene. — Geologica Romana, **15**:1–67, 56 fig, 2 tabs., 8 pls., Roma.
- LUMEY, de H. & DARLAS, A., 1994. Grotte de Kalamakia (Areopolis, Péloponnèse). — Bulletin de Correspondance Hellénique (BCH), **118**:535–559, Athens.
- MANOLESSOS, N.J., 1955. Beiträge zur Kenntnis der Geologie der Insel Kythera. — Doctorate Thesis, Ann. Géol. Pays Helléniques, **6**:51–80, 3 figs., 7 pls., Athens.
- MELENTIS, J., 1964. Die fossilen Rhinocerotiden, Hippopotamiden und andere Säugetiere aus dem Becken von Megalopolis im Peloponnes (Griech.). — Praktika Acad. Athinon, **39**:388–400, 1 pl., Athens.

- MELENTIS, J., 1965a. Die pleistozänen Cerviden des Beckens von Megalopolis im Peloponnes (Griechenland). — Ann. Géol. Pays Helléniques, **16**:1–92, 12 pls., Athens.
- MELENTIS, J., 1965b. Studien über fossile Vertebraten Griechenlands: 5. über *Hippopotamus antiquus* DESMARET aus dem Mittelpaläozän des Beckens von Megalopolis im Peloponnes (Griechenland). — Ann. Géol. Pays Helléniques, **16**:403–435, 3 pls., Athens.
- MELENTIS, J., 1966. Studien über fossile Vertebraten Griechenlands: 16. Die Pleistozäne Säugetierfauna des Beckens von Haliakmon (Griech.). — Ann. Géol. Pays Helléniques, **17**:247–266, 2 pls., Athens.
- MELENTIS, J., 1969. Die Quartären Vertebraten der Höhle der Seen von Kleitoria (im Gebiet der Aroania-Gebirge). — Prakt. Acad. Athinon **43**:350–363, 1 pl., Athens.
- NAGEL, D., 1995. Die Felidenreste aus dem Jungpleistozän von Vraona-Griechenland. — [in:] SYMEONIDIS, N. & RABEDER, G. (eds.). "Das Jungpleistozän in der "Höhle" von Vraona auf Attica in Griechenland" — Ann. Géol. Pays Helléniques, **36**(1993–95): 60–100, pl. 1, Athens.
- NAGEL, D., 1999. *Panthera pardus* vraonensis n. ssp., a new leopard from the Pleistocene of Vraona/Greece. — N. Jb. Geol. Paläont. Mh. **1999**(3):129–150, 5 figs., 4 tabs., Stuttgart.
- PITSIOS, Th., 1979. Paleoanthropological finds of Inner Mani. — Anthropos, **6**:98–105; Athens (in Greek).
- PITSIOS, Th., 1983. Paleoanthropologische Funde in Mittel Mani II. — Proceedings of 1st Local Meeting for Laconic Research (Molai, 6/1982) of Society of Peloponnesian Studies: 77–83, Athens (in Greek with German abstract).
- PITSIOS, Th., 1985a. Pleistocene fluctuation of Mediterranean Sea and contribution of this method to the datation of paleoanthropological finds of Inner Mani II. — Acta Anthropologica, **8**:23–32, Volos–Thessaloniki.
- PITSIOS, Th., 1985b. Paleoanthropologic research on the site "Apidima" Mesa Mani II. — Archaeology, **15**:26–33, Athens. (in Greek).
- PITSIOS, Th., 1996. The Taenarius Man. — Archaeologia **60**: 68–72, Athens.
- PITSIOS, Th. & LIEBHABER, B., 1995. Research conducted in Apidima and the surrounding region- Taenarius Man. — Acta Anthropologica, **1**:169–179. Proceedings of the Meeting on the Research Programme of Mani peninsula, 1978–1988, Athens.
- RABEDER, G., 1995. Jungpleistozäne und Frühholozäne Säugetierreste aus der Höhle von Vraona auf Attika, Griechenland. — [in:] SYMEONIDIS, N. & RABEDER, G. (eds.). "Das Jungpleistozän in der "Höhle" von Vraona auf Attica in Griechenland" — Ann. Géol. Pays Helléniques, **36**(1993–95): 47–58, Athens.
- RONDYANNI, Th., METTOS, A. & GEORGIU, Ch., 1995. Geological-morphological observations in the greater Oitilo–Diros area, Mani. — Acta Anthropologica, **1**:93–103. Proceedings of the Meeting on the Research Programme of Mani peninsula, 1978–1988, Athens.
- SICKENBERG, O., 1976. Eine Säugetierfauna des tieferen Bihariums aus dem Becken von Megalopolis (Peloponnes, Griechenland). — Ann. Géol. Pays Helléniques, **27**:25–73, 5 pls, Athens.
- STEENSMÅ, K.J., 1988. Plio-Pleistozäne Grosssaugtiere (Mammalia) aus dem Becken von Kastoria/Grevena, südlich von Neapolis-NW-Griechenland. — Dissertation, Math.-Naturwiss. Fak. Tech. Univ. Clausthal, 315 p., Clausthal.
- SYMEONIDIS, N. & THEODOROU, G., 1985/86a. New locations of fossil *Hippopotamus* on Northwest Peloponnese. — Ann. Géol. Pays Helléniques, **33**:51–67, 3 pls, Athens (in Greek).
- SYMEONIDIS, N., BACHMAYER, F. & ZAPFE, H., 1980. Ergebnisse weiterer Grabungen in der Höhle von Vraona (Attika, Griechenland). — Annales Géologiques des Pays Helléniques, **30**: 291–299, Athens.
- THENIUS, E., 1955. Studien über fossile Vertebraten Griechenlands. V. *Hippopotamus* aus dem Astien von Elis (Peloponnes). — Ann. Géol. Pays Helléniques, **6**: 206–212; Athens.
- THENIUS, E., 1956–59. Die Jungpleistozäne Wirbeltierfauna von Willendorf i. d. Wachau N.Ö. Österr. — Akad. Wiss., **VIII/IX**:133–170, Wien.
- THENIUS, E., 1972. Die Feliden (Carnivora) aus dem Pleistozän von Stranska Skala. — "Anthropos", 121–135, Vyjava, Brne.
- TSOUKALA, E., 1989. Contribution to the study of the Pleistocene fauna of large mammals (Carnivora, Perissodactyla, Artiodactyla) from Petralona Cave (Chalkidiki, N. Greece). — Doct. Thesis, Sci. Ann., Sch. Geology, Thes. Univ., **1**(8):1–360, 124 figs., 62 pls., 64 tabs., Thessaloniki (in Greek with English summary). — 1991, C.R.A.S. Paris **312** (II): 331–336.
- TSOUKALA, E., 1992a. The Pleistocene large mammals from the Agios Georgios Cave, Kilkis (Macedonia, N. Greece). — Geobios: **25**(3):415–433, 16 figs., 9 tabs., 3 pls., Lyon.
- TSOUKALA, E., 1992b. Quaternary faunas of Greece. — Courier Forsch. Inst. Seckenberg, **153**:79–92, 2 tabl., Frankfurt a.M.
- VERGINIS, S., VERICIOU, E. & PSARIANOS, P., 1975. New occurrences of Neogene deposits at Itylo area (W. Laconia) and contemporaneous continental movements of Laconic Peninsula. — Praktika Acad. Athinon, **50**:443–453, Athens (in Greek).
- WOLSAN, M., 1993. Evolution des carnivores Quaternaires en Europe centrale dans leur contexte stratigraphique et paleoclimatique. — L' Anthropologie, **97**(2/3): 203–222.

PLATE 1

Panthera pardus (LINNAEUS, 1758)

- Fig. 1. Maxilla fragment LAO-B-371a sin, occlusal (~1:1)
Fig. 2. C^s LAO-B-91dex, a) labial, b) lingual, c) posterior (~7.5:10)
Fig. 3. C^s fragment LAO-B-409dex, labial (~1:1)
Fig. 4. Radius LAO-B-388c+389sin, a) anterior, b) posterior (~6:10)
Fig. 5. Navicularis LAO-B-389b sin, a) proximal, b) anterior, c) posterior (~6:10)
Fig. 6. McV LAO-B-389a dex, a) dorsal, b) plantar, c) medial (~6:10)
Fig. 7. PhI LAO-B-410a, dorsal (~8:10)
Fig. 8. PhII LAO-B-348e, dorsal (~1:1)

Felis (Lynx) lynx (LINNAEUS, 1758)

- Fig. 9. C^s (~1:1)
- Vulpes vulpes* (LINNAEUS, 1758)
- Fig. 10. Mandible frag. with M₁, M₂ dex LAO-B-352c,353 dex
a) labial, b) occlusal, c) M₁ lingual (~1:1)
- Fig. 11. Tibia LAO-C-402a+404 sin (~7:10)

Capra ibex (LINNAEUS, 1758)

- Fig. 12. Horn fr. LAO-B-79 (~7:10)
- Fig. 13. Metacarpal (Mc III+IV) LAO-B-385 sin (~7:10)

Dama dama (LINNAEUS, 1758)

- Fig. 14. Metacarpal distal, LAO-B-87a, anterior (~7:10)

PLATE 1




PLATE 2

Capra ibex (LINNAEUS, 1758)

Fig. 1. Maxilla frag. with P⁴-M³ LAO-C-24 dex

Fig. 2. Maxilla frag. with P³-M³ LAO-C-26a sin

Fig. 3. Mandible frag. with P₄-M₃ LAO-D-18a sin

Fig. 4. Mandible frag. with P₄-M₃ LAO-D-79e dex


Cervus elaphus (LINNAEUS, 1758)

Fig. 5. Maxilla frag. with P²-M³ LAO-C-27dex.

a) labial, b) occlusal.

All Figures ~7.5:10

PLATE 2



ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Beiträge zur Paläontologie](#)

Jahr/Year: 1999

Band/Volume: [24](#)

Autor(en)/Author(s): Tsoukala Evangelia

Artikel/Article: [Quaternary large mammals from the Apidima Caves
\(Lakonia, S Peloponnese, Greece\) 207-229](#)