CONTRIBUTIONS TO ENTOMOLOGY

© 2022 Senckenberg Gesellschaft für Naturforschung

72 (2): 213–253 2022

SENCKENBERG

Supplement to the Staphylinidae fauna of Crete and the Aegean Islands, Greece (Coleoptera)

With 71 figures and 3 tables

Volker Assing[†]

[†]Volker Assing died completely unforeseen after sending the last corrections to this article. Unfortunately, he was no longer able to confirm their implementation. The editorial team hopes to have implemented everything in his will. We mourn the loss of an outstanding taxonomist and esteemed colleague.

Published on 2022–12–05

DOI: 10.3897/contrib.entomol.72.e97264

Abstract

Recent field trips to Crete, Samos, and Rhodos, as well as a revision of other material from various collections yielded numerous additional records, species, and taxonomic changes for the Staphylinidae faunas of Crete and the Aegean Islands. Fifteen species are described and illustrated for the first time, two of the Pselaphinae, two of the Oxytelinae, two of the Aleocharinae, one of the Leptotyphlinae, and eight of the Scydmaeninae: Paramaurops creticus BRACHAT spec. nov. (Crete: Psiloritis range); Afropselaphus chanianus BRACHAT spec. nov. (Crete); Euphanias ambulans Assing spec. nov. (Samos); Bledius (Hesperophilus) bedelianus SCHÜLKE spec. nov. (Samos, Rhodos, Karpathos, South Turkey, Lebanon, Israel); Hydrosmecta cultellata Assing spec. nov. (Samos); Atheta (Philhygra) tecta Assing spec. nov. (Samothraki, Samos, Kos, Turkey); Cretotyphlus kerkisicus Assıng spec. nov. (Samos); Cephennium amplexans Assıng spec. nov. (Samos); C. monstrans Assing spec. nov. (Samos); C. icariae Assing spec. nov. (Ikaría); C. rhodicum ASSING spec. nov. (Rhodos); Stenichnus (Stenichnus) samius MEYBOHM spec. nov. (Samos); S. (S.) amphimykalicus MEYBOHM spec. nov. (Samos, Southwest Turkey); Scydmoraphes amphimykalicus MEYBOHM spec. nov. (Samos, Southwest Turkey); Euconnus (Tetramelus) rhodicus MEYBOHM spec. nov. (Rhodos). Three new synonymies are proposed: Throbalium cycladicum (Косн, 1937) = T. biblicum (Косн, 1937), syn. nov., = T. adanense Косн, 1939, syn. nov., = Throbalium kosianum BORDONI, 2020, syn. nov. Two new combinations are proposed: Tychus kerkisicus (BRACHAT, 2017), comb. nov. (ex Paratychus BESUCHET, 1960) and Liogluta aloconotoides (BENICK, 1940), comb. nov. (ex Aloconota THOMSON, 1858). Several previous misidentifications are rectified. Numerous species are reported from Crete and the Aegean Islands for the first time, some even from Greece and from Europe. New checklists of the Staphylinidae of Samos and Rhodos are provided. Including the additions and changes, the faunas of Crete, Samos, and Rhodos are now represented by 403 identified named species (114 of them endemic), 199 named species (15 endemic), and 160 named species (10 endemic), respectively.

Taxonomic acts

Paramaurops creticus BRACHAT spec. nov. – urn:lsid:zoobank.org:act:0ADD6720-1EEC-4E67-90EE-F1DC5B1C9ACE Afropselaphus chanianus BRACHAT spec. nov. – urn:lsid:zoobank.org:act:7648FB38-4026-414C-8609-32447D514299 Euphanias ambulans Assing spec. nov. – urn:lsid:zoobank.org:act:4D363881-A549-4AA0-A824-66E2EBA02BD3 Bledius bedelianus SCHÜLKE spec. nov. – urn:lsid:zoobank.org:act:84848C14-D6BD-41DF-9D76-EFDDC83FE7CD Hydrosmecta cultellata Assing spec. nov. – urn:lsid:zoobank.org:act:014960F5-7A69-4D33-8B6F-2F543D07AF7D Atheta tecta Assıng spec. nov. – urn:lsid:zoobank.org:act:2794087C-8BD2-4278-A358-D0CE8A62C1DB Cretotyphlus kerkisicus Assıng spec. nov. – urn:lsid:zoobank.org:act:16BD4F7A-B7A6-47B3-8025-342CAD1855EF Cephennium amplexans Assıng spec. nov. – urn:lsid:zoobank.org:act:41A5DF62-9FFC-44D3-A57C-20B2B93720B9 Cephennium monstrans Assıng spec. nov. – urn:lsid:zoobank.org:act:86CB13EF-3F05-45A1-BB47-034707B226A5 Cephennium icariae Assıng spec. nov. – urn:lsid:zoobank.org:act:738FF668-DCED-4E48-808C-0DE0D67797CA Cephennium rhodicum Assıng spec. nov. – urn:lsid:zoobank.org:act:63142283-CD20-46E9-AB60-9F7AFA6DE29A Stenichnus samius MEYBOHM spec. nov. – urn:lsid:zoobank.org:act:D281CE0D-EEA6-45E7-BCB2-21734004DD80 Stenichnus amphimykalicus MEYBOHM spec. nov. – urn:lsid:zoobank.org:act:BC1B8722-6DC0-40C0-BC10-9A2E-673C9E4E

Scydmoraphes amphimykalicus MeyBOHM spec. nov. – urn:lsid:zoobank.org:act:2CDA0C85-BA74-46EF-8AA6-ADAFB1974DC2

Euconnus rhodicus MeyBOHM spec. nov. - urn:lsid:zoobank.org;act:A286F3E4-9AC8-434F-B473-96E388B238A8

Key words

Coleoptera, Staphylinidae, Palaearctic region, East Mediterranean, Greece, Crete, Aegean Islands, taxonomy, endemism, endogean fauna, new species, new synonymies, new combinations, additional records, checklists

Zusammenfassung

In den letzten Jahren durchgeführte Forschungsreisen nach Kreta, Samos und Rhodos sowie eine Revision von weiterem Material aus mehreren Sammlungen ergaben zahlreiche weitere Nachweise, Arten und taxonomische Änderungen für die Staphylinidenfauna Kretas und der ägäischen Inseln. Fünfzehn Arten aus den Unterfamilien Pselaphinae (2 Arten), Oxytelinae (2), Aleocharinae (2), Leptotyphlinae (1) und Scydmaeninae (8) werden erstmals beschrieben und abgebildet: Paramaurops creticus BRACHAT spec. nov. (Kreta: Psiloritis); Afropselaphus chanianus BRACHAT spec. nov. (Kreta); Euphanias ambulans Assing spec. nov. (Samos); Bledius (Hesperophilus) bedelianus SCHÜLKE spec. nov. (Samos, Rhodos, Karpathos, Südtürkei, Libanon, Israel); Hydrosmecta cultellata Assıng spec. nov. (Samos); Atheta (Philhygra) tecta Assing spec. nov. (Samothraki, Samos, Kos, Türkei); Cretotyphlus kerkisicus Assing spec. nov. (Samos); Cephennium amplexans Assing spec. nov. (Samos); C. monstrans Assing spec. nov. (Samos); C. icariae Assing spec. nov. (Ikaría); C. rhodicum Assing spec. nov. (Rhodos); Stenichnus (Stenichnus) samius Меувонм spec. nov. (Samos); S. (S.) amphimykalicus Меувонм spec. nov. (Samos, Südwesttürkei); Scydmoraphes amphimykalicus Меувонм spec. nov. (Samos, Südwesttürkei); Euconnus (Tetramelus) rhodicus Меувонм spec. nov. (Rhodos). Drei Namen werden synonymisiert: Throbalium cycladicum (Koch, 1937) = T. biblicum (KOCH, 1937), syn. nov., = T. adanense KOCH, 1939, syn. nov., = Throbalium kosianum BORDONI, 2020, syn. nov. Zwei Arten werden in andere Gattungen transferiert: Tychus kerkisicus (BRACHAT, 2017), comb. nov. (ex Paratychus BESUCHET, 1960) und Liogluta aloconotoides (BENICK, 1940), comb. nov. (ex Aloconota THOMSON, 1858). Einige auf Fehldetermination beruhende frühere Nachweise werden korrigiert. Zahlreiche Arten werden erstmals von Kreta und den ägäischen Inseln nachgewiesen, darunter auch eine Reihe von Erstnachweisen für Griechenland und sogar für Europa. Für Samos und Rhodos werden neue Checklisten erstellt. Einschließlich der Ergänzungen und taxonomischen Änderungen sind derzeit von Kreta 403 benannte (und identifizierte) Arten (davon 114 endemisch), von Samos 199 benannte Arten (15 davon endemisch) und von Rhodos 160 benannte Arten (10 davon endemisch) bekannt.

Schlüsselwörter

Coleoptera, Staphylinidae, Paläarktis, Ostmediterraneis, Griechenland, Kreta, Ägäische Inseln, Taxonomie, Endemismus, endogäische Fauna, Beschreibungen, neue Arten, neue Synonymien, weitere Nachweise

Introduction

The Staphylinidae of the Greek islands had received relatively little attention until little more than a decade ago, although, in view of the geographic, geological, and climatic history and conditions, many of them could be expected to host endemic species. In the past years, however, numerous articles have been published addressing the faunas particularly of Crete (see Assing 2019c, Assing et al. 2019, and references cited therein), Karpathos (Assing 2016a), and the eastern Aegean islands (only the respective latest contributions cited): Samothraki (Assing 2019b), Lesbos (Assing 2016b), Chios (Assing 2015b), Samos (Assing 2017b), Ikaría (2017b), and Rhodos (Assing 2013). While the first visits to these islands focused on the epigeic fauna, a survey of the endogean species began in 2018 in Crete (AssING 2018d, ASSING 2019c, ASSING et al. 2019) and continued in Samothraki (ASSING 2019b). Since then, two field trips have been conducted to Crete and Samos, again mainly targeting the endogean fauna, the former conducted by the author in December 2019 and the latter in cooperation with Volker Brachat (Geretsried) and Thomas Forcke (Hannover) in April 2022. Moreover, Thomas Forcke conducted a field trip to Rhodos from the end of December to the beginning of January, primarily aiming at studying Carabidae, but also collecting Staphylinidae.

Additional changes to the faunas of the islands have been brought about by several generic revisions, other taxonomic studies, and faunistic contributions. In consequence, the previously published species inventories are now more or less outdated, particularly those of Samos and Rhodos, for which updated checklists are provided in this paper. Also, various previously unnamed, mostly endemic species are described, partly by Volker Brachat and Heinrich Meybohm (Großhansdorf). Descriptions of Pselaphinae are exclusively authored by Volker Brachat, those of Scydmaeninae, except *Cephennium* MüLLER & KUNZE, 1822, by Heinrich Meybohm, and one of a more widespread species of *Bledius* LEACH, 1819 by Michael Schülke (Berlin). The remaining new species are described by the author.

Material and methods

The material treated in this study is deposited in the following public and private collections:

- IRSNB Institut Royal des Sciences Naturelles de Belgique, Bruxelles (Y. Gérard)
- MHNG Muséum d'Histoire Naturelle, Genève (G. Cuccodoro)
- MNB Museum für Naturkunde Berlin (incl. coll. Schülke)
- NHMB Naturhistorisches Museum Basel (M. Borer, I. Zürcher)
- NHMW Naturhistorisches Museum Wien (H. Schillhammer)
- SDEI Senckenberg Deutsches Entomologisches Institut, Müncheberg (M. Schröter)
- cAss private collection Volker Assing, Hannover
- cBra private collection Volker Brachat, Geretsried
- cDeg private collection Gregor Degasperi, Innsbruck
- cFel private collection Benedikt Feldmann, Münster
- cKle private collection Andreas Kleeberg, Berlin
- cMey private collection Heinrich Meybohm, Großhansdorf

For the morphological studies conducted by the author, a Stemi SV 11 microscope (Zeiss), a Discovery V12 microscope (Zeiss), and a Jenalab compound microscope (Carl Zeiss Jena) were used. The images were created using digital cameras (Nikon Coolpix 995, Axiocam ERc 5s), as well as Labscope and Picolay stacking software.

Measurements: body length was measured from the anterior margin of the mandibles (in resting position) or the labrum (whichever forms the anterior apex of the head) to the posterior apex of the elytra (Scydmaeninae) or to the abdominal apex (other subfamilies), the length of the forebody from the apex of the head to the posterior margin of the elytra (all subfamilies except Scydmaeninae), head length along the middle from the anterior margin of the clypeus (without ante-clypeus) (Aleocharinae) or the anterior margin of the frons (other subfamilies) to the posterior constriction of the head, head width across (and including) eyes, elytral length at the suture from the apex of the scutellum to the posterior margin of the elytra (except in Bledius), and the length of (the median lobe of) the aedeagus from the apex of the ventral process to the base of the aedeagal capsule unless stated otherwise. The "parameral" side (i.e., the side where the sperm duct enters) is referred to as the ventral, the opposite side as the dorsal aspect.

The Staphylinidae fauna of Crete

The fauna of Crete is remarkable not only regarding its diversity (397 named species prior to this study), but also with respect to the number of endemic species (111 named), many of them locally endemic. A comprehensive overview of the Staphylinidae of Crete was provided in a recent two-part monograph (Assing 2019, Assing et al. 2019). In the meantime an additional field trip was conducted to this island by the author in December 2019, with a special focus on the endogean fauna. Compared to previous field trips to Crete, the 2019 results were relatively poor, yielding a total of only 294 specimens, with merely two new species of Pselaphinae, a first record of a species of Tachyporinae, and few additional records of other species (Tab. 1). Moreover, two additional species, Carpelimus gusarovi GILDENKOV, 1992 and C. exiguus (ERICHSON, 1839), were reported from Crete for the first time by GILDENKOV & MANTIČ (2020). According to SCHÜLKE (2019), there is evidence that Planeustomus elegantulus (KRAATZ, 1857) is endemic to Crete. Andreas Kleeberg (pers. comm.) communicated the first record of Ochthephilus and alusiacus (FAGEL, 1957) from the island (see details below). Including the new species and the additional records, 403 named species are currently known from Crete, with 114 (28.2 %) of them endemic.

Localities/samples: December 2019, leg. Assing: 1: SW Chania, WSW Lakki, 35°23'53"N, 23°54'38"E, 390 m, stream valley, very old chestnut on slope near road, stony soil with little humus, soil washing, 21.XII.2019; **1a**: SW Chania, WSW Lakki, 35°23'53"N, 23°54'38"E, 390 m, stream valley, beneath *Platanus orientalis* near stream, fine soil with humus,

Tab. 1: Staphylinidae recorded from Crete in December 2019. Details on the localities/samples are provided at the end of the list. Specimen numbers are given in parentheses behind the sample number. First records from Crete are marked with an asterisk, endemics are preceded by a "+".

Species	Localities/samples
O m a l i i n a e	
+Boreaphilus fuelscheri ZERCHE, 1990	24(1), 29(1)
Proteininae	
+Proteinus creticus Assıng, 2004	19a(1)
P s e l a p h i n a e	
+*Afropselaphus chanianus BRACHAT, spec. nov.	1(1)
+Bryaxis meybohmianus BRACHAT, 2019	1(5), 4(1), 5(2), 5(2), 8(1), 9(3), 18(2)
+Euplectus assingi BRACHAT, 2019	1(1), 18(1)
+*Paramaurops creticus BRACHAT, spec. nov.	19a(1)
+ <i>Tychus reitterianus</i> Löbl, 1998	7(2), 11(2), 12(1)
Tachyporinae	
*Lordithon trinotatus (ERICHSON, 1839)	41a(1), 96a(1), 159(1), 172(1), 193(1), 212(1)
<i>Mycetoporus ignidorsum</i> Еррегsнеім, 1880	21a(1)
Tachyporus caucasicus (Kolenati, 1846)	17(1), 19a(1)
Habrocerinae	
Habrocerus pisidicus Korge, 1971	7(11), 15(1), 19a(26)
Aleocharinae	
+Alevonota cretica Assing & Wunderle, 2008	29(1)
Aloconota subgrandis (BRUNDIN, 1954)	7(1), 27(3)
Atheta aeneicollis (SHARP, 1869)	1(2), 1a(1), 6(1), 12(1), 13(1), 15(1), 20(1)
+Atheta cretica Brundin, 1944	7(1), 13(2)
Atheta (Mocyta) spp.	1(1), 15(1), 19a(2)
+Bellatheta albimontis Assıng, 2015	2(1)
Bolitochara obliqua Erichson, 1837	14a(4)
Cousya defecta (MULSANT & REY, 1875)	14a(1)
+Geostiba meybohmi Assıng, 2000	21a(3), 22(24), 24(1), 29(5), 30(2)
Geostiba oertzeni (Еррегsнеім, 1888)	2(14), 3(4), 5(3), 6(2), 15(3), 16(4), 17(1), 27(6), 28(2), 29(4)
Myllaena intermedia Erichson, 1837	7(36)
Oxypoda flavicornis KRAATZ, 1856	29(4)
Oxypoda rectacia Assing, 2019	7(1), 10(1), 19a(3)
O x y t e l i n a e	
Anotylus tetracarinatus (ВLOCK, 1799)	26(1)
Carpelimus corticinus (GRAVENHORST, 1806)	7(10)
Platystethus nitens (C.R. SAHLBERG, 1832)	19a(1)
Steninae	
Stenus ochropus Kiesenwetter, 1858	21a(2)
S c y d m a e n i n a e	
+Cephennium idanum Assıng, 2019	22(2), 25(2)
<i>Cephennium</i> spp. (♀)	28(1), 29(1)
+Scydmoraphes ziegleri Меувонм, 2008	26(1), 28(2), 29(3)
+ <i>Stenichnus brachati</i> Меувонм, 2019	21a(2)

Species	Localities/samples
L e p t o t y p h l i n a e	
+Allotyphlus foedatus Assing, 2019	1(1), 1a(2), 3(2), 5(1), 12(1), 16(2), 19(1)
+ <i>Cretotyphlus idanus</i> Assıng, 2019	19(1)
+Kenotyphlus virgatus Assıng, 2019	6(2)
Paederinae	
+ <i>Leptobium creticum</i> COIFFAIT, 1973	5(1), 15(1), 23(1)
+Medon cerrutii Coiffait, 1976	6(2), 7(1), 15(1)
Medon dilutus pythonissa (SAULCY, 1865)	7(1), 12(1), 13(1), 15(2), 18(1)
Sunius fallax (Lokay, 1919)	7(3)
Staphylininae	
<i>Gyrohypnus angustatus</i> STEPHENS, 1833	7(1)
Ocypus mus (Brullé, 1832)	20(1)
Othius lapidicola Märkel & Kiesenwetter, 1848	18(1), 23(2), 24(1), 25(2), 30(1)
Quedius nemoralis BAUDI DI SELVE, 1848	22(1)
+Quedius sigwalti COIFFAIT, 1972	7(2), 19a(1)
+ <i>Xantholinus creticus</i> Assıng, 2006	24(2)

washing, 21.XII.2019; 2: SW Chania, WSW soil Lakki, 35°23'53"N, 23°54'38"E, 390 m, stream valley, beneath Platanus orientalis near stream, soil washing, 22.XII.2019; **3**: SW Chania, W Omalos Plateau, 35°21'11"N, 23°48'42"E, 790 m, dry stream valley with old Platanus orientalis on slope with Juniperus and phrygana, soil washing beneath Platanus and between rocks, 21.XII.2019; 4: SW Chania, W Omalos Plateau, 35°20'50"N, 23°49'20"E, 710 m, moist slope with chestnut, soil washing, 21.XII.2019; 5: SW Chania, W Omalos Plateau, 35°20'21"N, 23°49'50"E, 640 m, ruderal slope with Quercus ilex and bushes, soil washing near dry stream bed, 21.XII.2019; 6: S Kissamos, Mili, 35°23'18"N, 23°39'49"E, 350 m, stream valley with Platanus orientalis, soil washing near stream, 22.XII.2019; 7: S Kissamos, Limni, 35°22'15"N, 23°37'55"E, 570 m, margin of artificial pond with old Platanus orientalis, litter sifted, 22.XII.2019; 8: NE Kandanos, Spina, 35°21'43"N, 23°46'51"E, 710 m, stream valley with oak, Platanus orientalis, other trees, and fern undergrowth, soil washing, 22.XII.2019; 9: NE Kandanos, Spina, 35°21'46"N, 23°46'38"E, 700 m, moist slope with very old chestnut, soil washing, 22.XII.2019; 10: SE Chania, SW Melidoni, 35°20'44"N, 24°05'49"E, 1030 m, stony slope with Cedrus and Quercus ilex, soil washing, 23.XII.2019; 11: SE Chania, W Melidoni, 35°23'35"N, 24°05'45"E, 400 m, small dry stream valley with Platanus orientalis, soil washing, 23.XII.2019; 12: SE Chania, Ramni, 35°23'51"N, 24°05'56"E, 400 m, grassy slope with old oak and Rubus undergrowth, sandy soil washing, 23.XII.2019; 13: SE Chania, Kambi, 35°24'57"N, 24°04'09"E, 550 m, ruderal slope with old oak, soil washing, 23.XII.2019; 14: SW Chania, S Nteres, 35°24'48"N, 23°50'46"E, 290 m, stream valley with very old chestnut trees, soil washing (stony soil), 24.XII.2019; 14a: same data, but litter sifted; 15: SW Chania, S Nteres, 35°24'15"N, 23°50'58"E, 320 m, stream valley with chestnut trees, soil washing (stony soil), 24.XII.2019; 16: SW Chania, N Nteres, 35°27'40"N, 23°50'18"E, 170 m, dry stream valley with various trees and fern, soil washing, 24.XII.2019; 17: SE Vrises, E Askifou, 35°18'32"N, 24°13'18"E, 880 m, scree slope with Quercus ilex and Arum undergrowth, soil washing, 25.XII.2019; 18: SE Vrises, E Askifou, 35°18'22"N, 24°14'06"E, 990 m, scree slope with Quercus ilex and Acer sp., soil washing, 25.XII.2019; 19: Psiloritis, NE Livadia, 35°18'56"N, 24°49'07"E, 440 m, stream valley with *Platanus orientalis*, soil washing near large rocks, 26.XII.2019; 19a: same data, but litter sifted; 20: E Malia, N Latsida, 35°16'08"N, 25°34'38"E, 260 m, dry stream valley between olive groves with oak and other trees, soil washing, 27.XII.2019; 21: SE Malia, Selena Oros, 35°14'50"N, 25°30'56"E, 840 m, rocky calcareous slope with Quercus ilex and phrygana, soil washing, 27.XII.2019; 21a: same data, but litter and soil sifted; 22: SE Malia, Selena Oros, 35°14'41"N, 25°31'03"E, 860 m, rocky calcareous slope with Quercus ilex and phrygana, litter and soil sifted, 27.XII.2019; 23: NW Ierapetra, NW Males, 35°05'03"N, 25°34'36"E, 510 m, steep stream valley, soil and gravel washing beneath old Platanus orientalis, 28.XII.2019; 24: NW Ierapetra, Selakano, 35°05'14"N, 25°32'48"E, 840 m, rocky stream valley with Quercus ilex, soil washing, 28.XII.2019; 25: NW Ierapetra, Selakano, 35°05'20"N, 25°32'51"E, 840 m, rocky slope near stream with Quercus ilex, soil washing, 28.XII.2019; 26: NW Ierapetra, NE Anatoli, 35°03'02"N, 25°38'59"E, 590 m, steep rocky slope with Quercus sp., Quercus ilex, and other bushes, stony soil washing near road margin, 28.XII.2019; **27**: E Agios Nikolaos, S Sfaka, 35°09'00"N, 25°55'26"E, 320 m, narrow dry stream valley with *Quercus ilex*, soil washing, 29.XII.2019; **28**: E Agios Nikolaos, E Mirsini, 35°10'06"N, 25°56'59"E, 320 m, narrow dry stream valley with *Platanus orientalis*, *Quercus ilex*, and bushes, soil and gravel washing, 29.XII.2019; **29**: SW Agios Nikolaos, SW Kroustas, 35°07'02"N, 25°37'56"E, 810 m, rocky slope with *Quercus ilex*, soil washing, 30.XII.2019; **30**: SW Agios Nikolaos, SW Kroustas, 35°06'42"N, 25°37'34"E, 930 m, rocky slope with *Quercus ilex* and *Pinus* sp., soil washing, 30.XII.2019.

Descriptions of new species

Paramaurops creticus BRACHAT spec. nov. urn:lsid:zoobank.org:act:0ADD6720-1EEC-4E67-90EE-F1DC5B1C9ACE (Figs 1, 21)

Type material: Holotype ♂: "GR – Crete [19a], Psiloritis, NE Livadia, 35°18'56"N, 24°49'07"E, 440 m, litter sifted, 26.XII.2019, V. Assing / *Paramaurops creticus* spec. nov. ♂, det. Brachat 2.2020 / Holotypus" (cBra).

Etymology: The specific epithet (Latin, adjective) alludes to the restricted distribution of this species.

Description: Body 2.40 mm long, pale reddish-brown, smooth, and with scattered yellowish pubescence. Habitus as in Fig. 1.

Head 0.52 mm long and 0.48 mm broad, postero-laterally with long erect pubescence. Eyes absent, replaced with a stout lateral spine-shaped extension; carinae extending from antennal protuberances very short; vertex posteriorly with a short median carina. Antennae slender, 1.27 mm long; all antennomeres distinctly oblong; antennomeres III–VII twice as long as broad. Maxillary palpomere IV elongate, 0.24 mm long.

Pronotum 0.48 mm long and 0.42 mm broad, without lateral carinae and without protuberances on disc; disc anteriorly with weakly pronounced median carina, basally with a small lateral fovea on either side and with a short median carina, ante-basally with a small median and two small lateral foveae.

Elytra 0.60 mm long and 0.72 mm broad, with three indistinct basal foveae, without impression on disc. Hind wings completely reduced. Metaventrite with a median fovea near posterior margin. Legs long and slender.

Abdomen: tergite IV 0.54 mm long and 0.78 mm broad, basally with a large median and two small and shallow lateral foveae, median fovea densely pubescent, sublateral carinae situated at some distance from lateral margins, oblique, and approximately half as long as tergite.

 σ : trochanters and femora unmodified; metatibia with short apical spine; aedeagus 0.30 mm long and shaped as in Fig. 21.

Comparative notes: The new species is distinguished from the similar *Paramaurops kaufmanni* (GANGLBAUER, 1895) and *P. commixtus* (J. MÜLLER, 1928) from the Balkans by distinctly larger body size (*P. kaufmanni* and *P. commixtus*: body length approximately 2.0 mm) and by the shape of the aedeagus.

Distribution: The type locality is situated near Livadia in the north slope of the Psiloritis range. The holotype was sifted from deep litter of *Platanus orientalis* in a stream valley with large rocks at an altitude of 440 m. For a photograph of this locality, which is also the type locality of *Cretotyphlus idanus* ASSING, 2019, *Cephennium idanum* ASSING, 2019, and *C. latius* ASSING, 2019 see figure 175 in ASSING et al. (2019).

Afropselaphus chanianus BRACHAT spec. nov. urn:lsid:zoobank.org:act:7648FB38-4026-414C-8609-32447D514299 (Figs 2, 22)

Type material: Holotypus ♂: "GR – Crete [1], SW Chania, WSW Lakki, 35°23'53"N, 23°54'38"E, 390 m, stream valley, 21.XII.2019, V. Assing / *Afropselaphus chanianus* spec. nov. ♂ det. Brachat 2.2020 / Holotypus" (cBra).

Etymology: The specific epithet is an adjective derived from Chania, the district where the type locality is situated.

Description: Body 1.60 mm long, reddish-brown, smooth, with scattered long yellowish setae. Habitus as in Fig. 2.

Head 0.37 mm long and 0.26 mm broad; frons with two carinae extending from antennal insertions to frontal foveae; posterior portion between frontal foveae and posterior margin of head shallowly impressed. Eyes small, composed of 7–8 ommatidia, less than half as long as postocular portion. Antenna 0.9 mm long; all antennomeres oblong, antennomeres IX–XI forming a club; antennomere XI as long as the combined length of antennomeres VIII–X. Maxillary palpomere IV club-shaped in apical half.

Pronotum 0.29 mm long and 0.25 mm broad, with three distinct ante-basal foveae. Elytra short, 0.34 mm long and 0.56 mm broad; posterior margin with dense sub-erect pubescence. Metaventrite distinctly domed.

Abdomen: tergite IV 0.34 mm long and 0.56 mm broad, anteriorly with dense scaly pubescence and a deep impression; sternite IV anteriorly without depression.

 σ : aedeagus 0.29 mm long and shaped as in Fig. 2; parameres each with three apical setae.

Comparative notes: The new species is highly similar to the geographically close *A*. *doernfeldi* BRACHAT, 2019 and *A*. *assingi* BRACHAT, 2019. It is reliably distinguished from them only based on the shape of the aedeagus. For illus-

trations of the aedeagi of *A. doernfeldi* and *A. assingi* see Assing et al. (2019).

In December 2018, an *Afropselaphus* male was collected practically in the same locality (see Assing 2019c). This male most likely represents an undescribed species distinguished from *A. chanianus* by a larger (0.33 mm), apically tapering aedeagus with a long internal structure. Since this structure is extruded in the only specimen available, a formal description is refrained from.

Distribution and natural history: The type locality is situated to the southwest of Chania, West Crete, at an altitude of 390 m. The holotype was collected by washing stony soil with little humus from a stream valley with a very old chestnut tree.

Notes on some species

Lordithon trinotatus (ERICHSON, 1839)

The specimens listed in Tab. 1 represent the first record from Crete.

Cephennium idanum Assıng, 2019 (Fig. 23)

The previously known distribution was confined to the type locality near Livadia in the north slope of the Psiloritis range, Central Crete. In December 2019, the species was recorded also from Selena Oros to the southeast of Malia and another locality to the Northwest of Ierapetra, records that significantly expand the known range towards the east of Crete. The aedeagus of a male from Selena Oros is illustrated in Fig. 23.

Allotyphlus foedatus Assing, 2019

This species was previously known only from West Crete eastwards to the Askifou plateau. The male from the environs of Livadia in the Psiloritis range considerably expands the known distribution eastwards to Central Crete.

Throbalium creticum COIFFAIT, 1980 and *T. dividuum dividuum* (ERICHSON, 1840)

The original description of *T. creticum* is based on a male holotype and four female paratypes from "Baie de Colpos-Soudas, Chypre" deposited in the Coiffait collection (COIFFAIT 1980). The type locality ("Chypre") is evidently a lapsus and was rectified by COIFFAIT (1982), who stated that the species was in fact distributed in "Crète". Nevertheless, it is still indicated for Cyprus in the Palaearctic Catalogue (SCHÜLKE & SMETANA 2015),

which partly explains why *T. creticum* was omitted from the checklist of the Staphylinidae of Crete in ASSING (2019c). Owing to the currently restricted loan policy of the Muséum National d'Histoire Naturelle, Paris, it was not possible to study the type material in order to clarify the status of the species and the question whether or not it is endemic to the island.

Based on old literature records by KRAATZ (1858) and OERTZEN (1887), *Throbalium dividuum dividuum* is the only *Throbalium* species listed for Crete in Assing (2019c). In view of the otherwise West Mediterranean distribution of *T. dividuum* there is little doubt that these records in fact refer to *T. creticum*.

Additional records

Andreas Kleeberg (Berlin) communicated additional records from the following localities:

1. S-Crete, Agios Galini, 3.IV.2002, leg. Kleeberg: Ochthephilus andalusiacus (FAGEL, 1957): 8 exs. [det. Makranczy] (cKle); Ochthephilus venustulus RosenнAUER, 1856: 4 exs. [det. Makranczy] (cKle); Gnypeta rubrior Тоттемнам, 1939: 1 ex. (cKle).

2. Central Crete, S Kaneros, Kotsifos Gorge, 5.IV.2002, leg. Kleeberg: *Lesteva nitidicollis* LOHSE & STEEL, 1961: 6 exs. [det. Schülke] (cKle); *Platystethus spinosus* ERICHSON, 1840: 2 exs. [det. Schülke] (cKle); *Aloconota subgrandis* (BRUNDIN, 1954): 1 ex. [det. Assing] (cKle).

3. West Crete, Lake Kournas, 4.IV.2002, leg. Kleeberg: *Carpelimus obesus* (KIESENWETTER, 1844): 3 exs. [det. Kleeberg] (cKle).

4. West Crete, Georgiopolis env., Aspropoliani, 30.IX.–6.X.2002, leg. Kleeberg: *Carpelimus bilineatus* STEPHENS, 1834: 2 exs. [det. Kleeberg] (cKle); *Stenus hospes* ERICHSON, 1840: 18 exs. [det. Puthz] (cKle, cPut); *Stenus brunnipes lepidus* WEISE, 1875: 1 ex. [det. Puthz] (cKle); *Stenus picipes* STEPHENS, 1833: 12 exs. [det. Puthz] (cKle, cPut); *Rugilus orbiculatus* (PAYKULL, 1789): 8 exs. [det. Kleeberg] (cKle); *Aloconota subgrandis* (BRUNDIN, 1954): 1 ex. [det. Assing] (cKle); *Gabrius nigritulus* (GRAVENHORST, 1802): 8 exs. [det. Schülke] (cKle); *Gabronthus maritimus* (MOTSCHULSKY, 1858): 5 exs. [det. Schülke] (cKle).

5. West Crete, Georgiopolis, beach, 25.IV.1992, leg. Rößner: *Stenistoderus cephalotes* (KRAATZ, 1858): 1 ex. [det. Janák] (cKle).

Ochthephilus andalusiacus is reported from Crete for the first time.

Tab. 2: Updated checklist of the Staphylinidae recorded from Samos and results of the field trip to Samos conducted by Volker Brachat, Thomas Forcke, and the author in April 2022. Details on the localities/samples are provided at the end of the list. Specimen numbers are given in parentheses behind the sample number. First records of non-endemic species from Samos are marked with an asterisk (*), first records from Greece with two, and first records from Europe with three asterisks. Endemic species are preceded by a "+". For additional footnotes and references for species recorded from Samos before 2017 see the footnotes and references in Assing (2017b).

The references are abbreviated as follows: A17b = Assing (2017b); A18c = Assing (2018c); A19a = Assing (2019a); Aip = Assing (in press); App = Assing (present paper); G19 = GILDENKOV (2019), M14 = MAKRANCZY (2014).

Species	Localities/samples	References
O m a l i i n a e		
Boreaphilus velox (HEER, 1839)		A17b
*Dropephylla perforata (ZANETTI, 1987)	4(1)	Арр
Omalium cinnamomeum KRAATZ, 1857	10(1), 10a(2), 11(1), 11a(17), 12a(1), 24(1), 29(3), 29a(1), 30(3), 37(1), 47a(3)	A17b, App
Omalium rhodicum ZANETTI & Assing, 2013		A17b
Omalium rugatum Mulsant & Rey, 1880	9a(1), 31(1), 60(17)	A17b, App
Philorinum hopffgarteni Eppelsheim, 1892		A17b
Proteininae		
Metopsia assingi ZERCHE, 1998		A17b
Proteinus utrarius Assıng, 2004	3(1), 4(1), 5(75), 6(2), 9(1), 11a(1), 32a(1)	A17b, App
Micropeplinae		
Micropeplus fulvus Erichson, 1840	3a(1), 4(3), 5(1), 8(2), 16a(3), 36(1)	A17b, App
Micropeplus staphylinoides (Макsнам, 1802)	27(2)	A17b, App
P s e l a p h i n a e		
*Batrisodes buqueti (AUBÉ, 1833)	36b(2)	Арр
*Batrisodes delaporti (Аив́е, 1833)	1d(1)	Арр
Bibloplectus parvulus BESUCHET, 1975		A17b
*Brachygluta foveola foveola (Motschulsky, 1840)	41(1)	Арр
*Brachygluta helferi longispina (REITTER, 1884)	50a(5)	Арр
Brachygluta spinicoxis fuchsii (PAGANETTI-HUMM- LER, 1899)	49(1)	A17b, App
*Brachygluta xanthoptera (REICHENBACH, 1816)	23b(1)	Арр
Bryaxis anatolicus (SAULCY, 1878)	3(1), 3a(4), 5(1), 8(2), 9a(3), 10a(6), 13a(4), A17b, 16a(7), 18a(2), 19(1), 21(1), 24a(4), 29a(2), 30a(1), 34(3), 34a(2), 37(5), 37b(3), 39(1), 40(3), 40a(5), 43(1), 51(2) A17b,	
Bryaxis pumilus Bekchiev & Brachat, 2015	8(7), 10(2), 10a(4), 11(2), 11a(32), 12(2), 12a(11), A17b, A 14a(5), 16a(2), 24a(8), 29a(2), 30(6), 30a(2), 34(1), 36(3), 47(4), 47a(13), 48(9), 51a(1), 54(2)	
Bythinus simplicipalpis BRACHAT, 2016		A17b
Claviger spp.	36a(3), 36b(1), 40b(2)	A17b, App
*Euplectus kirbii kirbii Denny, 1825	36a(3), 36b(2)	Арр
+Euplectus meybohmi BRACHAT, 2017	3(1)	A17b, App
Faronus distinctus Besuchet, 1999		A17b
+Namunia cavernicola Besuchet, 1978		A17b
+Paramaurops spp. (undescribed) (9)	10(1), 13(1)	A17b, App
Paratychus mendax (KIESENWETTER, 1858)		A17b

Species	Localities/samples	References
Tribatus creticus REITTER, 1884		A17b
Trimium caucasicum Kolenati, 1846	2(1), 3(3), 3a(1), 4(2), 7(1), 8(1), 11a(9), 12(4), 12a(10), 13a(2), 21a(1), 24a(9), 29a(4), 30(2), 30a(3), 33(1), 34(1), 36a(2), 51a(3), 52b(1), 53a(1), 54(1)	A17b, App
+ <i>Trimium</i> sp. (undescribed) (♀)		A17b
+ <i>Tychobythinus brachati</i> Besucнет, 2008		A17b
<i>Tychus anatolicus</i> Besucher, 1964		A17b
+ <i>Tychus kerkisicus</i> (BRACHAT, 2017), comb. nov.	37(1)	A17b, App
<i>Tychus</i> sp. (♀)	12a(1), 37(1)	Арр
Phloeocharinae		
Phloeocharis longipennis FAUVEL, 1875		A17b
T a c h y p o r i n a e		
Bryoporus multipunctus HAMPE, 1867		A17b
Lordithon bimaculatus (SCHRANK, 1798)		A17b
*Lordithon trinotatus (ERICHSON, 1839)	1d(1)	Арр
Mycetoporus confinis Rey, 1883		A17b
Mycetoporus ignidorsum Eppelsheim, 1880	12a(1)	A17b, App
Mycetoporus imperialis Bernhauer, 1902	11a(1), 12a(2), 31(1), 38(1), 46a(2), 47a(3), 51(1)	A17b, App
Mycetoporus macrocephalus Bernhauer, 1917		A17b
Mycetoporus monticola Fowler, 1888		A17b
Mycetoporus reichei (PANDELLÉ, 1869)	3a(1), 8(3), 12a(2), 53(1)	A17b, App
Mycetoporus simillimus FAGEL, 1965	1b(12), 19(1), 33(1), 40(1), 46(1), 46a(11), 47a(19), 48(1), 55(1), 57(7)	A17b, App
<i>Mycetoporus</i> (<i>nigricollis</i> group) sp. (♀)	30(1)	Арр
Sepedophilus immaculatus (STEPHENS, 1832)		A17b
Sepedophilus obtusus (Luze, 1902)		A17b
Sepedophilus testaceus (FABRICIUS, 1792)	1a(1), 1d(2), 52a(1)	A17b, App
Tachyporus abner SAULCY, 1865		A17b
Tachyporus caucasicus Kolenati, 1846		A17b
<i>*Tachyporus hypnorum</i> (FABRICIUS, 1775)	1c(1)	Арр
Tachyporus nitidulus (FABRICIUS, 1781)		A17b
Habrocerinae		
Habrocerus pisidicus Korge, 1971	9a(1), 12a(1), 52a(1)	A17b, App
Aleocharinae		
Acrotona muscorum (BRISOUT, 1860)		A17b
Aleochara cf. conviva Eppelsheim, 1890		A17b
*Aleochara bipustulata (LINNAEUS, 1760)	22(1)	Арр
Aleochara haematoptera KRAATZ, 1858	23a(3), 49(1)	A17b, App
Aleochara rambouseki Likovský, 1981	1b(1), 13a(1), 14a(1)	A17b, App
Alevonota rufotestacea (KRAATZ, 1856)		A17b
Aloconota aegea Assing, 2016	23a(1), 23b(3)	A17b, App
*Aloconota cambrica (WOLLASTON, 1855)	23b(3)	App
Aloconota samia Assing, 2016	60(2)	A17b, App

Species	Localities/samples	References
Atheta aeneicollis (SHARP, 1869)	$\begin{array}{l} 3(1), \ 8(1), \ 9(6), \ 10(3), \ 10a(2), \ 11(1), \ 12(1), \\ 12a(7), \ 13(4), \ 13a(1), \ 14a(3), \ 23a(1), \ 24a(3), \\ 27(1), \ 28(2), \ 29(17), \ 30(3), \ 31(2), \ 32a(3), \ 33(2), \\ 36(1), \ 36b(1), \ 37(1), \ 40a(1), \ 43(4), \ 44(1), \ 46a(2), \\ 47(4), \ 47a(2), \ 48(4), \ 51(1), \ 55(1), \ 60(2) \end{array}$	A17b, App
*Atheta amicula (STEPHENS, 1832)	47(1)	App
Atheta speculum (KRAATZ, 1856)		A17b
*Atheta tecta Assing, spec. nov.	41(2), 41a(3), 50a(1), 56(1)	Арр
Atheta trinotata (KRAATZ, 1856)		A17b
Atheta pulchra (KRAATZ, 1856)		A17b
Atheta (Microdota) sp. (?)		A17b
Atheta (Mocyta) spp.	8(1), 11a(2), 12a(1), 24a(2), 36a(1), 47a(29), 50a(1), 56(11), 60(5)	A17b, App
Brundinia meridionalis (Mulsant & Rey, 1853)	35(1), 35a(2), 45(3), 50(2), 50a(4)	A17b, App
Cousya defecta (MULSANT & REY, 1875)	1c(1)	A17b, App
Cypha spathulata Assing, 2007		A17b
Cypha cf. tarsalis (Luze, 1902) (♀)		A17b
Dinusa cretica Assıng, 2013		A17b
Geostiba maxiana (Тікномігоvа, 1973)	1b(4), 42(1), 46(2), 46a(12)	A17b, App
Geostiba oertzeni (ЕрреLsнеім, 1888)	(4), 9(1), 9a(1), 10(1), 10a(5), 11(1), 24a(1), 28(3), 29(2), 30(3), 30a(1), 33(1), 34(7), 36(1), 37(7), 37a(1), 37b(4), 39(1), 40a(5), 51(9), 51a(1), 52a(1), 52b(1)	A17b, App
+Geostiba plicipennis Assıng, 2015	1a(1), 3(1)	A17b, App
*Hydrosmecta cultellata Assing, spec. nov.	60(4)	Арр
*Hydrosmecta insularum Assıng, 2019	23b(4)	Арр
* <i>Hydrosmecta</i> sp. [♀]	23b(6)	Арр
*Ischnoglossa minor (AuBé, 1863)		Aip, App
Ischnoglossa turcica Wunderle, 1992		A17b
Leptusa samia Assing, 2004	1d(1), 11a(1), 36(1)	A17b, App
Liogluta longiuscula (GRAVENHORST, 1802)	12a(3), 47a(1)	A17b, App
<i>Liogluta</i> sp.	60(1)	Арр
Maurachelia roubali (Lонse, 1970)		A17b
*** <i>Meotica truncata</i> Assing, 2004	26(2)	Арр
*Myllaena graeca KRAATZ, 1858	31(2), 31a(3)	Арр
*Myllaena infuscata KRAATZ, 1858	50(1)	Арр
Myllaena intermedia Erichson, 1837	41a(1)	A17b, App
Myrmecopora convexula Assıng, 1997	17(1)	A17b, App
Ocalea brachyptera FAGEL, 1971		A17b
Oligota pumilio Kiesenwetter, 1858	24a(1)	A17b, App
Oxypoda acutissima Assıng, 2006		A17b
Oxypoda flavicornis KRAATZ, 1856		A17b
***Oxypoda flavocaudata Assing, 2006	24a(1), 47a(1), 48(1)	Арр
***Oxypoda gladiatoria Assıng, 2006	12a(1), 46a(1), 57(1)	Арр
*Oxypoda cf. haemorrhoa (Маппегнеім, 1830)	47(1)	Арр

Species	Localities/samples	References
+Oxypoda kerkisica Assıng, 2015	57(3)	A17b, App
Oxypoda libanotica FAGEL, 1965		A17b
Oxypoda lurida WOLLASTON, 1857	47(2), 53a(1), 57(1), 60(2)	A17b, App
Oxypoda rectacia Assıng, 2019	$ \begin{array}{c} 1a(1), \ 1b(12), \ 3a(1), \ 12a(1), \ 14a(1), \ 32a(1), \\ 40a(2), 42(1), 46a(7), 47a(1), 47a(1), 60(1) \end{array} $	A17b, A19a, App
Oxypoda scheerpeltziana (FAGEL, 1968)		A17b
Oxypoda turcica Zercне, 1999		A17b
Oxypoda vicina KRAATZ, 1858		A17b
Parocyusa longitarsis (Erichson, 1839)	23(1), 23a(2), 23b(1), 49(3), 60(4)	A17b, App
Piochardia reitteri (WASMANN, 1894)	58(1)	A17b, App
Pseudosemiris kaufmanni (Eppelsheim, 1887)		A17b
Taxicera moczarskii (Bernhauer, 1914)		A17b
O x y t e l i n a e		
*Anotylus clypeonitens (PANDELLÉ, 1867)	23b(1), 49(1)	Арр
*Anotylus complanatus (ERICHSON, 1839)	8(1)	Арр
Anotylus inustus (Gravenhorst, 1806)	1b(1), 8(5), 10a(3), 11a(1), 11b(1), 12as(4), 13(1), 14a(1), 22(6), 23(2), 23a(11), 23b(3), 24a(2), 29(1), 33(1), 34(1), 36(1), 45(1), 49(1), 60(12)	A17b, App
*Bledius bedelianus SCHÜLKE, spec. nov.	23a(3), 23b(2), 49(9), 60(1)	Арр
Bledius frisius Lonse, 1978	35(14), 35a(2)	A17b, App
Bledius furcatus (OLIVIER, 1811)	35(23), 35a(12)	A17b, App
Bledius unicornis (GERMAR, 1825)	35(27), 35a(12), 50(4), 50a(3)	A17b, App
*Carpelimus bilineatus (STEPHENS, 1834)	60(2)	Арр
*Carpelimus corticinus (GRAVENHORST, 1806)	41a(1), 60(6)	Арр
*Carpelimus fuliginosus (GRAVENHORST, 1802)	60(1)	Арр
*Carpelimus gracilis (MANNERHEIM, 1830)	60(1)	Арр
*Carpelimus similis (SMETANA, 1967)	60(1)	Арр
* <i>Carpelimus simplicicollis</i> (WOLLASTON, 1857) ¹⁾		A17b, App, G19
*Euphanias ambulans Assing, spec. nov.	35a(2)	Арр
Ochthephilus and alusiacus (FAGEL, 1957)	49(1), 60(3)	App, M14
Planeustomus cephalotes (ERICHSON, 1840)		A17b
Platystethus spinosus Erichson, 1840		A17b
*Thinodromus bodemeyeri (Bernhauer, 1902)	60(1)	Арр
S t e n i n a e		
Stenus aceris Stephens, 1833	8(1), 9a(1), 24a(1), 30(1), 31(8), 31a(7), 37a(1), 41(1), 41a(1), 50a(1), 56(2)	A17b, App
Stenus brunnipes lepidus WEISE, 1875		A17b
*Stenus callidus BAUDI DI SELVE, 1848	41(1), 41a(2)	Арр
Stenus glacialis HEER, 1839	1c(1)	A17b, App
Stenus hospes Erichson, 1840		A17b
Stenus maculiger WEISE, 1875	9a(1), 23a(1), 31(5), 31a(2), 49(2), 59(1), 60(14)	A17b, App
Stenus parcior Bernhauer, 1929		A17b

Species	Localities/samples	References
Stenus turbulentus Bondroit, 1912	8(2), 9a(6), 9b(1), 12a(1), 36b(1), 48(1), 37a(1), 39(1), 41(1), 46a(8), 55(1), 57(1)	A17b, App
Stenus turcicus Bernhauer, 1912	23(1)	A17b, App
Scydmaeninae		
+ <i>Cephennium ambulans</i> Assing, spec. nov. ¹⁾		A17b, App
+ <i>Cephennium monstrans</i> AssING, spec. nov. ¹⁾		A17b, App
<i>Chevrolatia</i> sp. (♀)		A17b
+ <i>Euconnus ambelosicus</i> Меувонм, 2017	1(1), 8(9), 10a(13), 11a(4), 30(7), 30a(1), 36(1), 40b(1), 51(1), 52(3)	A17b, App
Euconnus intrusus (SCHAUM, 1844)	50a(1)	A17b, App
+ <i>Euconnus kerkisicus</i> Меувонм, 2017	13a(2), 24a(1)	A17b, App
<i>Euconnus samius</i> Меувонм, 2017	10a(1), 11(3), 12(2), 14a(1), 16a(1), 24a(1), $34a(1), 54(1)$	A17b, App
<i>Euconnus</i> sp. (♀; undescribed)		A17b
Eutheia sp.		A17b
Leptomastax simonis Stussiner, 1881	12a(1), 13(1), 24(2), 28(1), 37(1), 37a(1), 53(1)	A17b, App
*Scydmaenus cornutus Motschulsky, 1844	36a(3), 36b(3)	Арр
<i>Scydmoraphes amphimykalicus</i> Меувонм, spec. nov.	14a(2)	A17b ¹⁾ , App
Stenichnus amphimykalicus Меувонм, spec. nov.		A17b ¹⁾ , App
**Stenichnus cribrum (SAULCY, 1880)	36a(10), 36b(3), 40b(1)	A17b ¹⁾ , App
+ <i>Stenichnus samius</i> Меувонм, spec. nov.	13a(1), 24a(1)	A17b ¹⁾ , App
P s e u d o p s i n a e		
Pseudopsis sulcata NEWMAN, 1834		A17b
Leptotyphlinae		
+Cretotyphlus kerkisicus Assıng, spec. nov.	18(1)	Арр
+ <i>Cretotyphlus</i> sp.	5(1), 12(1), 38(1)	Арр
Paederinae		
Astenus bimaculatus (ERICHSON, 1840)		A17b
Astenus procerus (GRAVENHORST, 1806)		A17b
Astenus thoracicus (BAUDI DI SELVE, 1857)		A17b
Domene stilicina (ERICHSON, 1840)	56(1)	A17b, App
Leptobium gracile (GRAVENHORST, 1802)		A17b
+Leptobium samium Assıng, 2009	1a(3), 1c(1)	A17b, App
Lobrathium rugipenne (Носннитн, 1851)	11b(2), 23(/2), 23a(2), 31a(1), 60(5)	A17b, App
Medon dilutus pythonissa (SAULCY, 1865)	47a(1), 53(2)	A17b, App
Medon fusculus (MANNERHEIM, 1830)		A17b
Medon lydicus Bordoni, 1980	8(4), 10a(2), 11a(9), 12a(9), 14a(3), 29(1), 30(3), 32a(1), 37b(1), 40a(1)	A17b, App
Medon maronitus (SAULCY, 1865)	52a(1)	A17b, App
Medon semiobscurus (FAUVEL, 1875)	4(1), 8(1), 9a(1), 10(3), 10a(6), 12a(1), 20(2), 24a(3), 32a(1), 40a(2), 53(1), 54(1), 55(1)	A17b, App
Medon subfusculus FAGEL, 1969		A17b
Ochthephilum brevipenne (MULSANT & REY, 1861)	35(2), 40a(2), 45(1)	A17b, App

Species	Localities/samples	References
Paederus littoralis GRAVENHORST, 1802	40a(1)	A17b, App
*Pseudomedon obscurellus (Erichson, 1840)	60(15)	Арр
Rugilus lesbius Assıng, 2005	9a(3), 11a(1), 23(1), 31(2), 31a(5)	A17b, App
*Scopaeus cameroni COIFFAIT, 1968	60(1)	Арр
*Scopaeus gracilis (SPERK, 1835)	23b(2), 60(1)	Арр
Scopaeus haemusensis Frisch, 1997		A17b
+Sunius ambelosicus Assıng, 2015		A17b
+Sunius geiseri Assıng, 2009		A17b
*Throbalium cycladicum (Косн, 1937)	35(2)	Арр
Staphylininae		
*Acylophorus glaberrimus (HERBST, 1784)	41(1)	Арр
Astrapaeus ulmi (Rossi, 1790)		A17b
Erichsonius subopacus (Носннитн, 1851)	31a(1), 60(2)	A17b, App
Gabrius astutoides (Strand, 1946)		A17b
Gabrius nigritulus (GRAVENHORST, 1802)	31(1), 60(1)	A17b, App
*Gauropterus fulgidus (FABRICIUS, 1787)	23(2)	Арр
* Gyrohypnus fracticornis (Müller, 1776)	11a(1)	Арр
Heterothops dissimilis (GRAVENHORST, 1802)	13b(1)	A17b ²⁾ , App
<i>Heterothops</i> sp. (♀)	47a(1)	Арр
Ocypus mus (Brullé, 1832)	18a(1), 60(1)	A17b, App
Orthidus cribratus cribratus (ERICHSON, 1840)	45(1)	A17b, App
Othius laeviusculus Stephens, 1833		A17b
Othius lapidicola Märkel & Kiesenwetter, 1848	1b(1), 4(1), 42(1), 44(2), 46(1), 46a(6), 47a(2), 57(5)	A17b, App
Philonthus concinnus (GRAVENHORST, 1802)		A17b
Philonthus quisquiliarius (GYLLENHAL, 1810)	22(3), 35a(6), 41(5), 41a(4), 45(6), 49(6), 50a(2)	A17b, App
*Philonthus rufimanus rufimanus Erichson, 1840	23(3), 23a(7), 49(4)	Арр
Quedius cohaesus Eppelsheim, 1888	1a(2), 1b(2)	A17b ⁴⁾
Quedius curtidens Smetana, 1967		A17b
*Quedius fissus Gridelli, 1938	60(1)	Арр
Quedius cf. hellenicus Assıng, 2017 (♀)		A17b
Quedius hermonensis COIFFAIT, 1963		A17b ³⁾ , A18c
*Quedius humeralis STEPHENS, 1832	31(2), 60(4)	Арр
Quedius job COIFFAIT, 1963		A17b, App
*Quedius levicollis (Brullé, 1832)	56(4)	Арр
Quedius nemoralis BAUDI DI SELVE, 1848	11a(1), 31a(1), 40a(1), 47a(1), 57(1), 60(3)	A17b, App
Quedius ortrudae Korge, 1971	33(1)	A17b ⁵⁾ , A18c, App
Quedius scintillans (GRAVENHORST, 1806)	31(2), 31a(1), 45(1), 56(22), 60(5)	A17b, App
Quedius semiaeneus (Stephens, 1833)		A17b
Quedius umbrinus Erichson, 1839	31(9), 31a(5),	A17b, App
**Quedius vicinus (Ménétriés, 1832)	41a(2)	Арр
*Stenistoderus cephalotes (KRAATZ, 1858)	35(1), 45(3)	Арр

Species	Localities/samples	References
Xantholinus chiosicus Assıng, 2015	1a(1), 1b(1), 3a(1), 4(2)	A17b, App
Xantholinus rufipennis Erichson, 1839	56(1)	A17b, App
Xantholinus varnensis COIFFAIT, 1972		A17b

Footnotes: ¹⁾ listed as unnamed species in ASSING (2017b); ²⁾ listed as *Heterothops minutus* WOLLASTON, 1860 in ASSING (2017b) (misidentication); ³⁾ listed as *Quedius rugosipennis* FAGEL, 1969 in ASSING (2017b) (synonym); ⁴⁾ listed as *Quedius pseudonigriceps* REITTER, 1909 in ASSING (2017b) (synonym); ⁵⁾ listed as *Quedius nivicola* KIESENWETTER, 1858 in ASSING (2017b) (misidentification).

The Staphylinidae fauna of Samos

According to the latest contribution to the fauna of Samos (Assing 2017b), 157 species, 17 of them endemic, were known from the island. Revisions and a faunistic article published in the meantime (Assing 2018c, 2019a, GILDENKOV 2019) and a previously overlooked revision (MAKRANCZY 2014) resulted in taxonomic changes and records of three additional species.

A field trip conducted to Samos in April 2022 by Volker Brachat (primary focus on Pselaphinae and Scydmaeninae), Thomas Forcke (primary focus on Carabidae), and the author (primary focus on endogean Staphylinidae) yielded more than 1500 specimens belonging to at least 136 species, with nearly 50 species recorded from Samos for the first time, some of them even representing first records from Greece and/or even the whole of Europe (Tab. 2).

Despite exhaustive efforts at taking soil samples (in all more than 50 large samples) and extracting them by soil washing, the results of the 2022 field trip were remarkably poor regarding the endogean fauna, a phenomenon most likely caused by the preceding, exceptionally hot and dry summer of 2021. Only four specimens of Leptotyphlinae, each of them in a different sample, and not a single specimen of *Cephennium* MÜLLER & KUNZE, 1822 or Anillina (Carabidae) were collected, although at least two *Cephennium* and three Anillina species were already known from the island (ASSING 2017b, GIACHINO & VAILATI 2019). Similarly, only few Staphylinidae were found by turning stones, much fewer than during similar field trips in 2015 and 2017.

As many as ten species, two of Aleocharinae, two of Oxytelinae, one of Leptotyphlinae, and five of Scydmaeninae are newly described. Some additional species of Pselaphinae (*Claviger* spp., *Paramaurops* spp., *Trimium* sp., *Tychus* sp.?) and Scydmaeninae (*Euconnus* sp.) are most likely undescribed, but cannot be named for want of males or require a DNA-based approach (*Claviger* spp.).

Currently including a total of 199 named species, 15 of them endemic, plus additional unnamed or unidentified species, the known diversity of the Staphylinidae fauna of Samos is greater than those of all other Aegean islands. For a compilation of the species numbers recorded from other Mediterranean islands see ASSING (2019c). Localities/samples: April 2022: 1: SSW Agios Konstantinos, Oros Ambelos: N-slope, 37°46'33"N, 26°48'38"E, 940 m, grassy clearing with Pinus and Quercus ilex, soil washing, 2.IV.2022, leg. Assing; 1a: same data, but litter and pine bark sifted, and under stones, leg. Brachat & Forcke; 1b: same data, but litter of Quercus ilex sifted, 15.IV.2022, leg. Assing & Forcke; 1c: same data, but under stones, 15.IV.2022, leg. Assing & Forcke; 1d: same data, but dead pine sifted, 15.IV.2022, leg. Assing; 2: SSW Agios Konstantinos, Oros Ambelos: N-slope, 37°46'41"N, 26°48'30"E, 930 m, N-slope with large rocks, pine and Quercus ilex, and fern undergrowth, soil washing, 2.IV.2022, leg. Assing; 3: SSW Agios Konstantinos, Oros Ambelos: N-slope, 37°46'56"N, 26°48'11"E, 910 m, N-slope with Castanea, soil washing, 2.IV.2022, leg. Assing; 3a: same data, but litter sifted, leg. Brachat; 4: SSW Agios Konstantinos, Oros Ambelos: N-slope, 37°47'00"N, 26°48'05"E, 900 m, N-slope with Castanea, soil washing, 2.IV.2022, leg. Assing; 5: SSW Agios Konstantinos, Oros Ambelos: N-slope, 37°47'13"N, 26°47'59"E, 870 m, soil washing beneath old Platanus orientalis, 2.IV.2022, leg. Assing; 6: SSW Agios Konstantinos, Oros Ambelos: N-slope, 37°47'26"N, 26°48'00"E, 770 m, dry stream valley with bushes and small trees, soil washing, 2.IV.2022, leg. Assing; 7: SSW Agios Konstantinos, Oros Ambelos: N-slope, 37°47'30"N, 26°47'51"E, 750 m, Castanea forest, soil washing near large rock, 2.IV.2022, leg. Assing; 8: SSW Agios Konstantinos, Oros Ambelos: N-slope, 37°47'48"N, 26°48'23"E, 370 m, Platanus orientalis litter near small stream sifted, 2.IV.2022, leg. Assing, Brachat & Forcke; Nightingale 9: SE Agios Konstantinos, Valley, 37°47'25"N, 26°49'58"E, 110 m, stream valley with Platanus orientalis, soil washing, 3.IV.2022, leg. Assing; 9a: same data, but litter sifted, leg. Brachat & Forcke; 9b: same data, but dead *Platanus* trunk sifted, leg. Brachat; 10: SE Agios Konstantinos, road to Manolates, 37°47'14"N, 26°49'23"E, 230 m, stream valley with Platanus orientalis, soil washing (scree) near large rock, leg. Assing; 10a: same data, but litter near large rock sifted, leg. Assing, Brachat & Forcke; 11: SE Agios Konstantinos, road to Manolates, 37°47'33"N, 26°49'30"E, 120 m, stream valley with Platanus orientalis, soil washing near large rocks, 3.IV.2022, leg. Assing; 11a: same data, but litter near large rocks sifted, leg. Assing, Brachat &

Forcke; 11b: same data, but stream bank, leg. Assing & Forcke; 12: W Pirgos, 37°43'21"N, 26°45'46"E, 390 m, stream valley on S-slope, soil washing near stream, 4.IV.2022, leg. Assing; 12a: same data, but litter near stream sifted, leg. Assing, Brachat & Forcke; 13: SW Karlovassi, N Kastania, 37°45'36"N, 26°40'55"E, 250 m, stream valley with Platanus, Acer, and Populus, soil washing, 4.IV.2022, leg. Assing; 13a: same data, but litter sifted, leg. Brachat & Forcke; 13b: running on stone, leg. Assing; 14: Oros Kerkis: NW-slope, NE Kalithea, track to Agios Dimitrias, 37°44'35"N, 26°37'02"E, 600 m, stream valley with very old Platanus and old Pinus, soil washing near Platanus trunks, 5.IV.2022, leg. Assing; 14a: same data, but litter near Platanus trunks sifted, leg. Brachat; 15: Oros Kerkis: NW-slope, NE Kalithea, track to Agios Dimitrias, 37°44'35"N, 26°37'02"E, 620 m, stream valley with very old Platanus, soil washing near Platanus and near rocks above track, 5.IV.2022, leg. Assing; 16: Oros Kerkis, NE-slope, 37°44'25"N, 26°39'24"E, 780 m, dry stream valley with pine and Platanus orientalis, soil washing, 6.IV.2022, leg. Assing; 16a: same data, but litter sifted, leg. Assing & Brachat; 17: Oros Kerkis, NE-slope, 37°44'22"N, 26°39'14"E, 740 m, track margin, nest of black Messor, 6.IV.2022, leg. Assing; 18: Oros Kerkis, NE-slope, 37°44'33"N, 26°39'28"E, 740 m, dry stream valley with Quercus ilex and Pinus, soil washing, 6.IV.2022, leg. Assing; 18a: same data, but track margin, under stone, leg. Brachat; 19: Oros Kerkis, NE-slope, 37°44'36"N, 26°39'29"E, 750 m, pine forest, dry stream valley with old Platanus orientalis, soil washing beneath Platanus, 6.IV.2022, leg. Assing; 20: Oros Kerkis, NE-slope, 37°44'42"N, 26°39'21"E, 680 m, dry stream valley near road margin, soil washing (fine gravel), 6.IV.2022, leg. Assing; 21: Oros Kerkis, NE-slope, 37°45'09"N, 26°39'15"E, 640 m, pine forest, dry stream valley with old Platanus orientalis, soil washing, 6.IV.2022, leg. Assing; 21a: same data, but dry stream valley, litter sifted, 6.IV.2022, leg. Brachat; 22: Karlovassi, 37°47'54"N, 26°42'03"E, 5 m, river bank, 6.IV.2022, leg. Forcke; 23: E Karlovassi, 37°47'14"N, 26°43'40"E, 30 m, river bank, 6.IV.2022, leg. Forcke; 23a: same data, but 14.IV.2022, leg. Forcke; 23b: same data, but treading, washing, and floating, 16.IV.2022, leg. Assing & Forcke; 24: W Karlovassi, Potami Waterfalls, 37°47'11"N, 26°40'05"E, 30 m, stream valley with mixed forest, soil washing, 7.IV.2022, leg. Assing; 24a: same data, but litter sifted, 7.IV.2022, leg. Assing & Brachat; 25: Oros Kerkis, NE-slope, 37°45'14"N, 26°39'13"E, 630 m, dry stream valley on slope with old Platanus orientalis, soil washing, 7.IV.2022, leg. Assing; 26: Oros Kerkis, NE-slope, 37°45'24"N, 26°39'34"E, 610 m, ruderal stream valley, soil washing, 7.IV.2022, leg. Assing; 27: Oros Kerkis, NE-slope, just above Agios Nikolaos, 37°46'09"N, 26°39'51"E, 390 m, ruderal slope with chestnut and olive trees, soil washing, 7.IV.2022, leg. Assing; 28: S Agios Konstantinos, 37°48'18"N, 26°48'45"E, 50 m, ruderal trees and bushes, soil washing, 8.IV.2022, leg. **29**: S Agios Assing; Konstantinos, 37°48'09"N,

26°48'46"E, 120 m, stream valley in pine forest with Platanus orientalis, soil washing near rock, 8.IV.2022, leg. Assing; 29a: same data, but litter near rock sifted, 8.IV.2022, leg. Brachat; 30: SE Agios Konstantinos, road to Manolates, 37°47'14"N, 26°49'23"E, 230 m, stream valley with Platanus orientalis, soil washing (scree) near large rock, 8.IV.2022, leg. Assing; 30a: same data, but litter near large rock sifted, 8.IV.2022, leg.Brachat; 31: SE Karlovassi, 37°47'16"N, 26°43'54"E, 60 m, stream bank, under reed, 8.IV.2022, leg. Forcke; 31a: same data, but 14.IV.2022; 32: S Vourliotes, 37°46'02"N, 26°51'23"E, 540 m, stream valley, soil washing near stream and near large rocks (coarse sand), 9.IV.2022, leg. Assing; 32a: same data, but litter near large rocks sifted, leg. Brachat; 33: S Vourliotes, 37°46'06"N, 26°51'09'E, 600 m, soil washing on slope near large rocks (dark soil with stones), 9.IV.2022, leg. Assing; 34: S Vourliotes, 37°45'52"N, 26°50'26'E, 670 m, stream valley in shade of large rock wall, soil washing beneath Quercus ilex (dark soil with stones), 9.IV.2022, leg. Assing; 34a: same data, but Platanus bark and litter around old Platanus orientalis sifted, 9.IV.2022, leg. Brachat; 35: Psili Ammos, Alyki wetland, salt meadow, 37°42'23"N, 27°00'31"E, 1 m, 9.IV.2022, leg. Forcke; 35a: Psili Ammos, Alyki wetland, 37°42'31"N, 27°00'27"E, 5 m, salt meadow, 10.IV.2022, leg. Forcke; 36: S Agios Konstantinos, Manolates, 37°47'02"N, 26°49'28"E, 330 m, soil washing near old Platanus orientalis, 10.IV.2022, leg. Assing; 36a: same data, but dead wood with Lasius brunneus sifted, leg. Assing & Brachat; **36b**: same data, but dead wood with Lasius brunneus sifted, 13.IV.2022, leg. Brachat; 37: S Agios Konstantinos, SW Manolates, 137°46'50"N, 26°49'20"E, 430 m, dry stream valley, soil washing under deciduous trees with rocks (scree), 10.IV.2022, leg. Assing; 37a: same data, but soil washing near large rocks in ivy (dark soft soil with few stones); 37b: same data, but litter on slope with deciduous trees sifted, leg. Brachat; 38: S Agios Konstantinos, SW Manolates, 37°46'50"N, 26°48'53"E, 640 m, dry stream valley, soil washing (stony soil on slope with small trees), 10.IV.2022, leg. Assing; 39: S Agios Konstantinos, SW Manolates, 37°46'47"N, 26°48'56"E, 660 m, soil washing in dry stream valley beneath old Platanus orientalis (stony soil), 10.IV.2022, leg. Assing; 40: S Agios Konstantinos, SW Manolates, 37°46'38"N, 26°48'58"E, 710 m, soil washing beneath old oak near rocks (stony dark soil), 10.IV.2022, leg. Assing; 40a: same data, but oak litter sifted, leg. Brachat; 40b: track margin, under stone with Lasius sp., leg. Brachat; 41: W Psili Ammos, 37°42'43"N, 26°59'06"E, 3 m, wetland, 10.IV.2022, leg. Forcke; 41a: same data, but 11.IV.2022; 42: SE Paleokastro, 37°43'26"N, 27°02'40"E, 110 m, grassy slope with old oak tree. soil washing, 11.IV.2022, leg. Assing; 43: SE Paleokastro, 37°43'24"N, 27°02'34"E, 80 m, dry stream valley, soil washing on slope beneath Quercus ilex, 11.IV.2022, leg. Assing; 44: SE Paleokastro, 37°43'25"N, 27°02'29"E, 80 m, grassy slope with young Quercus ilex, soil washing (small sample), 11.IV.2022, leg. Assing; 45: W Pythagorio, 37°41'29"N, 26°55'44"E, 5 m, swamp, 11.IV.2022, leg. Forcke; 46: SE Paleokastro, 37°43'31"N, 27°03'22"E, 200 m, calcareous grassy slope with abandoned olive and other trees, soil washing near dead tree (shallow reddish soil with small stones), 12.IV.2022, leg. Assing, 46a: same data, but litter near rocks and old olive tree sifted, leg. Assing & Brachat; 47: W Vathi, Kamara, 37°44'55"N, 27°00'30"E, 150 m, between large vertical rock and road margin, soil washing below Quercus ilex (deep soft black soil with large stones), 12.IV.2022, leg. Assing; 47a: same data, but litter of Quercus ilex sifted, 12.IV.2022, leg. Assing & Brachat; 48: N Vathi, coast road, 37°47'12"N, 26°57'29"E, 10 m, slope with various deciduous trees, bushes, and Rubus, soil washing (deep soft black soil with large stones), 12.IV.2022, leg. Assing; 49: E Myli, 37°40'39"N, 26°52'19"E, 15 m, river bank, 12.IV.2022, leg. Forcke; 50: W Psili Ammos, 37°42'32"N, 26°59'20"E, 3 m, reed and salt meadow 12.IV.2022, leg. Forcke; 50a: same data, but 13.IV.2022; 51: S Agios Konstantinos, E Manolates, 37°46'57"N, 26°49'51"E, 370 m, dark stream valley, soil washing (deep stony soil), 13.IV.2022, leg. Assing; 51a: same data, but litter sifted, leg. Brachat; **52**: S Agios Konstantinos, E Manolates, 37°46'51"N, 26°49'55"E, 380 m, dark dry stream valley, soil washing (stony soil), 13.IV.2022, leg. Assing; 52a: same data, but soil washing under dead trunk (dark stony soil); 52b: same data, but litter under dead trunk sifted, leg. Brachat; 53: S Agios Konstantinos, E Manolates, 37°46'47"N, 26°50'02"E, 380 m, soil washing beneath oak tree, 13.IV.2022, leg. Assing; 53a: same data, but under stone, leg. Brachat; 54: S Agios Konstantinos, E Manolates, 37°46'35"N, 26°49'55"E, 420 m, stream valley, soil washing, 13.IV.2022, leg. Assing; 55: S Agios Konstantinos, 37°47'13"N, 26°49'46"E, 280 m, grassy slope with trees and bushes, soil washing beneath tree (deep dark stony soil), 13.IV.2022, leg. Assing; 56: W Psili Ammos, 37°42'43"N, 26°59'18"E, 10 m, arable land, 13.IV.2022, leg. Forcke; 57: Oros Kerkis, 37°43'26"N, 26°38'10"E, 1210 m, litter, grass, and moss beneath Juniperus and cushion plants sifted, 14.IV.2022, leg. Assing; 58: Oros Kerkis, 37°45'N, 26°40'E, 820 m, under stone, Cataglyphis nodus nest, 14.IV.2022, leg. Brachat; 59: Agios Konstantinos, 37°48'13"N, 26°49'37"E, 20 m, stream bank, 15.IV.2022, leg. Forcke; 60: W Karlovassi, Potami, 37°47'11"N, 26°40'05"E, 30 m, stream bank, treading, washing, and floating, 16.IV.2022, leg. Assing & Forcke.

Notes on some species and additional records

Euplectus meybohmi Brachat, 2017

According to Volker Brachat (e-mail 22 July, 2022), the elytra have four basal foveae, not three as indicated in the original description.

Tychus kerkisicus (BRACHAT, 2017), comb. nov.

Paratychus kerkisicus BRACHAT, 2017: 107.

This species was originally attributed to *Paratychus* BESUCHET, 1960 primarily because of the elongated maxillary palpomeres IV. Based on a reassessment of other external and sexual characters, however, it belongs to the genus *Tychus* LEACH, 1817 (Volker Brachat, e-mail 22 July, 2022).

Liogluta aloconotoides (BENICK, 1940), comb. nov. (Figs 4, 24)

Atheta (Glossola) aloconotoides BENICK, 1940: 180 f.

Aloconota (Aloconota) aloconotoides: SCHÜLKE & SMETANA (2015)

Type material examined: Holotype \mathfrak{P} : " \mathfrak{P} / Trento, Pechlaner / Ci. Tombea, 13.6.35 / Atheta grandicornis Fauv. / Atheta aloconotoides \mathfrak{P} , Typus, G. Bck / wohl zu *Liogluta* J. V. / aloconotoides, det. Zanetti '96 / MHNG ENTO 00096512 / Liogluta aloconotoides (Benick), det. V. Assing 2022" (MHNG).

Comment: The original description is based on a female holotype and an unspecified number of paratypes ("in mehreren Exemplaren") from "Ci. Tombea" (BENICK 1940). In order to clarify if the female of *Liogluta* sp. (see Tab. 2) is conspecific with this species, the holotype was examined. It undoubtedly belongs to *Liogluta* THOMSON, 1858. The habitus and the shape of the spermatheca of the holotype are illustrated in Figs 4, 24.

Ischnoglossa (Xenodota) minor (AUBÉ, 1863)

When preparing a taxonomic note on the genus *Ischno-glossa* KRAATZ, 1856, a female of *I. minor* from Samos, the first record of this species from Greece, was examined (ASSING in press). The data are as follows: $1 \, \varphi$, Samos, N Psili Ammos, 37°43'N, 26°59'E, 90 m, under bark near stream, 19.IV.2003, leg. Brachat & Meybohm (cAss).

Meotica truncata Assing, 2004

Meotica truncata was originally described based on material from Antalya (AssING 2004b) and subsequently reported also from other localities in various South Turkish provinces (Izmir, Aydın, Isparta, Niğde, Mersin, Kahramanmaraş, Kayseri) (ASSING & VOGEL 2019). The specimens from Samos represent the first records from Samos, Greece, and Europe.

Oxypoda flavocaudata Assing, 2006

This species was previously known only from South Turkey, from Muğla in the west to Mersin in the east (AssING 2006, 2007). The specimens from Samos represent the first records from Samos, Greece, and Europe.

Oxypoda gladiatoria Assing, 2006

The previously known distribution was confined to South Turkey (Muğla, Antalya) and Cyprus (Assing 2006). The specimens from Samos represent the first records from Samos, Greece, and Europe.

Oxypoda rectacia Assing, 2019

According to a recent revision (AssING 2019a), the material listed in AssING (2017b) as *Oxypoda* cf. *nova* BERNHAUER, 1902 belongs to *O. rectacia*.

Carpelimus simplicicollis (WOLLASTON, 1857)

The female listed in AssING (2017b) as *Carpelimus* sp. was identified by GILDENKOV (2019) as *C. simplicicollis*.

Ochthephilus andalusiacus (FAGEL, 1957)

This species was not listed in Assing (2017b), but had been recorded from Samos by MAKRANCZY (2014).

Stenichnus cribrum (SAULCY, 1880)

According to SCHÜLKE & SMETANA (2015), this species had been reported only from the Caucasus region (Georgia; Russian South European territory). The specimens from Samos represent the first records from Greece.

Throbalium cycladicum (Косн, 1937) (Figs 40–57)

Lathrobium (Throbalium) cycladicum Косн, 1937: 260 ff. Lathrobium (Throbalium) biblicum Косн, 1937: 262 f.; syn. nov. Throbalium adanense Косн, 1939: 436 f.; syn. nov. Throbalium kosianum BORDONI, 2020: 223 f.; syn. nov.

Type material examined: *Lathrobium cycladicum*: Holotype σ [abdomen (except segment VIII) and aedeagus missing]: "Naxos, Graecia / Prep. microscp. No. 1429 / Cotype" (NHMB).

Lathrobium biblicum: Holotype ♂ [aedeagus missing]: "96 / Attlith, Palestine, 22–27.8.1935, A. Rabinovitch cep.

/ Lathrobium (Throbalium) biblicum Koch, det. C. Koch / Type / Holotype 1956 det. Kamp / Coll. G. Frey NMB / Throbalium cycladicum (Koch), det. V. Assing 2022" (NHMB).

Throbalium adanense: Holotype ♂: "Adana, Asm., 4.VI.1934, Neubert / Museum Frey Tutzing / Th. adanense Koch, det. C. Koch / Type / Coll. G. Frey NMB / Throbalium cycladicum (Koch), det. V. Assing 2022" (NHMB).

Additional material examined: GREECE: mainland: 1 J, Pieria, E Korinos, 40°19'N, 22°38'E, 0 m, salt meadow, 20.X.2021, leg. Forcke (cAss); 3 9 9, Chalkidiki, Kassandra, 15.VIII.1994, leg. Eifler (cAss); 6 ♂ ♂, 1 ♀, Makhedonía, Serrai, Strimonas delta, 28.VIII.1998 (cAss); 1 &, Thráki, Nomós Évros, Évros delta, Salicornia, 22.IV.1994, leg. Bense (cAss). Samos: see Tab. 2. Kos: 1 d, Tigaci, 18.IX.1989, leg. Jørum (cAss). CYPRUS: 1 d', Lemesos salt lake, 5 m, 9.III.1996, leg. Frisch (cAss). TURKEY: Mersin: 1 J, 1 9, Silifke, 1990, leg. Lödl (NHMW, cAss); 1 º, S Silifke, Göksu delta, 36°20'N, 34°01'E, 9.V.2004, leg. Brachat & Meybohm (cAss). Adana: 1 d, Ceyhan, Devecusta delta, 13–15.IV.1992, leg. Hovorka (cAss). Kayseri: 1 J, SW Seyhşaban, 38°28'N, 35°18'E, 1150 m, 11.VI.1998, leg. Bartsch & Salmen (cAss). ISRAEL: 1 &, Dead Sea region, lake shore near Newe Zohar, 31°09'N, 35°22'E, -350 m, 27.II.2009, leg. Aßmann (cFel).

Comment: KOCH (1937) failed to specify any type material of *Lathrobium cycladicum*. It can be inferred from the specific epithet that the syntype(s) he based the description on originated from the Cyclades. Only one male syntype from Naxos was found in the Koch collection at NHMB. Nearly all of the abdomen and the aedeagus is missing. The latter was apparently mounted on a slide and separated from the specimen, most likely by H. Coiffait.

The original description of *L. biblicum* is based on a male and a female from "Attlith (südlich von Haifa, Palästina)". In stating that he illustrated the aedeagus of the type in the caption of figure 10, KOCH (1937) designated the male as the holotype. The aedeagus of the examined holotype is missing, suggesting that it was lost or separated from the specimen either by C. Koch or by H. Coiffait. *Throbalium adanense* was described based on a unique male holotype from "Adana" (KOCH 1939). Very recently, BORDONI (2020) described *Throbalium kosianum* based on two males from the Aegean island Kos.

A comparison of the examined type material, of additional material from Greece (including Kos), Cyprus, Turkey, and Israel, and of the illustrations provided with the respective original descriptions revealed some intraspecific variation of the shapes of the male sternites VII and VIII (Figs 48–57), but yielded no evidence that *T. cycladicum*, *T. biblicum*, *T. adanense*, and *T. kosianum* should refer to different species. Similarly, no significant differences were found in the structure of the aedeagus (Figs 40–47). However, there appears to be some clinal variation with the ventral process of the aedeagus tending to become broader towards the east of the distribution. In consequence, *T. cycladicum* is designated as the senior synonym of *T. biblicum*, which was described in the same article. *Throbalium adanense* and *T. kosianum* were made available subsequently and are thus junior synonyms of *T. cycladicum*.

Quedius cohaesus Eppelsheim, 1888

This species was listed as *Q. pseudonigriceps* REITTER, 1909 in Assing (2017b). The name was recently synonymised with *Q. cohaesus* by Assing (2019e).

Quedius hermonensis COIFFAIT, 1963

This species was listed as *Q. rugosipennis* FAGEL, 1969 in ASSING (2017b). The name was recently synonymised with *Q. hermonensis* by ASSING (2018c).

Quedius job COIFFAIT, 1963

The previously known distribution included Lesbos, Samos, Karpathos, South Turkey, and Lebanon (ASSING 2018c). Some paratypes of *Q. lydicus* FAGEL, 1968, a name recently synonymised with *Q. job* by ASSING (2018c), had been collected in Samos (FAGEL 1968). Additionally, the following material was examined from Samos: $2 \sigma \sigma$, $1 \circ$, Valeonates, Nightingale Valley, 37.784°N, 26.818°E, 70 m, stream bank, 9.VI.2013, leg. Degasperi (cDeg, cAss); 1 ex., Mili env., Imuresos, 5.IV.2008, leg. Rossi (cAss).

Quedius ortrudae Korge, 1971

According to a recent revision (AssING 2018c), this species was previously confounded with *Q. nivicola* KIESENWETTER, 1858, as which it was recorded from Samos by AssING (2017b).

Quedius vicinus (Ménétriés, 1832)

The previously known distribution included Turkey, the Middle East, the Caucasus region, and Middle Asia (SCHÜLKE & SMETANA 2015). The material from Samos (and Rhodos; see the section on the Staphylinidae of Rhodos) represents the first records from the Aegean Islands (Samos, Rhodos) and Greece.

Descriptions of new species

Euphanias ambulans ASSING spec. nov. urn:lsid:zoobank.org:act:4D363881-A549-4AA0-A824-66E2EBA02BD3 (Figs 3, 18–20)

Type material: Holotype \Im : "GREECE: Samos [35a], Psili Ammos, Alyki wetld, 37°42'31"N, 27°00'27"E, 5 m, salt meadow, 10.IV.2022, T. Forcke / Holotypus \Im *Euphanias ambulans* sp. n., det. V. Assing 2022" (cAss). Paratype \Im : same data as holotype (cAss).

Etymology: The specific epithet is the present participle of the Latin verb ambulare (to walk) and alludes to the completely reduced hind wings.

Description: Body length 2.7–2.9 mm; length of forebody 1.4–1.6 mm. Habitus as in Fig. 3. Colouration: body, including the appendages, blackish; whole body, including eyes, covered with dense, somewhat irregularly distributed pale scales.

Head (Figs 19-20) strongly domed in median dorsal portion, this elevation sharply delimited from (and by) lateral and posterior sulci; above antennal insertion strongly domed. Eyes large, strongly convex, and composed of coarse ommatidia, approximately three times as long as the distance between posterior margin of eye and posterior constriction of head in dorsal view. Antenna (Fig. 18) 1.0-1.1 mm long; antennomeres I large and long, approximately three times as long as broad, subapically constricted and apically somewhat flattened, covered with moderately dense scales not concealing all of integument, II much narrower and shorter than I and nearly twice as long as broad, III-IV of subequal length, very slender, strongly oblong, and longer and narrower than II, V of similar shape and II and III, but slightly longer, VI-VII much shorter than III-V, less slender, of conical shape, and approximately twice as long as broad, VIII of similar shape as VI-VII but slightly less oblong, IX significantly larger than VIII, of distinctly conical shape, and approximately as broad as long; X transverse, approximately 1.5 times as broad as long, and XI transverse and apically constricted.

Pronotum (Figs 19–20) 0.55–0.60 mm broad, 1.29–1.36 times as broad as long and 1.31–1.33 times as broad as head; middle of anterior margin, lateral and posterior margins strongly bulging; middle of disc with pair of pronounced oblong and narrowly separated elevations, area around these elevations deeply impressed.

Elytra (Figs 19–20) 1.17–1.19 times as long as pronotum, each elytron 1.55–1.60 times as long as broad and with a sutural, a humeral, and two intermediate carinae, all of them pronounced and reaching posterior margin of elytra (sutural and third carina), or nearly so (second and humeral carina); humeral carina most pronounced basally, gradually becoming less pronounced posteriad. Hind wings completely reduced.

Abdomen not distinctive.

♂: unknown.

Comparative notes: The enigmatic genus *Euphanias* FAIRMAIRE & LABOULBÈNE, 1856 was previously represented in the Palaearctic region by only four species, all of them extremely rarely found: *E. insignis* (MULSANT & REY, 1856) (North Africa: Algeria, Tunisia, Egypt; France, Italy; Middle East: Syria, Saudi Arabia) with the subspecies *E. insignis insularis* KOCH, 1934 endemic to Sardinia; *E. boiteli* NORMAND, 1936 (Tunisia); *E. pliginskii* BERNHAUER, 1912 (Ukraine; Russian West Caucasus); *E. pusanovi* BLINSTEIN, 1976 (Ukraine). Using the key to species in BLINSTEIN (1976), *E. ambulans* would key out together with *E. pliginskii* and *E. pusanovi*, mainly based on the longer antennomere I (*E. insignis* and *E. boiteli*: antennomere I only twice as long as broad). It is distinguished from *E. pliginskii* and *E. pusanovi* as follows:

- from *E. pusanovi* by larger size (*E. pusanovi*: body length 2.2 mm), significantly larger eyes (*E. pusanovi*: eyes twice as long as postocular region), and much shorter median elevations of the pronotum (*E. pusanovi*: median elevations reaching anterior and posterior margins);

- from *E. pliginskii* by shorter elytra (*E. pliginskii*: elytra > 1.25 times as long as pronotum; each elytron approximately 1.65 times as long as broad) with a less oblique humeral carina, more oblong and more distinctly separated median elevations of the pronotum, and longer and more slender antennomeres III–X.

For illustrations of *E. pusanovi* and *E. pliginskii* see BLINSTEIN (1976) and https://www.zin.ru/animalia/ coleoptera/eng/eupplikg.htm.

Distribution and natural history: The type locality is situated near Psili Ammos, Southeast Samos. The specimens were hand-collected in a salt meadow near sealevel.

Hydrosmecta cultellata ASSING spec. nov. urn:lsid:zoobank.org:act:014960F5-7A69-4D33-8B6F-2F543D07AF7D (Figs 5, 25–32)

Type material: Holotype σ : "GREECE: Samos [60], W Karlovassi, Potami, 37°47'11"N, 26°40'05"E, 30 m, stream bank, 16.IV.2022, V. Assing / Holotypus σ *Hydrosmecta cultellata* sp. n., det. V. Assing 2022" (cAss). Paratypes: 3 φ φ : same data as holotype (cAss).

Etymology: The specific epithet is an adjective derived from the Latin noun cultellum (small knife). It alludes to the shape of the conspicuous process at the base of the ventral process of the aedeagus.

Description: Body length 2.5–2.8 mm; length of forebody 1.1–1.2 mm. Habitus as in Fig. 5. Colouration: body blackish; legs yellow; antennae blackish.

Head weakly transverse; punctation fine and moderately dense; interstices with shallow microsculpture. Eyes large, longer than distance from posterior margin of eye to posterior constriction of head in dorsal view. Antenna 0.8–0.9 mm long; antennomeres III–IV weakly oblong, V–X weakly oblong or as long as broad.

Pronotum 1.15–1.20 times as broad as long and approximately 1.1 times as broad as head; punctation and microsculpture similar to those of head; pubescence along midline directed anteriad, posteriad only near posterior margin.

Elytra slightly longer than pronotum; punctation extremely dense and fine, barely visible in the pronounced microsculpture even at high magnification (100 x).

Abdomen with dense and fine punctation; interstices with microsculpture.

 σ : tergite VIII transverse, posterior margin weakly concave in the middle (Fig. 27); sternite VIII longer than tergite VIII, posterior margin strongly convex (Fig. 28); median lobe of aedeagus (Figs 29–30) 0.27 mm long and of very distinctive shape, at base of ventral process with additional knife-shaped process (best visible in lateral view).

 $\$: tergite VIII of similar shape as in male (Fig. 31); sternite VIII (Fig. 32) slightly longer than tergite VIII, posterior margin distinctly concave in the middle; spermatheca shaped as in Figs 25–26, with long apical cuticular invagination.

Comparative notes: This species is reliably distinguished from its congeners only by the shapes of the distinctive aedeagus and of the spermatheca.

Distribution and natural history: The type locality is situated near Potami Waterfalls to the west of Karlovassi, Northwest Samos. The specimens were floated from fine gravel on a stream bank at an altitude of 30 m.

On the species of the *Atheta* (*Philhygra*) *elongatula* group

In a comprehensive and, especially regarding the illustrations, superb revision of the *Atheta* subgenus *Hygroecia* MULSANT & REY, 1873, now a junior synonym of *Philhygra* MULSANT & REY, 1873, BRUNDIN (1944) described three subspecies of *Atheta elongatula* (GRAVENHORST, 1802), all of them currently regarded as distinct species: *Atheta cretica*, *A. balcanica* (a preoccupied name subsequently replaced with *A. balcanicola* SCHEERPELTZ, 1968), and *A. transcaspica* (today a junior synonym of *A. laevigata* (HOCHHUTH, 1849)). An additional species belonging to this group, *A. monstruosa*, was rather recently described by ASSING (2011). In external and the secondary sexual characters, the species of the *A. elongatula* group are highly similar, so that a reliable identification is possible only based on the shape of the aedeagus.

While *A. elongatula* is widspread in nearly all of the Palaearctic region from West Europe in the west to East Siberia and North Korea in the east, the other species of this group have more restricted distributions, with *A. cretica* endemic to Crete, *A. balcanicola* distributed in

the Balkans and southern Central Europe, *A. monstruosa* recorded from North Turkey and Armenia, and the previously known distribution of *A. laevigata* ranging from the East Aegean Islands across Turkey to the Caucasus region.

Doubts regarding the identity of the specimens collected in Samos, together with the somewhat implausible distribution of *A. laevigata*, initiated a thorough revision of material previously identified as *A. cretica*, *A. balcanicola*, and *A. laevigata*. This study revealed that what had previously been recorded as *A. laevigata* from the Aegean Islands and South Turkey represented a distinct species distinguished from *A. laevigata* by slight, but constant differences in the shape of the aedeagus and also by external characters. This species is described below. Confirmed records of *A. laevigata* were seen only from the Caucasus region (Russian West Caucasus, Armenia, Northeast Turkey).

For an adequate assessment of the shape of the aedeagus, it should be noted that the large apical structure of the median lobe may be more or less extruded, thus sometimes conveying a misleading impression of aedeagal size and shape. Therefore, measurements of the aedeagus are always given as the length from the apex of the ventral process to the base of the aedeagal capsule.

For comparison, new illustrations of the aedeagi of *A. balcanicola*, *A. cretica*, *A. laevigata*, and *A. monstruosa* are provided in Figs 33–36.

Atheta (Philhygra) tecta ASSING spec. nov. urn:lsid:zoobank.org:act:2794087C-8BD2-4278-A358-D0CE8A62C1DB (Figs 6, 37–39)

Type material: Holotype ♂: "GR – Samothraki [16], W Lakkoma, 40°25'57"N, 25°30'54"E, 20 m, gravel river bank, 9.IV.2019, V. Assing / Holotypus & Atheta tecta sp. n., det. V. Assing 2022" (cAss). Paratypes: 1 ♂, 2 ♀ ♀: same data as holotype (cAss); 2 ♂ ♂, 2 ♀ ♀: "GR – Samothraki [38a], SE Kamariotissa, 40°26'48"N, 25°29'53"E, 40 m, stream gravel, 16.IV.2019, V. Assing" (cAss); 1 J, 1 9: "GREECE: Samos [41], W Psili Ammos, 37°42'43"N, 26°59'06"E, 3 m, wetland, 10.IV.2022, T. Forcke" (cAss); 2 ♂ ♂, 1 ♀: same data, but "[41a] ... 11.IV.2022" (cAss); 1 9: "GREECE: Samos [50a], W Psili Ammos, 37°42'32"N, 26°59'20"E, 3 m, reed & salt meadow 13.IV.2022, T. Forcke" (cAss); 1 ♂: "GREECE: Samos [56], W Psili Ammos, 37°42'43"N, 26°59'18"E, 10 m, arable land, 13.IV.2022, T. Forcke" (cAss); 3 ♂ ♂, 1 ♀: "GREECE: Kos [14], 9.6 km WSW Kos, NW Zia, 36°50'58"N, 27°11'43"E, 210 m, leaf litter sifted, 26.XII.2016, V. Assing" (cAss); $13 \circ \circ$, $7 \circ \circ$: "TR. - Muğla; No. 13, SE Dalaman, 10 m, flood-plain wood, meadows, 36°47'02N, 28°50'03E, 28.III.2002, V. Assing" (cAss, MNB); 1 &: "N37°16' E028°10', Türkei Umg. Mugla, Bayir, 400 m, Meybohm 1.5.2001" (cAss); 1 ♂, 2 ♀ ♀: "TR [13a]- Konya, 35 km ENE Beyşehir, **Etymology**: The specific epithet (Latin, adjective: hidden, secret) alludes to the fact that this species was previously confounded with *Atheta laevigata*.

Comment: The above type specimens from Kos and Samothraki were reported as *A. laevigata* by Assing (2017a, 2019b).

Description: Body length 3.5–4.2 mm; length of forebody 1.5–1.8 mm. Habitus as in Fig. 6. Colouration: head black; pronotum dark-brown to black; elytra yellowish-brown to dark-brown; abdomen black, usually with the apex (segments IX–X and posterior portion of segment VIII) yellowish-brown to reddish-brown; legs yellow; antennae blackish-brown to black. Head and pronotum with fine, moderately dense punctation and with weak shine owing to very fine and shallow microsculpture. Other external characters not distinctive. ♂: sternite VIII longer than tergite VIII, posterior margin strongly convex; median lobe of aedeagus relatively small, 0.40–0.47 mm long, and shaped as in Figs 37–39.

Comparative notes: Regarding the shape of the aedeagus, *A. tecta* most resembles *A. cretica* and *A. laevigata*. The new species is distinguished from both species by a smaller and less robust body and by a smaller aedeagus with a smaller crista apicalis, an apically less strongly curved ventral process (lateral view), a ventro-apically more longer and more acute apical structure of the median lobe (lateral view), from *A. laevigata* additionally by paler colouration (especially of the elytra; *A. laevigata*: elytra usually black) and by a less matt forebody. For illustrations of the aedeagi of the other species of the *A. elongatula* group see Figs 33–36 and BRUNDIN (1944).

Distribution and natural history: The known distribution ranges from the East Aegean Islands (Samothraki, Samos, Kos) in the west across South Turkey (Muğla, Konya) northeastwards to Tokat. The specimens were collected on stream and river banks, on the shore of reservoirs, in wetlands and salt meadows, in a floodplain forest, and on arable land. The altitudes range from near sealevel to 1270 m. *Cretotyphlus kerkisicus* ASSING spec. nov. urn:lsid:zoobank.org:act:16BD4F7A-B7A6-47B3-8025-342CAD1855EF (Figs 7, 58–59)

Type material: Holotype ♂: "GREECE: Samos [18], Oros Kerkis, NE-slope, 37°44'33"N, 26°39'28"E, 740 m, soil washing, 6.IV.2022, V. Assing / Holotypus ♂ *Cretotyphlus kerkisicus* sp. n. det. V. Assing 2022" (cAss).

Etymology: The specific epithet is an adjective derived from Kerkis, the name of the mountain on whose north slope the holotype was collected.

Description: Body length 1.35 mm (abdomen extended), length of forebody approximately 0.6 mm. Habitus as in Fig. 7. Colour of body dark-yellowish.

Head with sparse, barely visible punctation and with very shallow microsculpture. Pronotum approximately as long as broad and slightly narrower than head; microsculpture slightly more distinct than that of head.

Abdomen with shallow microsculpture composed of large meshes; punctation extremely fine and moderately dense; posterior margin of tergite VIII truncate.

 σ : sternite VIII (Fig. 59) nearly symmetric, weakly oblong, with unmodified chaetotaxy, and with a broadly concave posterior excision; aedeagus (Fig. 58) 0.28 mm long, weakly asymmetric; ventral process long, slender, and bisinuate, with a pronounced ventral tooth approximately in the middle.

 $^{\mathrm{Q}}\colon unknown.$

Comparative notes: The genus *Cretotyphlus* AssING, 2019 previsously included three species, all of them locally endemic to Crete. Regarding the shape of the aedeagus, *C. kerkisicus* is very similar to *C. hamatus*, the type species of the genus. It differs from this species by a more slender and strongly bisinuate ventral process of the aedeagus and a slightly more oblong male sternite VIII with unmodified pubescence. For illustrations of *C. hamatus* see AssING (2019d).

Distribution and natural history: Despite extensive soil sampling, *C. kerkisicus* was found in only one locality on the northeastern slope of Oros Kerkis, West Samos. The holotype was collected by washing soil from a dry stream valley with *Quercus ilex* and *Pinus* sp. at an altitude of 740 m.

Cephennium amplexans ASSING spec. nov. urn:lsid:zoobank.org:act:41A5DF62-9FFC-44D3-A57C-20B2B93720B9 (Figs 10, 60)

Type material: Holotype ♂: "GR Samos, Avlakia 100 m, N37°47' E26°51' / 28.4.2003, leg. Brachat & Meybohm, aus Streu und Boden in Bachnähe / Holotypus ♂ *Cephennium amplexans* sp. n. det. V. Assing 2022" (cAss). Paratypes:

1 ♂: "GR Samos, Psili Amos 1 m, N37°42' E27°00' / 20.4.2003, leg. Brachat & Meybohm, am Rand der Saline unter Mastix-Strauch" (cAss); 2 ♀ ♀: "GR Samos, Manolates 300 m, N37°47' E26°49' / 29.4.2003, leg. Brachat & Meybohm, aus Streu und Boden unter Sträuchern" (cAss); 1 ♂: "GR Samos, Kosmadei 500 m, Höhle Tsitse Tripa, N37°45' E26°39' / 23.4.2003, leg. Brachat & Meybohm, bei der Höhle" (cAss); 1 ♂: "GR Samos, Platanakia 100 m, N37°47' E26°50' / 29.4.2003, leg. Brachat & Meybohm, Nachtigallental" (cAss).

Etymology: The specific epithet is the present participle of the Latin verb amplexare (to encompass, to embrace). It alludes to the shapes of the internal structures of the aedeagus, with one of the claw-shaped structures enveloping the other.

Description: Body length 0.8–1.0 mm. Habitus as in Fig. 10. Punctation of pronotum more distinct than that of elytra. Eye rudiments minute, composed of 2–4 ommatidia without pigmentation. Other external and male secondary sexual characters as in other species of the *C. unguicum* subgroup of the *C. corruptum* group.

 σ : aedeagus (Fig. 60) approximately 0.23 mm long, broad, and apically weakly pointed in the middle; internal sac with a pair of strongly curved claw-shaped structures; parameres apically extending slightly beyond apex of median lobe.

Comparative notes: Based on the similar external characters and particularly the similarly derived structure of the aedeagus (internal sac with two claw-shaped structures), C. amplexans undoubtedly belongs to the C. unguicum subgroup of the C. corruptum group (see Assing & Meybohm 2021a). Among the species of the C. unguicum subgroup, whose previously known representatives are all distributed in Southwest Turkey, the aedeagus of C. amplexans is most similar to that of C. unguicum Assing & MEYBOHM (2021) from Dilek Dağı, a mountain in the Turkish province Aydın separated from the east coast of Samos only by the narrow Mycale Strait. However, in C. unguicum, the claw-shaped structures are of nearly identical shape and the left structure is much less strongly curved. For illustrations of the aedeagi of C. unguicum and other species of the C. ungui*cum* subgroup see Assing & Меувонм (2021а).

Based on the morphology of the aedeagus and external characters, *C. amplexans* is closely allied neither to any of the species known from mainland Greece (see ASSING & MEYBOHM 2021b), nor to those from Crete and Samothraki (ASSING 2019b, ASSING et al. 2019).

Distribution and natural history: The type material was found in several localities across Samos at altitudes of close to sea-level to 500 m. The specimens were sifted from litter and soil near a stream, near a salt-meadow, and near a cave. *Cephennium monstrans* ASSING spec. nov. urn:lsid:zoobank.org:act:86CB13EF-3F05-45A1-BB47-034707B226A5 (Figs 11, 61)

Type material: Holotype ♂: "GR Samos, 430 m, 6 km westl. Pirgos, N37°43' E26°45' / 24.4.2003, leg. Brachat & Meybohm, Bachtal, Kiefernwald / Holotypus ♂ *Cephennium monstrans* sp. n. det. V. Assing 2022" (cAss). Paratype ♂: "GR Samos, Platanakia 100 m, N37°47' E26°50' / 29.4.2003, leg. Brachat & Meybohm, Nachtigallental" (cAss).

Etymology: The specific epithet is the present participle of the Latin verb monstrare (to show, to point). It alludes to the shapes of the internal structures of the aedeagus, both of which seem to point in the same direction.

Description: Body length 0.85–0.90 mm. Habitus as in Fig. 11. Punctation of pronotum more distinct than that of elytra. Eye rudiments minute, composed of 2–3 ommatidia without pigmentation. Other external and male secondary sexual characters as in *C. amplexans* and other species of the *C. unguicum* subgroup.

 σ : aedeagus (Fig. 61) approximately 0.23 mm long, broad, and apically truncate; internal sac with a pair of claw-shaped structures, both of them pointing to the left in ventral view.

Comparative notes: Like *C. amplexans, C. monstrans* belongs to the *C. unguicum* subgroup. It is reliably distinguished from *C. amplexans* and other species of the *C. unguicum* subgroup only based on the shapes of the internal structures of the aedeagus.

Distribution and natural history: The type material was found in two localities in Central and North Samos, in one of the localities together with *C. amplexans*. The specimens were sifted from litter and soil at altitudes of 100 and 430 m.

Stenichnus (Stenichnus) samius MEYBOHM spec. nov. urn:lsid:zoobank.org:act:D281CE0D-EEA6-45E7-BCB2-21734004DD80 (Figs 14, 64)

Type material: Holotype σ : "GR Samos Potami 20 m N37°47' E26°40' / leg. 22.4.2003 Brachat & Meybohm im Schluchtwald / *Stenichnus samius* m. Meybohm 2022 det. / Holotypus" (cMey). Paratypes: 1 \circ : same data as holotype; 1 \circ : "GREECE: Samos [24a], W Karlovassi, Potami, 37°47'11"N 26°40'05"E, 30 m, litter sifted, 7.IV.2022, V. Assing" (cMey); 1 σ : "N37°45'36 E26°40'55 GR Samos (13a) Kastania N 250 m leg. Brachat 4.4.2022" (cMey).

Etymology: The specific epithet (Latin, adjective) alludes to the fact that this species is most likely endemic to Samos.

Description: Body length \circ 1.50–1.60 mm, \circ 1.65–1.73 mm. Male habitus as in Fig. 14. Body weakly bicolored; head and pronotum brown, elytra black; legs, antennae, and palpi light brown.

Head strongly transverse, 1.3 times as broad as long, 0.8 times as broad as pronotum; head width 0.30 mm (σ) to 0.31 mm (φ), head length 0.23 mm (σ) to 0.24 mm (φ); postocular region as long as diameter of eyes; vertex and frons distinctly flattened, with punctures and pubescence dimorphic; punctures in female distinctly finer and sparser than in male; setae in female long, in male on vertex shorter and between eyes missing; supra-antennal tubercles very weakly pronounced. Eyes composed of approximately 20 ommatidia, in male slightly larger and with slightly more ommatidia than in female. Antenna slender, 0.85 mm (σ) to 0.90 mm (φ) long; antennomeres I–VII elongate, up to twice as long as broad, VIII about as long as broad, IX and X moderately transverse, XI distinctly shorter than the combined length of IX and X.

Pronotum weakly oblong, 0.41–0.42 mm long and 0.36– 0.39 mm broad in male, 0.43 mm long and 0.39 mm broad in female; punctures very fine; pubescence very long and suberect; setae directed mediad in lateral portion of posterior half, otherwise directed posteriad; basally only with two minute pits separated from each other by half the distance as between pits and posterior margin.

Elytra broadly oval, dorsally flattened, 0.95–1.04 mm long and 0.68 mm broad in male, 1.06–1.10 mm long and 0.71–0.72 mm broad in female, broadest at anterior third, height approximately one-fourth of elytral length (lateral view); humeral angle and humeral fold absent; at base of each elytron with a large deep inner impression, lateral impressions absent; pubescence long and suberect, longest setae distinctly longer than combined length of antennomeres III–IV; punctation on elytra distinct and moderately fine; elytral apices obliquely truncate. Legs relatively long and slender, with dilated femora; profemora with sexual dimorphism and more dilated than mesofemora, the latter more dilated than metafemora.

♂: profemur distally broadly dilated externally, forming an angle of approximately 135°; sternites III and V with a median impression, that of sternite III circular and shallow, that of sternite V transverse and deeper; aedeagus (Fig. 64) 0.29 mm long and 0.14 mm broad; sclerotised portion of dorsal aspect relatively long; dorsal plate distinctly longer than broad, apically emarginate; ventral plate much broader than dorsal plate, apically broadly truncate; lateral contours straight in dorsal view, only apex broadly rounded; parameres each with two apical setae.

Comparative notes: *Stenichnus samius* is most similar to *S. brachati* MEYBOHM, 2019 (Crete). Both species share broadly oval, dorsally weakly flattened elytra, broadly dilated male profemora forming an obtuse angle distally, and a similar aedeagus. *Stenichnus samius* is easily distinguished from *S. brachati* by larger body size, a bicolored body, and longer and more erect pubescence

of pronotum and elytra. The new species differs from *S. chius* MEYBOHM, 2016 (Chios), with which it shares a bicolored body and long suberect pubescence of the elytra, by broader and dorsally more flattened elytra, and from *S. lesbius* MEYBOHM, 2016 by smaller body size and denser and shorter pubescence of the elytra.

Distribution and natural history: *Stenichnus samius* is most likely endemic to Samos. The specimens were collected in northwestern Samos, three of them in the valley of Potami river by sifting litter of *Platanus*. The altitudes range from 20 to 250 m.

Stenichnus (Stenichnus) amphimykalicus MEYBOHM spec. nov. urn:lsid:zoobank.org:act:BC1B8722-6DC0-40C0-BC10-9A2E673C9E4E (Figs 15, 65)

Type material: Holotype ♂: "Greece: Samos [16+1], Oros Kerkis: NW-slope, E Kalithea, 37°44'36"N 26°37'02"E, 580 m, 1.IV.2014, V. Assing / Stenichnus amphimykalicus m. Meybohm 2022 det. / Holotypus" (cMey). Paratypes 1 9: "N37°40'54 E027°10'03 (1) Türkei Aydin Dilek Dagi Kanyon 70-370 m 16.4.2006 l. Brachat & Meybohm" (cMey); 2 ♂ ♂: "N37°40'19 E027°10'41 (2) Türkei Aydin Dilek Dagi s Kanyon 670 m 17.4.2006 l. Brachat & Meybohm" (cMey); 1 9: "N37°40'56 E027°13'36 (25) Türkei Aydin Dilek Dagi s Güzelcamli 440 m 26.4.2006 l. Brachat & Meybohm" (cMey); 1 9: "N37°41'29 E027°09'33 (26) Türkei Aydin Dilek Dagi Kanyon 70-200 m 29.4.2006 l. Brachat & Meybohm" (cMey); 1 J: "N37°26'35 E028°18'07 (4) Türkei Mugla Yatagan-Bozdogan 825 m 19.4.2006 l. Brachat & Meybohm" (cMey); 1 J: "N37°55'45 E027°53'46 (9) Türkei Aydin Passayaylasi 1115 m 20.4.2006 l. Brachat & Meybohm" (cMey); 1 9: "TR [30] - Manisa, 10 km S Manisa, Karadağ, 1200 m, roadside, 38°33'26N 27°23'13E (26), 15.IV.2006 V. Assing" (cMey).

Etymology: The specific epithet (adjective) alludes to the fact that this species is distributed on both sides of the Strait of Mykali (the straight separating Samos from Turkey).

Description: Mean body length 1.29 mm. Male habitus as in Fig. 15. Body weakly bicolored; head and pronotum more or less distinctly dark-brown; elytra black; legs, antennae, and palpi pale-brown. Head strongly transverse, about 1.4 times as broad as long and 0.7 times as broad as pronotum; mean head width 0.26 mm, mean head length 0.19 mm; postocular region slightly longer than diameter of eyes; vertex and frons distinctly flattened, punctures and pubescence not dimorphic, punctures very fine and sparse, setae long; supra-antennal tubercles pronounced. Eyes composed of approximately 20 ommatidia, in male slightly larger and with slightly more ommatidia than in female. Antenna on average 0.87 mm long; antennomeres I–VI elongate, III–VI up to 1.5 times as long as broad, VII about as long as broad, VIII moderately transverse, IX and X distinctly transverse, X about twice as broad as long, XI distinctly shorter than the combined length of IX and X.

Pronotum weakly transverse, on average 0.33 mm long and 0.36 mm broad, pubescence long and suberect; basally only with two minute pits separated from each other by the same distance as from posterior margin.

Elytra broadly oval, dorsally weakly flattened, on average 0.81 mm long and 0.52 mm broad, broadest at anterior third, height approximately one-third of elytral length (lateral view); humeral angles and humeral folds absent; at base of each elytron with a large deep inner impression, lateral impressions absent; pubescence long and suberect, longest setae about as long as combined length of antennomeres III and IV; punctation on elytra distinct and moderately fine; elytral apices truncate. Legs relatively long and slender, with dilated femora; profemora with sexual dimorphism, more dilated than mesofemora, the latter more dilated than metafemora.

 σ : profemur distally broadly dilated externally, forming an angle of approximately 120°; sternite IV and posterior portion of sternite V with a shallow median, more or less transverse impression; aedeagus (Fig. 65) on average 0.23 mm long and 0.13 mm broad; sclerotised portion of dorsal aspect relatively short; dorsal plate as long as broad, apically more or less emarginate; ventral plate much broader than dorsal plate, apically weakly convex; aedeagus in dorsal view rounded basally, constricted at apical third, apex broadly convex; parameres each with two apical setae.

Intraspecific variation: The specimens of Turkey differ from the holotype by slightly larger body size, slightly less erect and slightly shorter and denser pubescence of the elytra, and an apically slightly less emarginate dorsal plate of aedeagus. These differences, however, are insignificant and consequently interpreted as intraspecific variation.

Comparative notes: Based on the similar structure of the aedeagus and the modifications of the male sternites III-V, S. amphimykalicus belongs to a species group including S. brachati and S. orientalis MEYBOHM, 2019 from Crete, as well as S. samius, S. chius, and S. lesbius from the Aegean Islands. Stenichnus amphimykalicus additionally shares distally broadly dilated male profemora with S. brachati, S. orientalis, and S. samius. Of all these species, S. amphimykalicus is most similar to S. brachati, with which it also shares broadly oval, dorsally weakly flattened elytra and the male profemora forming an obtuse angle distally. The new species is distinguished from all the species of this species group by shorter antennae with antennomeres IX and X approximately twice as broad as long and by smaller body size (exception: S. lesbius).

Distribution and natural history: *Stenichnus amphimykalicus* is known from one locality in Samos and several localities in Aydın, Muğla, and Manisa provinces, Southwest Turkey. The specimens were sifted from litter in various shrub and forest habitats at altitudes between 70 and 1200 m.

Scydmoraphes amphimykalicus MEYBOHM spec. nov. urn:lsid:zoobank.org:act:2CDA0C85-BA74-46EF-8AA6-ADAFB1974DC2 (Figs 16, 66)

Type material: Holotype ♂: "Greece: Samos [3], Oros Kerkis: Prof. Ilias, 37°43'34"N 26°38'02"E, 1210 m, sifted, 7.IV.2017, V. Assing / Scydmoraphes amphimykalicus m. Meybohm 2022 det. / Holotypus" (cMey). Paratypes: 1 3, $3 \circ \circ$: same data as holotype (cMey); $1 \circ$: "GR Samos Potami 20 m N37°47' E26°40' / leg. 22.4.2003 Brachat & Meybohm im Schluchtwald" (cMey); 1 ♂, 1 ♀: "Greece: Samos [16+1], Oros Kerkis: NW-slope, E Kalithea, 37°44'36"N, 26°37'02"E, 580 m, 1.IV.2014, V. Assing" (cMey); 2 ♂ ♂: N37°44'35"E 26°37'02" GR Samos (14a) Kerkis Oros 600 m leg. Brachat 5.4.2022 (cMey); 2 ♂ ♂: "GR Samos Kosmadei 560 m N37°45' E26°39' / 1.5.2003 Brachat & Meybohm aus Farn und Boden am Platanenstumpf" (cMey); 1 J, 1 9: "GR Samos Kosmadei 600 m N37°44' E26°38' / 30.4.2003 Brachat & Meybohm vor dem Felspfad zur Höhle Kakoperato" (cMey); 1 9: "N37°40'19 E027°10'41 (2) Türkei Aydin Dilek Dagi s Kanyon 670 m 17.4.2006 l. Brachat & Meybohm" (cMey); 2 ♂ ♂, 2 ♀ ♀: "N37°41'26 E027°13'43 (24) Türkei Aydin Dilek Dagi s Güzelkamli 410 m 28.4.2006 l. Brachat & Meybohm" (cMey); 2 ♂ ♂, 1 9: "N37°40'56 E027°13'36 (25) Türkei Aydin Dilek Dagi s Güzelkamli 440 m 28.4.2006 l. Brachat & Meybohm" (cMey).

Etymology: The specific epithet (adjective) alludes to the fact that this species is distributed on both sides of the Strait of Mykali.

Description: Habitus as in Fig. 16. External characters as in *S. samothracicus* MEYBOHM, 2019 from Samothraki, except as follows:

Body larger, length 1.05-1.12 mm. Pubescence of elytra shorter, setae in basal third more erect and in apical half more bent downwards. Head width 0.21 mm in male and 0.20 mm in female, head length 0.16 mm. Eyes sexually dimorphic, in female much smaller than in male; female eyes composed of approximately 10 ommatidia and diameter of eyes about as long as antennomere V, distance between eyes about four times as long as diameter of eyes; male eyes composed of approximately 15 ommatidia and diameter of eyes about 1.5 times as long as antennomere V, distance between eyes about 2.5 times as long as diameter of eyes. Antenna longer (0.52 mm) and stouter; antennomeres II almost 1.5 times as long as broad, III as long as broad, IV and V slightly longer than broad, VI as long as broad, VII to X of increasing width, VIII to X about twice as broad as long. Pronotum 0.31 mm long and 0.27 mm

 σ : aedeagus (Fig. 66) with slightly convex apico-lateral parts; dorso-apical plate of internal sac more slender, broadened to base, laterally without concavity; small horseshoe-shaped plate situated dorsally of frondshaped median sclerites indistinct; central projection of ventral border of ostium broader and laterally less emarginated.

Intraspecific variation: The specimens from Dilek Dağı differ from all except one of the specimens from Samos by slightly less curved setae on the elytra and by less distinctly broadened (dorsal view) and apically less convex (lateral view) elytra. The aedeagi, however, are identical.

Comparative notes: *Scydmoraphes amphimykalicus* belongs to a group including numerous species distributed in the northern Mediterranean (from Spain to Turkey) and in Georgia. This group is mainly characterised by a pair of frond-shaped sclerites in the internal sac of the aedeagus and by a pronotum with carinate lateral margins only in the posterior half. *Scydmoraphes amphimykalicus* is distinguished from all of them by the combination of characters indicated in the description above.

Distribution and natural history: This species is probably endemic to Samos and the geographically close Dilek Dağı in the Turkish province Aydın. The specimens were sifted from litter in various shrub and forest habitats at a wide range of altitudes (20 and 1210 m).

The Staphylinidae fauna of Ikaría

According to a checklist compiled in AssING (2017b), 70 species of Staphylinidae were previously known from Ikaría, seven of them endemic.

Hydrosmecta insularum Assıng, 2019

The specimens recorded as *Hydrosmecta* sp. in Assing (2017b) belong to *H. insularum* (see Assing et al. 2019).

Oxypoda rectacia Assing, 2019

According to a recent revision (Assing 2019a), the material listed in Assing (2017b) as *Oxypoda* cf. *nova* BERNHAUER, 1902 belongs to *O. rectacia*.

Description of new species

Cephennium icariae ASSING spec. nov. urn:lsid:zoobank.org:act:738FF668-DCED-4E48-808C-0DE0D67797CA (Figs 12, 62)

Type material: Holotype ♂: "GR Ikaria, Steli 290 m, N37°35' E26°09' / 27.4.2003, leg. Brachat & Meybohm, am grasigen Hang unter Gebüsch / Holotypus & Cephennium icariae sp. n. det. V. Assing 2022" (cAss). Paratypes: 1 J: "GR - Ikaria, Steli, 290 m, N37°35'42" E26°9'44", 27.IV.2003, leg. Meybohm/Brachat" (cAss); 5 ♂ ♂, 2 ♀ ♀: "GR Ikaria, Nas 10–100 m, N37°37' E26°03' / 26.4.2003, leg. Brachat & Meybohm, im Bachtal aus Gras unter Mastix-Strauch" (cAss); 1 9: "GR - Ikaria, westl. Arministis, Nas, 290 m, N37°37'03" E26°03'46", 10–100 m, 26.IV.2003, leg. Meybohm/Brachat" (cAss); 1 9: "GREECE: Ikaría [7], S Evdilos, pass env., 37°35'20"N, 26°10'22"E, 570 m, sifted, 10.IV.2017, V. Assing" (cAss).

Etymology: The specific epithet is the genitive of Icaria, the island where this species is most likely endemic.

Description: Body length approximately 1.0 mm. Habitus as in Fig. 12. Punctation of pronotum fine, only slightly more distinct than that of elytra. Eye rudiments minute, composed of 2–5 ommatidia without pigmentation.

♂: metaventrite flat, not impressed; aedeagus (Fig. 62) approximately 0.25 mm long, broad, and apically weakly convex; internal structure relatively large and symmetric, not claw-shaped; paramere apically extending slightly beyond apex of median lobe.

Comparative notes: Like the two species from Samos (*C. amplexans*, *C. monstrans*), *C. icariae* belongs to the *C. corruptum* group. It is reliably distinguished from other species of this group only by the shapes of the internal structures of the aedeagus. For illustrations of the aedeagi of the Turkish representatives of the *C. unguicum* subgroup see ASSING & MEYBOHM (2021a).

Distribution and natural history: The type material was found in three localities in Central Ikaría at altitudes between 10 and 570 m. The specimens were sifted from grass, litter, and roots in a grassy slope, in a stream valley, and at the margin of a pasture.

The Staphylinidae fauna of Samothraki

According to Assing (2019b), the fauna of Samothraki was composed of at least 123 species, 114 of them named and six endemic. In the meantime, a revision of some of the material has revealed some misidentifications, one of them already published (Assing 2020). The material previously identified as *Atheta laevi*- *gata* now belongs to the type series of *A. tecta* (see the description of this species above). Other identification errors are rectified below. Including the additions, the fauna of Samothraki now includes at least 125 species, 116 of them named.

Aloconota eichhoffi (Scriba, 1868)

Two of the specimens reported as *Aloconota cambrica* (WOLLASTON, 1855) from localities 9 and 9a in ASSING (2019b) refer to *A. eichhoffi*.

Aloconota planifrons (WATERHOUSE, 1863)

One of the specimens reported as *Aloconota cambrica* from locality 12a in AssING (2019b) belongs to *A. planifrons*.

Cypha spathulata Assing, 2007

The record of *C. graeca* Assing, 2004 in Assing (2019b) in fact refers to *C. spathulata* (see Assing 2020).

The Staphylinidae fauna of Lesbos

According to a recent study of the fauna of Lesbos (ASSING 2016b), 199 species, 184 named and 15 unnamed, had been recorded from this island. Nine of the named and at least two of the unnamed Pselaphinae and Scydmaeninae were considered endemics.

Hydrosmecta insularum Assıng, 2019

The specimens recorded as *Hydrosmecta* sp. 2 in Assing (2016b) belong to *H. insularum* (see Assing et al. 2019).

Myllaena graeca Kraatz, 1858

A revision of some Mediterranean and Caucasian species of *Myllaena* ERICHSON, 1837 (ASSING 2018b) revealed that the specimen listed as *Myllaena* sp. in ASSING (2005) belongs to *Myllaena graeca*.

Pella discolor Assing, 2008

The record of *Pella cinctipennis* (EPPELSHEIM, 1884) from Lesbos in ASSING (2016b) is based on misidentification. A revision of the material revealed that it belongs to *P. discolor*, the first record of this species from Greece and Europe. **Tab. 3**: Updated checklist of the Staphylinidae recorded from Rhodos and results of the field trip conducted to Rhodos by Thomas Forcke in December 2019. Details on the localities/samples are provided at the end of the list. Specimen numbers are given in parentheses behind the sample number. First records from Rhodos are marked with an asterisk, first records from Greece with two, and first records from Europe with three asterisks. For additional footnotes and references for species recorded from Rhodos before 2013 see the footnotes and references in Assing (2013).

The references are abbreviated as follows: A13 = Assing (2001); A17a = Assing (2017a); A18a = Assing (2018a); A19a = Assing (2019a); A19c = Assing (2019c); A21 = Assing (2021); App = Assing (present paper); M14 = MAKRANCZY (2014).

Species	Localities/samples	References
O m a l i i n a e		
Aphaenostemmus rhodicus Assıng, 2006		A13
Dialycera aspera (Eppelsheim,1889)		A13
Omalium rhodicum ZANETTI & Assing, 2013		A13
Omalium rivulare (PAYKULL, 1789)		A13
Omalium rugatum MULSANT & REY, 1880		A13
Proteininae		
Metopsia assingi ZERCHE, 1998		A13
Proteinus atomarius Erichson, 1840		A13
Proteinus utrarius Assıng, 2004		A13
Micropeplinae		
Micropeplus fulvus Erichson, 1840		A13
Micropeplus staphylinoides (MARSHAM, 1802)		A13
P s e l a p h i n a e		
Brachygluta cavernosa (SAULCY, 1876)		A13
+Bryaxis nov. sp.		A13
Faronus distinctus Besucher, 1999		A13
Namunia myrmecophila Reitter, 1884		A13, A17a
Reichenbachia chevrieri (AUBÉ, 1844)		A13
*Rybaxis longicornis (LEACH, 1817)	1(2), 1a(7), 1b(4), 1c(4)	Арр
***Sognorus calcaratus (BAUDI DI SELVE, 1870)	5(1)	Арр
Tribatus creticus Reitter, 1884		A13
Trimium libani J. SAHLBERG, 1908		A13
Tychus rhodensis SABELLA et al., 1998		A13
Tachyporinae		
Lordithon thoracicus (FABRICIUS, 1777)		A13
<i>Mycetoporus ignidorsum</i> Еррегsнеім, 1880	16(1)	A13, App
Mycetoporus reichei (PANDELLÉ, 1869)		A13
Mycetoporus cf. simillimus FAGEL, 1965		A13
Parabolitobius inclinans (GRAVENHORST, 1806)		A13
Sepedophilus immaculatus (STEPHENS, 1832)	13(1), 13a(1)	A13, App
*Sepedophilus obtusus (Luze, 1902)	10b(1)	Арр
Tachyporus abner SAULCY, 1865	la(1)	A13, App
Tachyporus caucasicus Kolenati, 1846	10b(1)	A13, App
Tachyporus hypnorum (FABRICIUS, 1775)		A13
Tachyporus nitidulus (FABRICIUS, 1781)	1b(1), 2(1), 16(1)	A13, App

Species	Localities/samples	References
Habrocerinae		
Habrocerus cyprensis Assing & Wunderle, 1995		A13
Habrocerus pisidicus Korge, 1971		A13
Aleocharinae		
Aleochara bipustulata (LINNAEUS, 1760)		A13
Aleochara laticornis KRAATZ, 1856		A13
Aleochara maculipennis BAUDI DI SELVE, 1857		A13
Aleochara tristis (GRAVENHORST, 1802)		A13
Alevonota libanotica (FAGEL, 1965)		A13
*Aloconota aegaea Assing, 2016		Арр
**Aloconota pfefferi (ROUBAL, 1929)		Арр
Aloconota sulcifrons (STEPHENS, 1832)		A13
Atheta aegra (HEER, 1841)		A13
Atheta aeneicollis (SHARP, 1869)	13(1)	A13, App
*Atheta amicula (STEPHENS, 1832)	1b(1)	Арр
Atheta clientula (Erichson, 1839)		A13
Atheta coriaria (KRAATZ, 1856)		A13
Atheta longicornis (GRAVENHORST, 1802)		A13
Atheta meybohmi Assıng, 2011		A13
Atheta rhodiensis SCHEERPELTZ, 1963		A13
Atheta trinotata (KRAATZ, 1856)		A13
Atheta (Mocyta) spp.	1(12), 1a(1), 1b(17), 1c(1), 2(2), 6(1), 7(2)	A13, App
Cousya araxis (Bernhauer, 1902)		[A13] ¹⁾ , A18a
Cousya defecta (MULANT & REY, 1875)		[A13] ¹⁾ , A18a
Dinusa sp. (º)		A13
Geostiba lucens (BENICK, 1970)		A13
Geostiba oertzeni (Eppelsheim, 1888)		A13
Geostiba rhodiensis PACE, 1983		A13
Liogluta longiuscula (GRAVENHORST, 1802)		A13
<i>Myrmecopora fugax</i> (Erichson, 1839)		A13
<i>Myrmecopora laesa</i> (Erichson, 1839)		A13
+ <i>Myrmecopora rhodica</i> Assıng, 2013		A13
Nehemitropia lividipennis (MANNERHEIM, 1830)	1(1), 1b(1)	A13, App
Oligota pusillima (Gravenhorst, 1806)		A13
Oxypoda bimaculata BAUDI DI SELVE, 1870	1b(1), 2(1), 7(3)	A13j, App
Oxypoda exoleta Erichson, 1839		A13
Oxypoda lesbia Assing, 2005		A13
Oxypoda obscuricollis Assıng, 2007		A13
Oxypoda rectacia Assıng, 2019		[A13] ¹⁾ , A19a
<i>Oxypoda</i> sp.	1b(1)	App
Parocyusa longitarsis (ERICHSON, 1839)		A21
***Pseudosemiris atavia Assıng, 2001	7(1)	App

Species	Localities/samples	References
O x y t e l i n a e		
Anotylus clypeonitens (PANDELLÉ, 1867)		A13
Anotylus complanatus (ERICHSON, 1839) (♀)		A13
Anotylus inustus (GRAVENHORST, 1806)	2(1), 6(1)	A13, App
Anotylus sculpturatus (GRAVENHORST, 1806)		A13
Anotylus tetracarinatus (BLOCK, 1799)		A13
*Bledius bedelianus SCHÜLKE, spec. nov.		Арр
Bledius bicornis (GERMAR, 1823)		A13
Bledius fossor HEER, 1839		A17a
Bledius spectabilis (KRAATZ, 1857)		A13
Bledius unicornis (GERMAR, 1825)		A13
*Bledius verres Erichson, 1840	11(6)	Арр
***Carpelimus prudeki GILDENKOV, 2013	1c(1)	Арр
Ochthephilus andalusiacus (FAGEL, 1957)		M14
Ochthephilus venustulus (Rosenhauer, 1856)		M14
*Platystethus brevipennis BAUDI DI SELVE, 1857	2(9), 6(3), 7(1)	Арр
Platystethus nitens (SAHLBERG, 1832)		A13
*Platystethus spinosus Erichson, 1840	7(10)	Арр
Steninae		
Stenus aceris Stephens, 1833	1(2), 1a(4), 1b(1), 1c(11)	A13, App
Stenus assequens assequens REY, 1884		A13
Stenus brunnipes lepidus WEISE, 1875	6(1)	A13, App
Stenus glacialis HEER, 1839		A13
Stenus guttula Müller, 1821	11(1), 16(2)	A13, App
Stenus hospes Erichson, 1840	2(1), 10b(1)	A13, App
Stenus impressus GERMAR, 1824		A13
Stenus maculiger WEISE, 1875		A13
Stenus subaeneus Erichson, 1840		A13
Stenus turbulentus BONDROIT, 1912	16(5)	A13, App
S c y d m a e n i n a e		
+ <i>Cephennium rhodicum</i> Assıng, nov. spec.		A13 ¹⁾ , App
+Euconnus dodecanicus Franz, 1966		A13
+Euconnus oblitus FRANZ, 1972		A13
+Euconnus rhodensis FRANZ, 1966		A13
+ <i>Euconnus rhodicus</i> Меувонм, nov. spec.		A13
Scydmaenus menozzii FRANZ, 1966		A13
+Scydmoraphes rhodensis (FRANZ, 1966)		A13
Leptotyphlinae		
+Kenotyphlus rhodiensis COIFFAIT, 1973		A13
P a e d e r i n a e		
Achenium debile Erichson, 1840	1b(1), 2(3), 4(4), 6(5), 7(5)	A13, App
Achenium picinum Fauvel, 1875		A13
*Astenus bimaculatus bimaculatus (ERICHSON, 1840)	7(2)	Арр

Species	Localities/samples	References
*Astenus melanurus (Küster, 1853)	7(1)	Арр
Astenus procerus (GRAVENHORST, 1806)		A13
Astenus thoracicus (BAUDI DI SELVE, 1857)	5(9), 7(6), 17(1)	A13, App
Astenus rhodicus Assıng, 2013.	10(1)	A13, App
Domene stilicina (ERICHSON, 1840)	1(1), 2(5), 5(1), 6(5), 7(21), 12(1)	A13, App
Leptobium gracile (GRAVENHORST, 1802)	4(7)	A13, App
Lobrathium rugipenne (Носннитн, 1851)	1(2), 10(1), 10b(9), 11(1), 14(2), 16(2), 19(1)	А13, Арр
Medon dilutus pythonissa (SAULCY, 1865)	1a(1), 15(1), 16(2)	A13, App
Medon impar Assıng, 2004		A13
Medon lydicus Bordoni, 1980		A13
Medon maronitus (SAULCY, 1865)		A13
Medon semiobscurus (FAUVEL, 1875)		A13
Micranops pilicornis (BAUDI DI SELVE, 1870)		A13
Micrillus testaceus (Erichson, 1840)	1b(1)	A13, App
*Ochthephilum turkestanicum (Korge, 1968)	1(3), 1a(1), 1b(34), 1c(13), 6(3)	Арр
Pseudolathra quadricollis (FAUVEL, 1875)	1b(3), 1c(1)	A19c, App
+Scopaeus schusteri Scheerpeltz, 1965		A13
*Scymbalium anale (Nordmann, 1837)	4(5)	Арр
+Sunius rhodicus Assıng, 2013		A13
Staphylininae		
Bisnius fimetarius (GRAVENHORST, 1802)		A13
Bisnius sordidus (GRAVENHORST, 1802)		A13
Dinothenarus flavocephalus (GOEZE, 1777)		A13
*Gabrius latro Joy, 1913	1b(2)	Арр
*Gabronthus maritimus (MOTSCHULSKY, 1858)	1a(2), 1b(5), 1c(1)	Арр
*Gauropterus fulgidus (FABRICIUS, 1787)	3(9), 7(1)	Арр
*Heterothops dissimilis (GRAVENHORST, 1802)	1a(1), 1c(1)	Арр
Megalinus glabratus (GRAVENHORST, 1802)		A13
Megalinus scutellaris (FAUVEL, 1900)		A13
Ocypus curtipennis Motschulsky, 1849	5(1), 10(3), 10a(1), 13a(1), 20(16)	A13, App
Ocypus mus (Brullé, 1832)	1(1), 10(7), 10a(1), 13(3), 13a(12), 15(14), 18(2), 20(1)	А13, Арр
Ocypus orientis Smetana & Davies, 2000	3(1), 4(1), 5(3), 6(2)	A13, App
Othius laeviusculus STEPHENS, 1833		A13
Othius lapidicola Märkel & Kiesenwetter, 1848	(15(3), 16(1)	A13, App
Phacophallus parumpunctatus (Gyllenhal, 1827)		A13
Philonthus concinnus (GRAVENHORST, 1802)	1(1), 1b(1), 1c(2)	A13, App
Philonthus cruentatus (GMELIN, 1790)		A13
Philonthus ebeninus (GRAVENHORST, 1802)		A13
*Philonthus juvenilis Peyron, 1858	14(1)	Арр
*Philonthus rufimanus rufimanus Erichson, 1840	11(2)	Арр

Species	Localities/samples	References
Quedius acuminatus phenicius COIFFAIT, 1967		A13
Quedius cinctus (PAYKULL, 1790)		A13
Quedius fissus Gridelli, 1938	15(5)	А13, Арр
Quedius humeralis STEPHENS, 1832		A13
Quedius lateralis GRAVENHORST, 1802		A13
Quedius levicollis (Brullé, 1832)	15(46)	A13, App
Quedius nemoralis BAUDI DI SELVE, 1848	15(30), 16(6)	А13, Арр
*Quedius nitipennis (STEPHENS, 1833)	10(1), 10b(5),	Арр
*Quedius ortrudae Korge, 1971	1a(1)	Арр
Quedius scintillans (GRAVENHORST, 1806)	7(1), 10b(1), 16(1)	A13, App
Quedius semiobscurus (MARSHAM, 1802)	15(60)	A13, App
*Quedius suturalis Kiesenwetter, 1845	16(1)	Арр
**Quedius vicinus (Ménétriés, 1832)	1(1), 1a(2), 1b(1), 4(1), 6(4)	Арр
*Stenistoderus cephalotes (KRAATZ, 1858)	1b(3), 4(5)	Арр
Xantholinus rufipennis Erichson, 1839	1c(1), 6(1), 7(3)	A13, App
Xantholinus varnensis COIFFAIT, 1972	1(2), 1a(1), 1b(1), 1c(1), 2(3), 6(1), 7(17)	A13, App

Footnote: ¹⁾ listed as unnamed species in Assing (2013).

Cafius pruinosus (ERICHSON, 1840)

ASSING (2016b) recorded *Remus sericeus* HOLME, 1837 from Lesbos. A revision of the material revealed that it in fact belongs to *Cafius pruinosus*, a species recently moved from *Remus* HOLME, 1837 to *Cafius* STEPHENS, 1829 by Yoo et al. (2021).

The Staphylinidae fauna of Rhodos

According to a first comprehensive account, including an annotated checklist, the Staphylinidae fauna of Rhodos was composed of at least 130 species, with thirteen (three undescribed) of them endemic. In the meantime, several additional species have been reported (AssING 2017a, 2019c, 2021, MAKRANCZY 2014), some previously unnamed species have been revised (Assing 2018a, 2019a), and two species previously assumed to be endemic (*Omalium rhodicum, Astenus rhodicus*) have been shown to be more widespread.

Thomas Forcke (Hannover) conducted a field trip to Rhodos from 22 December 2019 to 1 January 2020, primarily aiming at collecting Carabidae. The staphylinid by-catches totalled 341 specimens belonging to 42 species, with as many as 18 of them recorded from Rhodos and five from Greece for the first time.

David Wrase (Gusow-Platkow), too, collected Carabidae in Rhodos in May, 2015, and forwarded the Staphylinidae (276 specimens of 21 species, four of them unidentified, six of them first records from Rhodos) to Michael Schülke. Including all the changes and additions, the staphylinid fauna of Rhodos is now composed of 160 identified named species (plus several unidentified and unnamed species), ten of which are currently known only from Rhodos and probably endemic.

In view of the number of additions and changes, an updated checklist is provided in Tab. 3.

Localities/samples: 22 December 2019-1 January 2020, leg. Forcke (sample numbers 1-9): 1: 2 km S Haraki, 36°08'58"N, 28°04'47"E, 4 m, beach, under debris and stones, 22.XII.2019; 1a: same data, but under debris, 27.XII.2019; 1b: same data, but debris sifted, 28.XII.2019; 1c: same data, but debris sifted, 1.I.2020; 2:3 km SW Apolakkia, mouth of Kourkourtahi, 36°03'03"N, 27°45'35"E, 3 m, arable land near beach, under debris and stones, 24.XII.2019; 3: northern part of Apolakkia dam, 36°06'47"N, 27°47'37"E, 105 m, wetland, under debris and stones, 25.XII.2019; 4: NW Siana, Akramitis north side, 36°09'58"N, 27°45'43"E, 290 m, partly flooded arable land, under stones and debris, 25.XII.2019; 5: NW Siana, Akramitis north side, 36°09'49"N, 27°44'27"E, 270 m, grassy area, under stones, 26.XII.2019; 6: 3 km N Apolakkia, 36°05'37"N, 27°47'53"E, 80 m, arable land, under stones, 29.XII.2019; 7: S Kissamos, Limni, 35°22'15"N, 23°37'55"E, 570 m, margin of artificial pond with old Platanus orientalis, litter sifted, 22.XII.2019; 8: NE Kandanos, Spina, 35°21'43"N, 23°46'51"E, 710 m, stream valley with oak, Platanus orientalis, other trees, and fern undergrowth, soil washing, 22.XII.2019; 9: 1 km N Apolakkia, 36°04'25"N, 27°47'24"E, 50 m, partly flooded arable land, under stones and debris, 31.XII.2019.

14-20 May, 2020, leg. Wrase & Laser (sample numbers 10-20): 10: ca 1 km SE Ialysos, 36°24'09"N, 28°09'19"E, 30 m, grassland with olive trees, under stones, 14-19.V.2015; 10a: same data, but pitfall traps with vinegar, 19-20.V.2015; 10b: ca 1 km SE Ialysos, 36°24'09"N, 28°09'19"E, 20 m, stream, under vegetation and stones, 20.V.2015; 11: W Kolymbia, Loutanis river at Epta Piges (Seven Springs), 36°15'34"N, 28°06'34"E, 80 m, river bank, in gravel, 15&19.V.2015; 12: ca 3.5 km SW Archangelos, 36°12'00"N, 28°03'59", 20 m, loamy field margin, under vegetation, 15.V.2015; 13: W Kolymbia, forest near Epta Piges, 36.04'08"N, 28°04'30"E, 90 m, nearly dry streambed, under stones, 16.V.2015; 13a: same data, but pitfall traps with vinegar; 14: NW Kiotari, Asklipinos river, near bridge, 36°04'01"N, 27°55'44"E, 60 m, river bank with gravel and mud, 16.V.2015; 15: Profitis Ilias, 36°16'25"N, 27°55'44", 690 m, pine forest, under stones and sifted from litter and moss, 17.V.2015; 16: ca 1.2 km SE Eleoússa, 36°16'14"N, 28°02'36"E, 200 m, bank of small stream with gravel, 17.V.2015; 17: 2.8 km W Ghadura Reservoir, 2.8 km SW Masari, 36°09'45'N, 28°03'E, 30 m, loamy wetland, 19.V.2015; 18: Petaloudes (Butterfly Valley), 36°12'16"N, 28°03'45"E, 230 m, forest, under stones along path, 20.V.2015; 19: W Ancient Kamiros, 36°20'09'N, 27°56'27"E, 5 m, gravel bank of small stream, 20.V.2015; 20: 3 km S Ródos, Princess Flora Hotel, 36°24'34'N, 28°13'29"E, 20 m, around buildings, collected at night, 14-20.V.2015.

Notes on some species

Sognorus calcaratus (BAUDI DI SELVE, 1870)

According to SCHÜLKE & SMETANA (2015), this species had been recorded only from Turkey and Cyprus. The specimen listed in Tab. 3 represents the first records from Rhodos, Greece, and Europe.

Pseudosemiris atavia Assıng, 2001

Only three specimens from the Turkish provinces Mersin and Osmaniye were known previously (Assing 2001, 2004). The male listed in Tab. 3 represents the first records from Rhodos, Greece, and Europe.

Carpelimus prudeki GILDENKOV, 2013

The original description is based on type material from the environs of Silifke in the Turkish province Mersin (GILDENKOV 2013). The male listed in Tab. 3, which was identified by Mikhail Gildenkov based on a photograph of the adeagus, represents the first records from Rhodos, Greece, and Europe.

Quedius vicinus (Ménétriés, 1832)

The previously known distribution included Turkey, the Middle East, the Caucasus region, and Middle Asia (SCHÜLKE & SMETANA 2015). The material from Rhodos (and Samos; see the section on the Staphylinidae of Samos) represents the first records from the Aegean Islands (Rhodos, Samos) and Greece.

Additional records

Aloconota aegea Assing, 2016

Material examined: Rhodos: 1 ex., 5 km W Kolimbia, 14.V.2013, leg. Ziegler (cAss).

This species was previously known only from Samos, Lesbos, and Samothraki. The above specimen represents the first record from Rhodos.

Aloconota pfefferi (ROUBAL, 1929)

Material examined: Rhodos: 1 ex., 5 km W Kolimbia, 14.V.2013, leg. Ziegler (cAss).

This species was previously known from Central Europe, Italy, the northern Balkans, and Ukraine (SCHÜLKE & SMETANA 2015). The above specimen represents the first record from Rhodos and Greece.

Parocyusa longitarsis (ERICHSON, 1839)

Material examined: Rhodos: 1 ex., Loutani, above Epta Piges, 7–19.IV.1995, leg. Zwick (cAss); 1 ex., 5 km W Kolimbia, 14.V.2013, leg. Ziegler (cAss); 1 ex., N Apolakkia, reservoir, 36°05'49"N, 27°47'29"E, 90 m, 9.IV.2007, leg. Bahr et al. (MNB).

The first two specimens listed above were already reported from Rhodos without specified localities by ASSING (2021).

Descriptions of new species

Cephennium rhodicum ASSING spec. nov. urn:lsid:zoobank.org:act:63142283-CD20-46E9-AB60-9F7AFA6DE29A (Figs 13, 63)

Type material: Holotype σ : "GR Rhodos 300 m, Eleousa Ag. Nikolaos, Fountoukli, Meybohm 10.4.1999 / Holotypus σ *Cephennium rhodicum* sp. n. det. V. Assing 2022" (cAss). Paratypes: 1σ , $6 \Leftrightarrow \phi$: same data as holotype; 1σ : same data as holotype, but 9.4.1999 (cAss); $1 \Leftrightarrow$: "GR Rhodos (6), Epta Piges 70 m, N36°15'22 E28°06'50,

leg. Meybohm 3.4.2012" (cAss); 1σ : "GR Rhodos 320 m (2), Ag. Nikolaos Fontoukli, N36°16'27 E27°59'51, leg. Meybohm 3.4.2012" (cAss); 1σ : "GR Rhodos 500 m, Salakos Prof. Ilias, östl. Hotels, Meybohm 9.4.1999" (cAss); 1φ : "GREECE – Rhodos [4], 5 km SW Embonas, Attaviros, 36°11'59"N, 27°48'56"E, 580 m, 17.III.2013, V. Assing" (cAss).

Etymology: The specific epithet (Latin, adjective) alludes to the distribution of this species, which is most likely confined to Rhodos.

Description: Body length approximately 1.0 mm. Habitus as in Fig. 13. Punctation of pronotum significantly denser and more distinct than that of elytra. Eye rudiments minute, composed of 3–5 ommatidia without pigmentation.

 σ : aedeagus (Fig. 63) approximately 0.25 mm long, broad, and apically weakly, obtusely pointed in the middle; internal structure relatively large and symmetric, not claw-shaped; paramere apically extending to apex of median lobe.

Comparative notes: Based on the structure of the aedeagus and external characters, *C. rhodicum* belongs to the *C. corruptum* group, whose representatives are distributed in the region from Samos and Ikaría across South Turkey to the Middle East (Lebanon, Jordan) (ASSING & MEYBOHM 2021a) and from which *C. rhodicum* is reliably distinguished only by the shapes of the internal structures of the aedeagus. For illustrations of the aedeagi of the previously described representatives of the *C. corruptum* group see ASSING & MEYBOHM (2021a).

Distribution and natural history: The known distribution is confined to several localities in Rhodos. The specimens were found at altitudes of 70–580 m. The specimen from Attaviros was sifted from litter in a pine forest.

Euconnus (Tetramelus) rhodicus MEYBOHM spec. nov. urn:lsid:zoobank.org:act:A286F3E4-9AC8-434F-B473-96E388B238A8 (Figs 17, 67)

Type material: Holotype σ : "Greece: Rhodos [12], N Monolithos, Akramitis, 36°09'56"N, 27°45'43"E, 300 m, pine forest, sifted 21.III.2013, V. Assing / *Euc.* (*Tetramelus*) *rhodicus* m. Meybohm 2022 det. / Holotypus" (cMey). Paratype: 1 \circ : same data as holotype (cMey).

Etymology: The specific epithet (Latin, adjective) alludes to the hypothesis that this species is probably endemic to Rhodos.

Description: Habitus as in Fig. 17. Body length 1.77 mm (σ) and 1.85 mm (φ). Head approximately 0.28 mm broad, 1.18 times as long as broad, postero-medially bulging and projecting beyond posterior constriction.

Eyes with three ommatidia, weakly pigmented in male and without pigmentation in female. Antenna 0.80 mm (\circ) and 0.85 mm (\circ) long; antennomeres II twice as long as broad, III-VI cylindrical and about 1.5 times as long as broad, VII intermediate between VI and VIII, VIII-X nearly as broad as long, and XI distinctly shorter than the combined length of IX and X. Pronotum nearly 1.15 times as long as broad. Elytra 1.6 times (\eth) and 1.5 times (\Im) as long as broad. Legs slender, mesotibiae 0.42 mm long. J: metasternum with long, broad, and deep median impression; mesotrochanter unmodified; aedeagus (Fig. 67) slender, 0.54 mm long; ventral process (dorsal view) nearly straight and only indistinctly tapering from base, on either side of middle without a series of microsetae; the two large sclerites slender, almost symmetrically and evenly curved and converging, without teeth; dorsal plate apically broadly truncate, and with large emargination, lateral wing-shaped structures large, distinctly projecting from lateral contours of capsule; parameres with three apical setae.

Comparative notes: *Euconnus rhodicus* is distinguished from the similar *E. kerkisicus* from Samos by slightly larger body size, slightly longer and especially more slender antennae with antennomeres VIII–X not broader than long, longer tibiae, a large impression on the male metasternum, and unmodified male mesotrochanters. The aedeagus differs from that of *E. kerkisicus* especially by its more slender shape, a large emargination of the dorsal plate, and large, slender, and almost symmetric internal sclerites.

Distribution and natural history: This species is most likely endemic to Rhodos. The specimens were sifted from litter in a pine forest with undergrowth at an altitude of 300 m.

On two Mediterranean *Bledius* species (authored by Michael Schülke)

Bledius (Hesperophilus) bedeli FAUVEL, 1878 (Fig. 8)

Type material examined: Syntypes: $1 \circleft : "Daya [round, violet] / Fauvel [round, red] / Bedeli Daya (Algerie) / P. de Borre / SYNTYPUS- <math>\circleft : Badeli Eauvel, 1878$ det. M. Schülke 2011" (MHNG); $1 \circleft : "Daya / Spec. typ. / J. Sahlb. / 9. / Bedeli Fvl. Alg. typ. / D.E.I. Coll. von Heyden / DEI Müncheberg Col-02277 [light green]" (SDEI); <math>1 \circleft : "Bedeli Fauv. Daya, Oran Bedel typ. / D.E.I. coll. von Heyden / Bledius Bedeli Fauv. / DEI Müncheberg Col-02276 [light green]" (SDEI); <math>1 \circleft : "Bedeli Fauv. Daya, Oran Bedel typ. / D.E.I. coll. von Heyden / Bledius Bedeli Fauv. / DEI Müncheberg Col-02276 [light green]" (SDEI); <math>1 \circleft : "Daya / Bedeli Fauv. / Coll. R. I. Sc. N. B." (IRSNB); <math>1 \circleft : "Daya (Algerie) / Bedeli / Fauvel Type / Coll. R. I. Sc. N. B." (IRSNB); 4 exs. [without locality data, but mounted on identical mounting cards]: "Coll. R. I. Sc. N. B." (IRSNB).$



Figs 1–13: Habitus of Paramaurops creticus (1), Afropselaphus chanianus (2), Euphanias ambulans (3), Liogluta aloconotoides, holotype (4), Hydrosmecta cultellata (5), Atheta tecta (6), Cretotyphlus kerkisicus (7), Bledius bedeli (8), B. bedelianus (9), Cephennium amplexans (10), C. monstrans (11), C. icariae (12), and C. rhodicum (13). Scale bars: 1.0 mm.



Figs 14–26: Stenichnus samius (14), S. amphimykalicus (15), Scydmoraphes amphimykalicus (16), Euconnus rhodicus (17), Euphanias ambulans (18–20), Paramaurops creticus (21), Afropselaphus chanianus (22), Cephennium idanum from Selena Oros (23), Liogluta aloconotoides, holotype (24), and Hydrosmecta cultellata (25–26): 14–17 – habitus; 18 – antenna; 19 – forebody in dorsal view; 20 – forebody in lateral view; 21–23 – aedeagus; 24–26 – spermatheca. Scale bars: 14–17: 1.0 mm; 18–20: 0.5 mm; 21–26: 0.1 mm.



Figs 27–39: *Hydrosmecta cultellata* (27–32), *Atheta laevigata*, Armenia (33), *A. cretica* (34), *A. monstruosa*, Armenia (35), *A. balcanicola*, Pelopónnisos (36), and *A. tecta* (37–39; 37: holotype; 38: Samos; 39: Kos): 27 – male tergite VIII; 28 – male sternite VIII; 29–30, 33–39 – median lobe of aedeagus; 31 – female tergite VIII; 32 – female sternite VIII. Scale bars: 27–28, 31–39: 0.2 mm; 29–30: 0.1 mm.



Figs 40–57: *Throbalium cycladicum* (**40–41**, **50–52**: mainland Greece; **42**, **48–49**, **53**: Samos; **43**: Kos; **44**, **54**: Cyprus; **45**, **56**: Turkey: Adana; **46**: holotype of *T. adanense*; **47**, **57**: Israel; **55**: Turkey: Kayseri): 40–47 – aedeagus in ventral view; **48** – male sternite VII; **49** – male sternite VIII; 50–57 – postero-median portion of male sternite VII. Scale bars: 0.2 mm.



Figs 58–67: Cretotyphlus kerkisicus (58–59), Cephennium amplexans (60), C. monstrans (61), C. icariae (62), C. rhodicum (63), Stenichnus samius (64), S. amphimykalicus (65), Scydmoraphes amphimykalicus (66), and Euconnus rhodicus (67): 58, 60–67 – aedeagus; 59 – male sternite VIII. Scale bars: 0.1 mm.



Figs 68–71: Bledius bedelianus: 68–69 – aedeagus in ventral view (68: Israel; 69: Turkey: Antalya); 70–71 – male sternite VII (70: Antalya; 71: Israel). Scale bars: 70–71: 0.5 mm; 68–69: 0.2 mm.

Additional material examined: ALGERIA: 1 ex., "Hussein-Dey, plage prés polygone" (IRSNB); 2 exs. [without labels but mounted on identical mounting cards] (IRSNB); 2 exs., Tlemcen env., 1932, leg. Lebrousse (IRSNB).

Distribution: *Bledius bedeli* was originally described from Algeria (FAUVEL 1878) and subsequently recorded from Morocco by KOCHER (1958), as well as from Lebanon and Turkey by HERMAN (1986). A revision of the material recorded by HERMAN (1986) revealed that the records from Lebanon and Turkey are based on misidentification and refer to an undescribed species (see material of *B. bedelianus* below). The known distribution of the species is restricted to North Africa. The habitus of a syntype from Daya is shown in Fig. 8.

Bledius (Hesperophilus) bedelianus SCHÜLKE spec. nov. urn:lsid:zoobank.org:act:84848C14-D6BD-41DF-9D76-EFDDC83FE7CD (Figs 9, 68–71)

Type material: Holotype ♂: "GREECE, Rhodos Lindos Saidouras river 18.04.1994 leg. J. Frisch / Holotypus *Bledius* (*Hesperophilus*) bedelianus spec. nov. det. M. Schülke 2011 [red]" (MNB). Paratypes: 26 exs.: same data as the holotype (MNB); 1 ex.: "GR Rhodos, Kolimbia, Strand, 36°15'17"N 28°10'06"E, 06.04.2007, leg. Bahr, Bayer, Brunner & Winkelmann (FO15)" (MNB); 5 exs.: "GREECE, Karpathos Aperi, 300 m NN 25.IV.1994, leg. J. Frisch" (MNB); 1 ex.: "GREECE, Karpathos Pigadia 10 m NN 22.IV.1994, leg. J. Frisch" (MNB); 1 ex.: "GREECE (Dodecanissa) Ródos, 2.8 km W Ghadura Reservoir, 2.8 km SW Masari, N36°09'44.6"/E 28°03' (loamy wet area) 32 m, 19.V.2015, Wrase & Laser [12]" (MNB); 3 exs.: "GREECE: Samos [23a] E Karlovassi, 37°47'14"N, 26°43'46"E, 40 m, river bank, 14.IV.2022, T. Forke" (cAss); 2 exs.: same data, but "[23b], 16.IV.2022, leg. Assing & Forcke" (cAss); 9 exs.: "GREECE: Samos [49], E Myli, 37°40'39"N, 26°52'19"E, 15 m, river bank, 12.IV.2022, T. Forke" (cAss); 1 ex.: "GREECE: Samos [60], W Karlovassi, Potami, 37°47'11"N, 26°40'05"E, 30 m, stream bank, 16.IV.2022, V. Assing" (cAss); 1 d: "ISRAEL, (Centr. distr.) N Tel Aviv, Breikhat Ya'ar wetland 28.III.2008, leg. T. Aßmann" (MNB); 2 exs.: "Liban: Nahr el Kelb V. 1964 G. Fagel / G. Fagel sp. apud. bedeli Fauv. / [green round label] / Coll. R. I. Sc. N. B." (IRSNB); 1 ex.: "Beirut / U. Sahlb. / 2322 / Bledius bedeli Fvl." (IRSNB); 1 ex.: "Turkey, Anatolia mer. Mersin; Kurtulus nr. Silifke, 12.-17.V.1994, leg. Prudek" (MNB); 16 exs.: "TURKEY: nr. Antalya Ihsaniye, 07.v.2001, 200 m 36°55'08"N 30°49'42"E ex flooded pasture, MLV Barclay & D J Mann" (BMNH, MNB); 28 exs.: "TURKEY: Antalya Dist. Ihsaniye, 13.v.2001, 36°55'08"N 30°49'42"E 200 m, ex flooded pasture, MLV Barclay & D J Mann" (BMNH, MNB); 1 ex.: "Turquie: Cilicia Adana S. Breuning / G. Fagel sp. apud. bedeli Fauv. / [green round label] / Coll. R. I. Sc. N. B." (IRSNB); 1 ex.: "Zypern-N. GIRNE/ KYRENIA. 3.-17.4.1992, leg. Winkelmann-Klöck" (MNB).

Etymology: The specific epithet is derived from the name of the similar *B. bedeli*.

Description: Habitus as in Fig. 9. Size similar to that of *B. gallicus* (GRAVENHORST, 1806), body length 4.50 to 5.55 mm, length of forebody 2.40 to 2.85 mm. Head and pronotum blackish; elytra pale-brown to dark-brown, sometimes slightly darkened along suture; abdomen pitchy-brown to blackish, with posterior segments from apex of segment VII somewhat paler; antennae yellowish, with the apical antennomeres sometimes slightly darker; legs yellowish to pale brown.

Head moderately large, approximately 0.95 mm broad (between eyes approximately 0.65 mm). Eyes short (eye length 0.23 mm) and distinctly protruding. Surface of head dull, with distinct microsculpture, setiferous punctation very indistinct, rough, and shallow. Labrum moderately transverse, with distinct microsculpture and two small and round tubercles at apical margin. Mandibles tridentate in male and bidentate in female. Antenna of male elongated, antennomeres IV to VI slightly longer than broad, VII to X as long as broad; antenna of female slightly shorter, penultimate antennomeres as long as wide or weakly transverse.

Pronotum bulging, slightly transverse, approximately 1.2 times as broad as long, midlongitudinal sulcus short, obsolete especially in anterior portion; explanate margins entirely visible in dorsal view; posterior angles obsolete; protergosternal suture present, procoxal fissure closed; dorsal surface with weak shine; microsculpture distinct, less fine than that of head; setiferous punctation distinct, of similar density, but much deeper than that of head.

Elytra moderately short, as wide as pronotum and weakly transverse (length from humeral angles/width 0.90–0.99), slightly extended backwards; epipleural ridge complete; posterior margin with small membranous lobe; surface glossy, without visible microsculpture; setiferous punctation finer and denser than that of the pronotum. Protibia with two rows of spine-like setae; tarsal formula 4-4-4.

Abdomen with distinct transverse microsculpture, meshes short in anterior tergal impressions, much more transverse in posterior tergal portions. Setiferous punctation inconspicuous, of similar density as that of the elytra, but finer; posterior margin margin of tergite VIII emarginate.

♂: posterior margin of sternite VII (Figs 70–71) almost completely rounded, with shallow membranous emargination in the middle of the posterior margin (membranous emargination mostly indistinct in specimens from Greek islands, more distinct in specimens from southern Anatolia and Lebanon); aedeagus as in Figs 68–69.

Remark: The specimens from the type locality and other material collected by J. Frisch are of generally darker colouration in all respects (elytra brown, antennae and legs dark brown). This difference of colouration may be an artefact resulting from the method of preservation.

Comparative notes: This species is distinguished from the similar *B. bedeli* by larger size (*B. bedeli*: body length 3.65–4.40 mm, length of forebody 2.00–2.15 mm), longer antennae with antennomeres VIII–X not (male) or less (female) transverse, and much denser and less coarse punctation of the elytra. As all dissected specimens of *B. bedeli* are females, a comparison of the aedoeagi is not possible. Like *B. bedeli*, *B. bedelianus* belongs to the group of species with a membranous posterior emargination of the male sternite VII (see HERMAN 1986). **Distribution and natural history:** The species is known from the Greek Southern Sporades (Samos, Rhodos, Karpathos), from central southern Turkey, Lebanon, Israel, and Cyprus. The material from Samos was collected on the banks of streams.

Acknowledgements

Several colleagues provided material from the respective public and private collections under their care (see material and methods section). Special thanks are due to Thomas Forcke (Hannover) for the generous gift of Staphylinidae collected in Rhodos (2019/2020) and in Samos (2022), to Volker Brachat (Geretsried) for Staphylinidae collected in Samos in 2022, and to Heinrich Meybohm (Großhansdorf) for the most generous gift of all his Cephennium material from Greece. Alexey Shavrin (Daugavpils) provided a translation of the description of Euphanias pusanovi and of the key to species in BLINSTEIN (1976). Volker Puthz (Schlitz) identified Stenus callidus females from Samos, Mikhail Gildenkov (Smolensk) identified Carpelimus prudeki based on a photo of the aedeagus. Michael Schülke (Berlin) identified a Carpelimus gracilis female and communicated records of Staphylinidae collected in Rhodos by David Wrase. Volker Brachat, Heinrich Meybohm, and Michael Schülke contributed descriptions of Pselaphinae, Scydmaeninae, and Bledius, respectively. The latter three colleagues and Benedikt Feldmann (Münster) proof-read the manuscript.

References

- Assing, V. 2001: A revision of *Pseudosemiris* MACHULKA, 1935. II. A new species from Turkey (Coleoptera, Staphylinidae, Aleocharinae). – Entomologische Blätter **97** (1): 79–82.
- Assing, V. 2004: A revision of *Callicerus* GRAVENHORST, *Pseudosemiris* MACHULKA, and related genera. III. A new genus and species from China, and additional records (Insecta: Coleoptera: Staphylinidae: Aleocharinae). – Zootaxa **767**: 1–8.
- Assing, V. 2005: Two new species and new records of Staphylinidae from the Greek island Lesbos (Insecta: Coleoptera). – Linzer Biologische Beiträge **37** (2): 1035–1046.
- Assing, V. 2006: On some species of *Oxypoda* MANNERHEIM from Turkey and adjacent regions (Insecta: Coleoptera: Staphylinidae, Aleocharinae). – Linzer Biologische Beiträge **38** (1): 277–331.
- Assing, V. 2007: On the *Oxypoda* species of Turkey and adjacent regions. II. Three new species, additional records, and a checklist (Coleoptera: Staphylinidae, Aleocharinae). Zootaxa **1411**: 1–24.

- Assing, V. 2011: On the Staphylinidae of Turkey VIII. Eleven new species, two new synonymies, a new combination, and additional records (Coleoptera: Staphylinidae). – Koleopterologische Rundschau **81**: 179–227.
- Assıng, V. 2013: On the Staphylinidae of Rhodes, Greece (Insecta: Coleoptera). – Linzer Biologische Beiträge **45** (2): 1587–1613.
- Assing, V. 2015a: On the Staphylinidae of the Greek island Samos (Coleoptera: Staphylinidae). Koleopterologische Rundschau **85**: 81–102.
- Assıng, V. 2015b On the Staphylinidae of the Greek island Chios (Insecta: Coleoptera). – Linzer Biologische Beiträge 47 (1): 43–55.
- Assing, V. 2016a: On the Staphylinidae of the Greek island Karpathos (Insecta: Coleoptera). – Linzer Biologische Beiträge **48** (1): 235–263.
- AssING, V. 2016b: On the Staphylinidae of the Greek island Lesbos II, with supplementary notes on the fauna of Samos and Chios (Coleoptera: Staphylinidae). – Koleopterologische Rundschau **86**: 103–138.
- Assing, V. 2017a: On the Staphylinidae of the Greek island Kos, with an appendix on Carabidae and additional records from other islands (Insecta: Coleoptera). – Linzer Biologische Beiträge **49** (1): 191–205.
- Assing, V. 2017b: On the Staphylinidae of the Greek island Ikaría, with supplementary notes on the fauna of Samos (Coleoptera: Staphylinidae). – Koleopterologische Rundschau **87**: 89–116.
- Assing, V. 2018a: Revision of the *Cousya* species of the West Palaearctic Region (Coleoptera: Staphylinidae: Aleocharinae). Integrative Systematics 1: 79–127.
- Assing, V. 2018b: On some *Myllaena* species in the East Mediterranean and Caucasus regions (Coleoptera: Staphylinidae: Aleocharinae). – Linzer Biologische Beiträge **50** (2): 1015–1032.
- AssING, V. 2018c: On the taxonomy and zoogeography of some West Palaearctic *Quedius* species, with a focus on the East Mediterranean and the species allied to *Quedius umbrinus* and *Q. nivicola* (Coleoptera: Staphylinidae: Staphylininae). – Linzer Biologische Beiträge **50** (1): 129–182.
- Assing, V. 2018d: On the Staphylinidae of Crete III. The first records of endogean fauna (Coleoptera: Staphylinidae: Leptotyphlinae, Aleocharinae). – Linzer Biologische Beiträge **50** (1): 7–15.
- AssING, V. 2019a: Revision of the *Baeoglena* species of the West Palaearctic Region (Coleoptera: Staphylinidae: Aleocharinae). Contributions to Entomology **69** (1): 1–32.
- Assıng, V. 2019b: On the Staphylinidae of the Greek island Samothraki (Insecta, Coleoptera). – Linzer Biologische Beiträge **51** (2): 881–906.
- Assing, V. 2019c: Monograph of the Staphylinidae of Crete (Greece). Part I. Diversity and endemism (Insecta: Coleoptera). – Contributions to Entomology 69 (2): 197–238.

- ASSING, V. 2019d: Leptotyphlinae; pp. 261–265. In: ASSING, V.; BRACHAT, V. & MEYBOHM, H.: Monograph of the Staphylinidae of Crete (Greece). Part II. Descriptions of new species (Insecta: Coleoptera). – Contributions to Entomology **69** (2): 239–289.
- Assing, V. 2019e: On the taxonomy of some West Palaearctic *Quedius* species, with descriptions of new species and new synonymies (Coleoptera: Staphylinidae: Staphylininae). – Linzer Biologische Beiträge **51** (1): 189–201.
- Assing, V. 2020: On the taxonomy and zoogeography of some West Palaearctic *Cypha* species (Coleoptera: Staphylinidae: Aleocharinae). – Acta Musei Moraviae, Scientiae Biologicae **105** (1): 11–26.
- AssıNG, V. 2021: On the taxonomy of Parocyusa, Tectusa, and miscellaneous genera of Oxypodina (Coleoptera: Staphylinidae: Aleocharinae: Oxypodini). – Annalen des Naturhistorischen Museums in Wien – Serie B **123**: 99–218.
- AssING, V. (in press): On the taxonomy of *Ischnoglossa* KRAATZ, 1856: two new synonymies and a new combination (Coleoptera: Staphylinidae: Aleocharinae: Oxypodini). – Linzer Biologische Beiträge **54** (2) (2022).
- ASSING, V.; BRACHAT, V. & MEYBOHM, H. 2019: Monograph of the Staphylinidae of Crete (Greece). Part II. Descriptions of new species (Insecta: Coleoptera). – Contributions to Entomology **69** (2): 239–289.
- Assing, V. & Метвонм, H. 2021a: The *Cephennium* fauna of Turkey and the Middle East (Coleoptera: Staphylinidae: Scydmaeninae). – Contributions to Entomology **71** (1): 1–29.
- ASSING, V. & MEYBOHM, H. 2021b: On the Cephennium fauna of Greece (Coleoptera, Staphylinidae, Scydmaeninae). – Linzer Biologische Beiträge **52** (2): 869–889.
- ASSING, V. & VOGEL, J. 2019: The mother of synonyms: on the *Meotica* species of the Palaearctic Region (Coleoptera, Staphylinidae, Aleocharinae, Oxypodini).
 – Linzer Biologische Beiträge 51 (2): 731–772.
- BENICK, G. 1940: Acht neue deutsche Atheten. (Col. Staph.). Entomologische Blätter **36** (6): 180–186.
- BLINSTEIN, S. Y. 1976: Novy i maloizvestiye vidy stafilinid (Coleoptera, Staphylinidae) iz severo-zapadnogo Prichernomorya i Prisivashya. – Vestnik Zoologii 1976 (5): 78–81.
- BORDONI, A. 2020: A new *Throbalium* MULSANT & REY, 1878 from the island of Kos (Insecta: Coleoptera: Staphylinidae: Paederinae). – Quaderno di Studi e Notizie di Storia Naturale della Romagna **52**: 223–225.
- BRUNDIN, L. 1944: Monographie der palaearktischen Arten der Atheta-Untergattung Hygroecia (Coleoptera, Staphylinidae). – Annalen des Naturhistorischen Museums in Wien 53 [1942]: 129–301, pl. 13–40.
- COIFFAIT, H. 1980: Staphylinides nouveaux de la région méditerranéenne. – Nouvelle Revue d'Entomologie **10**: 251–264.

- COIFFAIT, H. 1982: Coléoptères Staphylinidae de la région paléarctique occidentale. IV. Sous famille Paederinae.
 Tribu Paederini 1 (Paederi, Lathrobii). – Nouvelle Revue d'Entomologie, Supplément 12 (4): 1–440.
- FAGEL, G. 1968: Contribution a la connaissance des Staphylinidae CVI. – Remarques diverses sur des espèces de la région paléarctique occidentale. – Bulletin et Annales de la Société Royale d'Entomologie de Belgique **104**: 189–204.
- FAUVEL, A. 1878: Les Staphylinides de l'Afrique boréale.
 Bulletin de la Société Linnèenne de Normandie (ser. 3) 2: 83–162.
- GIACHINO, P. M. & VAILATI, D.: Nuovi Anillini (Coleoptera Carabidae) di Grecia e Anatolia. – Memorie della Società Entomologica Italiana 95: 3–44.
- GILDENKOV, M. Y. 2013: Novye vidy roda *Carpelimus* LEACH, 1819 iz Palearktiki i vostochnoy oblasti (Coleoptera, Staphylinidae, Oxytelinae). – Izvestia Smolenskogo Gosudarstvennogo Universiteta **22** (2013/2): 167–187.
- GILDENKOV, M. Y. 2019: Novye dannye o raspostraneny *Carpelimus* LEACH, 1819 (Coleoptera, Staphylinidae) v Palearktike. – Nauka, obrazovanie, oshchestvo 3. Vestnik hauchnykh konferenzy **43**: 56–58.
- GILDENKOV, M. Y. & MANTIČ M. 2020: New records of the genus *Carpelimus* LEACH, 1819 (Coleoptera: Staphylinidae: Oxytelinae) from the Palaearctic region. – Acta Biologica Universitatis Daugavpiliensis **20** (1): 13–20.
- HERMAN, L. H. 1986: Revision of *Bledius*. Part IV. Classification of species groups, phylogeny, natural history, and catalogue (Coleoptera, Staphylinidae, Oxytelinae). – Bulletin of the American Museum of Natural History **184** (1): 1–386.
- Косн, С. 1937: Ueber einige Staphylinidae aus dem östlichen Mediterrangebiet. – Pubblicazioni del Museo Entomologico "Pietro Rossi" (Duino) **2**: 229–264.

- Koch, C. 1939: Neue und wenig bekannte paläarktische Paederinae. (Col. Staph.). – Mitteilungen der Münchner Entomologischen Gesellschaft **29**: 429–442.
- KOCHER, L. 1958: Catalogue commenté des coléoptères du Maroc. Fascule II. Hydrocanthares, Palpicornes, Brachelytres. – Travaux de l'Institut Scientifique Chérifien. Série Zoologie no. 14: 5–246.
- KRAATZ, G. 1858: Beitrag zur K\u00e4ferfauna Griechenlands.
 Zweites St\u00fcck: Palpicornia, Silphales, Scydmaenidae,
 Pselaphidae, Staphylinidae. Berliner Entomologische Zeitschrift 2: 37–67.
- MAKRANCZY, GY. 2014: Revision of the genus Ochthephilus MULSANT & REY, 1856 (Coleoptera, Staphylinidae, Oxytelinae). – Revue Suisse de Zoologie **121** (4): 457–694.
- OERTZEN, E. v. 1887: Verzeichniss der Coleopteren Griechenlands und Cretas, nebst einigen Bemerkungen über ihre geographische Verbreitung und 4 die Zeit des Vorkommens einiger Arten betreffenden Sammelberichten. – Berliner Entomologische Zeitschrift **30** [1886] (2): 189–293.
- SCHÜLKE, M. 2019: Zur Taxonomie und Verbreitung einiger westpaläarktischer Arten der Gattung *Planeustomus* JACQUELIN DU VAL (Coleoptera, Staphylinidae, Oxytelinae). – Linzer Biologische Beiträge 51 (2): 1315–1324.
- SCHÜLKE, M. & SMETANA, A. 2015: Staphylinidae, pp. 304–1134. – In: LÖBL, I. & LÖB, D. (eds), Catalogue of Palaearctic Coleoptera. New, updated Edition. Volume 2. Hydrophiloidea – Staphylinoidea. Revised and updated edition. – Leiden, Brill: xxvi + 1702 pp.
- Yoo, I.-S.; SONG, J.-H. & AHN, K.-J. 2021: Total-evidencer phylogeny of the coastal Cafius complex (Coleoptera: Staphylinidae) and its new generic concepts. – Zoologica Scripta 50 (6): 734–751.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Beiträge zur Entomologie = Contributions to Entomology

Jahr/Year: 2022

Band/Volume: 72

Autor(en)/Author(s): Assing Volker

Artikel/Article: <u>Supplement to the Staphylinidae fauna of Crete and the Aegean</u> <u>Islands, Greece (Coleoptera) 213-253</u>