CONTRIBUTIONS TO ENTOMOLOGY

Research Article

Fossil Trichoptera embedded in mid-Cretaceous Burmese amber*

Wilfried Wichard¹

1 University of Koeln, Institute of Biology, Köln, Germany

https://zoobank.org/EC125FAC-9784-4575-A691-2FA6AEE0AA3D

Corresponding author: Wilfried Wichard (Wichard@uni-koeln.de)

Academic editor: Astrid Schmidt-Kloiber | Received 29 July 2023 | Accepted 15 October 2023 | Published 2 November 2023

Abstract

The paper gives an overview of Trichoptera found as adults in mid-Cretaceous Burmese amber from about 100 million years ago. Fifty-eight extinct species are listed, three of which are still described here: *Paduniella cretacea* **sp. nov.**, *Palerasnitsynus vilarinoi* **sp. nov.**, *Palleptocerus kuranishii* **sp. nov.** The extinct subfamily Palerasnitsyninae **stat. nov.** of the family Xiphocentronidae is established and the extinct *Bipectinata orientalis* **comb. nov.** is transferred from the family Calamoceratidae to the family Odonto-ceridae. The extinct family Lepidochlamidae Wang et al., 2022, **stat. nov.** is transferred to the superfamily Leptoceroidea.

The fifty-eight caddisflies of Burmese amber are distributed among twenty-one genera and fourteen families, of which fifteen genera and four families are also extinct. The large time distance between extinct and extant organisms makes the assignment to the extant genera and families difficult, because the higher taxa are defined according to the species living today and often do not or hardly correspond to the earlier species and their adaptations. Furthermore, in line with the hypothesis of a Gondwanan origin of Burmese amber, some embedded Trichopterans are discussed as relict descendants of Gondwanan Trichoptera, e.g. the family Palleptoceridae and the Xiphocentronid subfamily Palerasnitsyninae.

Key Words

Evolutionary history, Fossil record, Gondwana, Trichoptera checklist, West Burma Block

Introduction

TDA Cockerell described a fossil species embedded in Burmese amber, *Plecophlebus nebulosus* Cockerell (1917), and assigned it to the order Trichoptera. A later examination of the holotype by Botosaneanu (1981) showed that this species does not belong to the order Trichoptera, but to Homoptera (Auchenorrhyncha). However, at the same time Botosaneanu described another finding in Burmese amber that clearly belonged to the order Trichoptera: *Burminoptila bemeneha* Botosaneanu (1981) (Hydroptilidae). It took 24 years for a second description of a caddisfly, *Wormaldia myanmari* Wichard & Poinar, 2005 (Philopotamidae). This marked the beginning of the continuous study of the Trichoptera in mid-Cretaceous Burmese amber. To date, 58 species of caddisflies have been described, distributed among 21 genera and 14 families. In this paper, all described species are presented in an updated checklist, including the institutional repositories of all their deposited holotypes. In addition to this overview, special attention is given to the superfamilies Psychomyioidea and Leptoceroidea, which include remarkable extinct species embedded in mid-Cretaceous Burmese amber.

Materials and methods

The amber material was collected by local people in the Hukawng Valley of northern Myanmar, (Myitkyina District, Kachin State) (Fig. 1) and derives from an amber bearing layer, which is not exposed to surface but is extending at a depth of 2-15 m (Cruickshank and Ko 2003). The amber dates from the middle Cretaceous (Cenomanian) period about 98.8 ± 0.6 Ma ago (Shi et al. 2012).

* The paper is part of 17th International Symposium on Trichoptera, Edited by Simon Vitecek, Astrid Schmidt-Kloiber, Wolfram Graf, Hans Malicky.

Copyright Wilfried Wichard. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Figure 1. Camp and preparations for Burmese amber digging in the Hukawng Valley (from Bo Wang).

The Burmese amber with the embedded Trichopteran inclusions was cut, facegrinded and polished using a cutting machine and a polishing machine, a RotoPol-25 (Struers), with grinding paper for metallography: 800, 1200, 2500, and 4000 grit. Colour photographs were produced for the documentation of specimens. A Leica M420 macroscope with Apozoom 1:6 was used in combination with Canon EOS 80D and Canon EOS R, EOS 3.0 utility software and Zerene Stacker software. Measurements made with a Leica SApo ocular micrometer.

Depositories

All listed and described holotype specimens of Trichoptera in mid-Cretaceous Burmese amber are kept and preserved in the following institutional repositories:

NIGP	Nanjing Institute of Geology and Palae-
	ontology, Nanjing, China
CNUB	Capital Normal University, Beijing, China
RPX	Ruipoxuan Amber Museum Jinan, China
ZFMK	Zoological Research Museum Alexan-
	der Koenig, Bonn, Germany

SMNS	Staatliches Museum für Naturkunde
	Stuttgart, Germany
ZSM	Zoologische Staatssammlung München,
	Germany
MP.I	Museum für Naturkunde Berlin, Germany
B.M.(N.H.)	British Museum (Natural History),
	London, GB
Coll POINAR	Oregon State University, Corvallis, USA

Abbreviations

Wing venation:

I, II, III, IV, V	=	apical forks I, II, III, IV, V.
TC	=	thyridial cell.

Male genitalia:

inf app	=	inferior appendage.
pre app	=	preanal appendage (cercus).
phal	=	phallic apparatus
harp	=	harpago (apical segment of an infe-
		rior appendage).
coxo	=	coxopodite (basal segment of an in-
		ferior appendage).
int pro	=	intermediate process.
med pro	=	medio-distad process.
IX	=	abdominal segment IX.
		-

Checklist of the Trichoptera in mid-Cretaceous Burmese amber

The following current checklist (Table 1) includes 58 extinct species, distributed among 21 genera and 14 families. Of the 14 families, 6 families belong to the suborder Annulipalpia, the 8 other family to the suborder Integripalpia, of which 4 families are extinct and only found in Burmese Amber. Of the 21 listed genera, the majority of 15 genera is extinct and only 6 genera are extant.

Table 1. Fossil Trichoptera embedded in mid-Cretaceous Burmese amber. Species checklist with evidence of deposited holotypes.

ZFMK	TRI000817
CNUB	TRI-MA-2016505
NIGP	200021
SMNS	2305 22015
NIGP	200022
NIGP	200023
ZFMK	TRI000834
NIGP	200024
NIGP	170801
ZSM	TRI-AMB001
ZFMK	TRI000823
	ZFMK CNUB NIGP SMNS NIGP ZFMK NIGP NIGP ZSM ZFMK

Amberclipsis simplex Wichard, Müller & Fischer, 2022	ZFMK	TRI000824
Protoclipsis picteti Wichard, Müller & Fischer, 2022	ZFMK	TRI000826
Protoclipsis roeseli Wichard, Müller & Fischer, 2022	ZFMK	TRI000827
Protoclipsis ulmeri Wichard, Müller & Fischer, 2022	ZFMK	TRI000825
Family Psychomyiidae Walker, 1852		
Paduniella cretacea sp. nov.	ZFMK	TRI000835
Family Xinhocentronidae Ross 1949		
Subfamily *Palerasnitsvninae stat, nov.		
Palerasnitsvnus ohlhoffi Wichard Ross & Ross 2011	NIGP	157001
Palerasnitsynus furcatis Wichard Müller & Wang 2018	MBI	7304
Palorasnitsynus gracilis Wichard, Müller & Wang, 2018	NIGP	154981
Palarasnitsynus lanidus Wichard Müller & Wang 2018	MBI	7300
Palarasnitsynus spinosus Wichard Müller & Wang, 2010	MB.I	7200
Palayagnitaynus subalabalus Wighard Müller & Wang, 2018	NIGD	154096
Paloragnitannus subgrouodis Wichard, Müller & Wang, 2018	NICD	154980
Palanganitannus gubatahanga Wighard Müller & Wang, 2018	MDI	7208
Palerasniisynus sukaichevae wichaid, Muller & Wang, 2018	MD.I	7298
Palerashisynus vulgaris wichard, Muller & Wang, 2018	MB.I	/301
Palerasnitsynus vitarinoi sp. nov.	ZFMK	1 K1000836
Family Philopotamidae Stephens, 1829	NICD	150010
Wormaldia cercifurcata Wichard, Muller & Wang, 2020	NIGP	1/2212
Wormaldia cercilonga Wichard, Müller & Wang, 2020	NIGP	172211
Wormaldia cretacea Wichard & Wang, 2016	NIGP	156999
Wormaldia myanmari Wichard & Poinar, 2005	Coll POINAR	
Wormaldia resina Wichard & Wang, 2016	NIGP	157000
Wormaldia squamosa Wichard, Müller & Wang, 2020	ZFMK	TR1000821
Wormaldia transversa Wichard, Müller & Wang, 2020	ZFMK	TR1000820
Wormaldia diplobifurca Wang, Zhang, Shi & Ren, 2021	CNUB	TRI-MA-2016506
Wormaldia denticulata Wang, Zhang, Shi & Ren, 2021	CNUB	TRI-MA-2016507
Suborder Integripalpia Martinov, 1924:		
Family Hydroptilidae Stephens, 1836		
Subfamily "Burminoptilinae Wichard, 2021		
	DMOLU	20100
Burminoptila bemeneha Botosaneanu, 1981	B.M.(N.H.)	20180
Burminoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021	B.M.(N.H.) NIGP	20180 163573
Burminoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Breitentoria Weaver, 1984:	B.M.(N.H.) NIGP	20180 163573
Burminoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912	B.M.(N.H.) NIGP	20180 163573
Burminoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018	B.M.(N.H.) NIGP RPX	20180 163573 18001
Burminoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018 Family †Burmapsychidae Wichard, 2021	B.M.(N.H.) NIGP RPX	20180 163573 18001
Burminoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018 Family †Burmapsychidae Wichard, 2021 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018	B.M.(N.H.) NIGP RPX NIGP	20180 163573 18001 166872
Burminoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018 Family †Burmapsychidae Wichard, 2021 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche palpifurcata Wichard, Neumann, Müller, & Wang, 2018	B.M.(N.H.) NIGP RPX NIGP NIGP	20180 163573 18001 166872 166873
Burmapsyche palpifurcata Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche wolframmeyi Wichard, 2021	B.M.(N.H.) NIGP RPX NIGP NIGP SEHU	20180 163573 18001 166872 166873 54024
Burminoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018 Family †Burmapsychidae Wichard, 2021 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche palpifurcata Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche wolframmeyi Wichard & Kuranishi, 2023 Family †Cretapsychidae Wichard, 2021	B.M.(N.H.) NIGP RPX NIGP NIGP SEHU	20180 163573 18001 166872 166873 54024
Burminoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018 Family †Burmapsychidae Wichard, 2021 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche palpifurcata Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche wolframmeyi Wichard & Kuranishi, 2023 Family †Cretapsychidae Wichard, 2021 Cretapsyche circula Wichard, Neumann, Müller & Wang, 2018	B.M.(N.H.) NIGP RPX NIGP NIGP SEHU MB.I.	20180 163573 18001 166872 166873 54024 7271
Burminoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018 Family †Burmapsychidae Wichard, 2021 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche palpifurcata Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche volframmeyi Wichard & Kuranishi, 2023 Family †Cretapsychidae Wichard, 2021 Cretapsyche circula Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018	B.M.(N.H.) NIGP RPX NIGP SEHU MB.I. MB.I.	20180 163573 18001 166872 166873 54024 7271 7273
Burminoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018 Family †Burmapsychidae Wichard, 2021 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche palpifurcata Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche volframmeyi Wichard & Kuranishi, 2023 Family †Cretapsychidae Wichard, 2021 Cretapsyche circula Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche insueta Wichard, Neumann, Müller & Wang, 2018	B.M.(N.H.) NIGP RPX NIGP SEHU MB.I. MB.I. MB.I.	20180 163573 18001 166872 166873 54024 7271 7273 7272
Burminoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018 Family †Burmapsychidae Wichard, 2021 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche palpifurcata Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche wolframmeyi Wichard & Kuranishi, 2023 Family †Cretapsychidae Wichard, 2021 Cretapsyche circula Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche insueta Wichard, Neumann, Müller & Wang, 2018 Cretapsyche palpinova Wichard & Neumann, Müller & Wang, 2018	B.M.(N.H.) NIGP RPX NIGP NIGP SEHU MB.I. MB.I. MB.I. MB.I. MB.I.	20180 163573 18001 166872 166873 54024 7271 7273 7272 7340
Burminoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018 Family †Burmapsychidae Wichard, 2021 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche palpifurcata Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche wolframmeyi Wichard & Kuranishi, 2023 Family †Cretapsychidae Wichard, 2021 Cretapsyche circula Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche insueta Wichard, Neumann, Müller & Wang, 2018 Cretapsyche palpinova Wichard & Neumann, Müller & Wang, 2018 Cretapsyche palpinova Wichard & Neumann, 2019 Cretapsyche kachini Wichard & Espeland, 2022	B.M.(N.H.) NIGP RPX NIGP NIGP SEHU MB.I. MB.I. MB.I. MB.I. ZFMK	20180 163573 18001 166872 166873 54024 7271 7273 7272 7340 TRI000829
Burminoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018 Family †Burmapsychidae Wichard, 2021 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche palpifurcata Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche wolframmeyi Wichard & Kuranishi, 2023 Family †Cretapsychidae Wichard, 2021 Cretapsyche circula Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche insueta Wichard, Neumann, Müller & Wang, 2018 Cretapsyche palpinova Wichard & Neumann, Müller & Wang, 2018 Cretapsyche kachini Wichard & Espeland, 2022 Cretapsyche kachini Wichard & Espeland, 2022	B.M.(N.H.) NIGP RPX NIGP NIGP SEHU MB.I. MB.I. MB.I. ZFMK ZFMK	20180 163573 18001 166872 166873 54024 7271 7273 7272 7340 TRI000829 TRI000830
Burminoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018 Family †Burmapsychidae Wichard, 2021 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche palpifurcata Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche wolframmeyi Wichard & Kuranishi, 2023 Family †Cretapsychidae Wichard, 2021 Cretapsyche circula Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche insueta Wichard, Neumann, Müller & Wang, 2018 Cretapsyche palpinova Wichard & Neumann, Müller & Wang, 2018 Cretapsyche kachini Wichard & Espeland, 2022 Cretapsyche myanmari Wichard & Espeland, 2022 Family Calamoceratidae Ulmer, 1916	B.M.(N.H.) NIGP RPX NIGP SEHU MB.I. MB.I. MB.I. MB.I. ZFMK ZFMK	20180 163573 18001 166872 166873 54024 7271 7273 7272 7340 TRI000829 TRI000830
Burminoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018 Family †Burmapsychidae Wichard, 2021 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche palpifurcata Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche wolframmeyi Wichard & Kuranishi, 2023 Family †Cretapsychidae Wichard, 2021 Cretapsyche circula Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche insueta Wichard, Neumann, Müller & Wang, 2018 Cretapsyche palpinova Wichard, Neumann, Müller & Wang, 2018 Cretapsyche palpinova Wichard & Neumann, 2019 Cretapsyche kachini Wichard & Espeland, 2022 Cretapsyche myanmari Wichard & Espeland, 2022 Family Calamoceratidae Ulmer, 1916 Cretaganonema dongi Wichard, Espeland & Wang, 2018	B.M.(N.H.) NIGP RPX NIGP SEHU MB.I. MB.I. MB.I. MB.I. ZFMK ZFMK ZFMK NIGP	20180 163573 18001 166872 166873 54024 7271 7273 7272 7340 TRI000829 TRI000830 154571
Burminoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018 Family †Burmapsychidae Wichard, 2021 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche palpifurcata Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche volframmeyi Wichard & Kuranishi, 2023 Family †Cretapsychidae Wichard, 2021 Cretapsyche circula Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche insueta Wichard, Neumann, Müller & Wang, 2018 Cretapsyche palpinova Wichard, Neumann, Müller & Wang, 2018 Cretapsyche palpinova Wichard & Neumann, 2019 Cretapsyche kachini Wichard & Espeland, 2022 Cretapsyche myanmari Wichard & Espeland, 2022 Family Calamoceratidae Ulmer, 1916 Cretaganonema dongi Wichard, Espeland & Wang, 2018 Family Odontoceridae Wallengren, 1891	B.M.(N.H.) NIGP RPX NIGP SEHU MB.I. MB.I. MB.I. MB.I. ZFMK ZFMK ZFMK NIGP	20180 163573 18001 166872 166873 54024 7271 7273 7272 7340 TRI000829 TRI000830 154571
Burminoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018 Family †Burmapsychidae Wichard, 2021 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche palpifurcata Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche volframmeyi Wichard & Kuranishi, 2023 Family †Cretapsychidae Wichard, 2021 Cretapsyche circula Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche palpinova Wichard & Neumann, Müller & Wang, 2018 Cretapsyche palpinova Wichard & Neumann, 2019 Cretapsyche kachini Wichard & Espeland, 2022 Cretapsyche myanmari Wichard & Espeland, 2022 Family Calamoceratidae Ulmer, 1916 Cretaganonema dongi Wichard, Espeland & Wang, 2018 Family Odontoceridae Wallengren, 1891 Bipectinata orientalis Wichard, Espeland, Müller & Wang, 2020, comb. nov.	B.M.(N.H.) NIGP RPX NIGP SEHU MB.I. MB.I. MB.I. ZFMK ZFMK ZFMK NIGP NIGP	20180 163573 18001 166872 166873 54024 7271 7273 7272 7340 TRI000829 TRI000830 154571 172206
Burminoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018 Family †Burmapsychidae Wichard, 2021 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche palpifurcata Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche volframmeyi Wichard & Kuranishi, 2023 Family †Cretapsychidae Wichard, 2021 Cretapsyche volframmeyi Wichard & Kuranishi, 2023 Family †Cretapsychidae Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche insueta Wichard, Neumann, Müller & Wang, 2018 Cretapsyche palpinova Wichard & Neumann, 2019 Cretapsyche kachini Wichard & Espeland, 2022 Cretapsyche myanmari Wichard & Espeland, 2022 Family Calamoceratidae Ulmer, 1916 Cretaganonema dongi Wichard, Espeland & Wang, 2018 Family Odontoceridae Wallengren, 1891 Bipectinata orientalis Wichard, Espeland, Müller & Wang, 2020, comb. nov. Palaeopsilotreta xiai Wichard & Wang, 2017	B.M.(N.H.) NIGP RPX NIGP SEHU MB.I. MB.I. MB.I. MB.I. ZFMK ZFMK ZFMK NIGP NIGP	20180 163573 18001 166872 166873 54024 7271 7273 7272 7340 TRI000829 TRI000830 154571 172206 164781
Burminoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018 Family †Burmapsychidae Wichard, 2021 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche palpifurcata Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche volframmeyi Wichard & Kuranishi, 2023 Family †Cretapsychidae Wichard, 2021 Cretapsyche circula Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche insueta Wichard, Neumann, Müller & Wang, 2018 Cretapsyche kachini Wichard & Neumann, 2019 Cretapsyche kachini Wichard & Espeland, 2022 Cretapsyche myanmari Wichard & Espeland, 2022 Family Calamoceratidae Ulmer, 1916 Cretaganonema dongi Wichard, Espeland & Wang, 2018 Family Odontoceridae Wallengren, 1891 Bipectinata orientalis Wichard & Espeland, Müller & Wang, 2020, comb. nov. Palaeopsilotreta xiai Wichard & Wang, 2017 Palaeopsilotreta burmanica Wichard, Espeland, Müller & Wang, 2020	B.M.(N.H.) NIGP RPX NIGP SEHU MB.I. MB.I. MB.I. MB.I. ZFMK ZFMK NIGP NIGP ZFMK	20180 163573 18001 166872 166873 54024 7271 7273 7272 7340 TRI000829 TRI000830 154571 172206 164781 TRI000813
Burninoptila bemeha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018 Family †Burmapsychidae Wichard, 2021 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche palpifurcata Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche wolframmeyi Wichard & Kuranishi, 2023 Family †Cretapsychidae Wichard, 2021 Cretapsyche circula Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche insueta Wichard, Neumann, Müller & Wang, 2018 Cretapsyche insueta Wichard, Neumann, Müller & Wang, 2018 Cretapsyche kachini Wichard & Neumann, Müller & Wang, 2018 Cretapsyche kachini Wichard & Espeland, 2022 Cretapsyche kachini Wichard & Espeland, 2022 Family Calamoceratidae Ulmer, 1916 Cretaganonema dongi Wichard, Espeland & Wang, 2018 Family Odontoceridae Wallengren, 1891 Bipectinata orientalis Wichard & Espeland, Müller & Wang, 2020, comb. nov. Palaeopsilotreta xiai Wichard & Wang, 2017 Palaeopsilotreta burmanica Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta cretacea Wichard, Espeland, Müller & Wang, 2020	B.M.(N.H.) NIGP RPX NIGP NIGP SEHU MB.I. MB.I. MB.I. ZFMK ZFMK ZFMK NIGP NIGP NIGP ZFMK ZFMK	20180 163573 18001 166872 166873 54024 7271 7273 7272 7340 TRI000829 TRI000829 TRI000830 154571 172206 164781 TRI000813 TRI000814
Burninoptila bemeha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018 Family †Burmapsychidae Wichard, 2021 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche palpifurcata Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche wolframmeyi Wichard & Kuranishi, 2023 Family †Cretapsychidae Wichard, 2021 Cretapsyche circula Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche insueta Wichard, Neumann, Müller & Wang, 2018 Cretapsyche kachini Wichard & Neumann, Müller & Wang, 2018 Cretapsyche kachini Wichard & Espeland, 2022 Cretapsyche kachini Wichard & Espeland, 2022 Family Calamoceratidae Ulmer, 1916 Cretaganonema dongi Wichard, Espeland & Wang, 2018 Family Odontoceridae Wallengren, 1891 Bipectinata orientalis Wichard & Espeland, Müller & Wang, 2020, comb. nov. Palaeopsilotreta xiai Wichard & Wang, 2017 Palaeopsilotreta kiai Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta succini Wichard, Espeland, Müller & Wang, 2020	B.M.(N.H.) NIGP RPX NIGP SEHU MB.I. MB.I. MB.I. MB.I. ZFMK ZFMK ZFMK NIGP NIGP NIGP ZFMK ZFMK ZFMK	20180 163573 18001 166872 166873 54024 7271 7273 7272 7340 TRI000829 TRI000829 TRI000830 154571 172206 164781 TRI000813 TRI000814 TRI000822
Burminoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021Infraorder Brevitentoria Weaver, 1984:Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018Family †Burmapsychidae Wichard, 2021 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche palpifurcata Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche wolframmeyi Wichard & Kuranishi, 2023Family †Cretapsychidae Wichard, 2021 Cretapsyche circula Wichard, Neumann, Müller & Wang, 2018 Cretapsyche circula Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche insueta Wichard, Neumann, Müller & Wang, 2018 Cretapsyche insueta Wichard, Neumann, Müller & Wang, 2018 Cretapsyche kachini Wichard & Espeland, 2022 Cretapsyche myanmari Wichard & Espeland, 2022 Cretaganonema dongi Wichard, Espeland & Wang, 2018Family Odontoceridae Wallengren, 1891 Bipectinata orientalis Wichard, Espeland, Müller & Wang, 2020, comb. nov. Palaeopsilotreta xiai Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta succini Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta succini Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta succini Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta kachini Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta kachini Wichard, Müller & Xu, 2021	B.M.(N.H.) NIGP RPX NIGP SEHU MB.I. MB.I. MB.I. MB.I. ZFMK ZFMK ZFMK ZFMK ZFMK ZFMK ZFMK ZFMK	20180 163573 18001 166872 166873 54024 7271 7273 7272 7340 TRI000829 TRI000829 TRI000830 154571 172206 164781 TRI000813 TRI000814 TRI000822 175454
Burminoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021Infraorder Brevitentoria Weaver, 1984:Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018Family †Burmapsychidae Wichard, 2021 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche palpifurcata Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche wolframmeyi Wichard & Kuranishi, 2023Family †Cretapsychidae Wichard, 2021 Cretapsyche circula Wichard, Neumann, Müller & Wang, 2018 Cretapsyche circula Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche insueta Wichard, Neumann, Müller & Wang, 2018 Cretapsyche insueta Wichard, Neumann, Müller & Wang, 2018 Cretapsyche palpinova Wichard & Espeland, 2022 Cretapsyche myanmari Wichard & Espeland, 2022 Cretaganonema dongi Wichard, Espeland & Wang, 2018Family Odontoceridae Wallengren, 1891 Bipectinata orientalis Wichard, Espeland, Müller & Wang, 2020, comb. nov. Palaeopsilotreta xiai Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta succini Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta succini Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta kachini Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta kachini Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta kachini Wichard, Müller & Xu, 2021 Palaeopsilotreta kachini Wichard, Müller & Xu, 2021	B.M.(N.H.) NIGP RPX NIGP SEHU MB.I. MB.I. MB.I. MB.I. ZFMK ZFMK ZFMK ZFMK ZFMK ZFMK ZFMK ZFMK	20180 163573 18001 166872 166873 54024 7271 7273 7272 7340 TRI000829 TRI000829 TRI000830 154571 172206 164781 TRI000813 TRI000813 TRI000814 TRI000822 175454 175453
 Burminoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018 Family †Burmapsychidae Wichard, 2021 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche palpifurcata Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche wolframmeyi Wichard & Kuranishi, 2023 Family †Cretapsychidae Wichard, 2021 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche linsueta Wichard, Neumann, Müller & Wang, 2018 Cretapsyche linsueta Wichard, Neumann, Müller & Wang, 2018 Cretapsyche kachini Wichard & Neumann, Müller & Wang, 2018 Cretapsyche kachini Wichard & Espeland, 2022 Cretapsyche mannari Wichard & Espeland, 2022 Family Calamoceratidae Ulmer, 1916 Cretaganonema dongi Wichard, Espeland, Wang, 2018 Family Odontoceridae Wallengren, 1891 Bipectinata orientalis Wichard, Espeland, Müller & Wang, 2020, comb. nov. Palaeopsilotreta xiai Wichard & Wang, 2017 Palaeopsilotreta siaccini Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta succini Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta succini Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta kachini Wichard, Müller & Xu, 2021 Palaeopsi	B.M.(N.H.) NIGP RPX NIGP SEHU MB.I. MB.I. MB.I. MB.I. ZFMK ZFMK ZFMK ZFMK ZFMK ZFMK ZFMK ZFMK	20180 163573 18001 166872 166873 54024 7271 7273 7272 7340 TRI000829 TRI000829 TRI000830 154571 172206 164781 TRI000813 TRI000813 TRI000814 TRI000822 175454 175453
Burninoptila bemeneha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018 Family †Burmapsychidae Wichard, 2021 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche palpifurcata Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche wolframmeyi Wichard & Kuranishi, 2023 Family †Cretapsychidae Wichard, 2021 Cretapsyche volframmeyi Wichard & Kuranishi, 2023 Family †Cretapsychidae Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche palpinova Wichard & Neumann, Müller & Wang, 2018 Cretapsyche palpinova Wichard & Neumann, 2019 Cretapsyche kachini Wichard & Espeland, 2022 Cretapsyche kachini Wichard & Espeland, 2022 Family Calamoceratidae Ulmer, 1916 Cretaganonema dongi Wichard, Espeland, 2022 Family Odontoceridae Wallengren, 1891 Bipectinata orientalis Wichard & Espeland, Müller & Wang, 2020, comb. nov. Palaeopsilotreta xiai Wichard & Espeland, Müller & Wang, 2020 Palaeopsilotreta succini Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta succini Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta kachini Wichard, Müller & Xu, 2021 Palaeopsilotreta kachini Wichard, Müller & Xu, 2021	B.M.(N.H.) NIGP RPX NIGP SEHU MB.I. MB.I. MB.I. MB.I. ZFMK ZFMK ZFMK ZFMK ZFMK ZFMK ZFMK ZFMK	20180 163573 18001 166872 166873 54024 7271 7273 7272 7340 TRI000829 TRI000829 TRI000830 154571 172206 164781 TRI000813 TRI000813 TRI000814 TRI000822 175454 175453 TRI-MA-2015501
Burminoptila bemenha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018 Family †Burmapsychidae Wichard, 2021 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche wolframmeyi Wichard & Kuranishi, 2023 Family †Cretapsychidae Wichard, Neumann, Müller & Wang, 2018 Cretapsyche circula Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche alpinova Wichard & Neumann, Müller & Wang, 2018 Cretapsyche palpinova Wichard & Neumann, 2019 Cretapsyche myanmari Wichard & Espeland, 2022 Cretapsyche myanmari Wichard & Espeland, 2022 Family Calamoceratidae Ulmer, 1916 Cretaganonema dongi Wichard, Espeland & Wang, 2018 Family Odontoceridae Wallengren, 1891 Bipectinata orientalis Wichard, Espeland, Müller & Wang, 2020, comb. nov. Palaeopsilotreta xiai Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta suicini Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta succini Wichard, Espeland, Müller & Wan	B.M.(N.H.) NIGP RPX NIGP SEHU MB.I. MB.I. MB.I. MB.I. ZFMK ZFMK ZFMK ZFMK ZFMK ZFMK ZFMK ZFMK	20180 163573 18001 166872 166873 54024 7271 7273 7272 7340 TRI000829 TRI000830 154571 172206 164781 TRI000813 TRI000814 TRI000814 TRI000822 175454 175453 TRI-MA-2015501
Burminoptila bemenha Botosaneanu, 1981 Cretacoptila botosaneanui Wichard, 2021 Infraorder Brevitentoria Weaver, 1984: Family Helicopsychidae Ulmer, (1906) 1912 Cretahelicopsyche liuyani Wichard, Espeland, & Wang, 2018 Family †Burmapsychidae Wichard, 2021 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche comosa Wichard, Neumann, Müller, & Wang, 2018 Burmapsyche wolframmeyi Wichard & Kuranishi, 2023 Family †Cretapsychidae Wichard, 2021 Cretapsyche volframmeyi Wichard & Kuranishi, 2023 Family †Cretapsychidae Wichard, Neumann, Müller & Wang, 2018 Cretapsyche circula Wichard, Neumann, Müller & Wang, 2018 Cretapsyche elegans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche legans Wichard, Neumann, Müller & Wang, 2018 Cretapsyche alpinova Wichard & Neumann, 2019 Cretapsyche kachini Wichard & Espeland, 2022 Cretapsyche myanmari Wichard & Espeland, 2022 Family Calamoceratidae Ulmer, 1916 Cretaganonema dongi Wichard, Espeland & Wang, 2018 Family Odontoceridae Wallengren, 1891 Bipectinata orientalis Wichard, Espeland, Müller & Wang, 2020, comb. nov. Palaeopsilotreta xiai Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta sucini Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta kurmanica Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta sucini Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta kachini Wichard, Espeland, Müller & Wang, 2020 Palaeopsilotreta kachini Wichard, Müller & Xu, 2021 Palaeopsilotreta kachini Wichard, Müller & Xu, 2021 Family †Lepidochlamidae Wang et al., 2022 Family †Palleptoceridae Wichard & Müller, 2022 Palleptoceridae Wichard & Müller, 2022	B.M.(N.H.) NIGP RPX NIGP SEHU MB.I. MB.I. MB.I. MB.I. ZFMK ZFMK ZFMK ZFMK ZFMK ZFMK ZFMK ZFMK	20180 163573 18001 166872 166873 54024 7271 7273 7272 7340 TRI000829 TRI000829 TRI000830 154571 172206 164781 TRI000813 TRI000813 TRI000814 TRI000822 175454 175453 TRI-MA-2015501 TRI000831

Systematic palaeontology

Order Trichoptera Kirby, 1813 Suborder Annulipalpia Martynov, 1924 Superfamily Psychomyioidea Walker, 1852 Family Psychomyiidae Walker, 1852 Genus *Paduniella* Ulmer, 1913

Paduniella cretacea sp. nov.

https://zoobank.org/8CE659BD-2BCD-4080-ABDC-41B6DF7DCB17 Fig. 2

Holotype. Male-specimen deposited Zoological Research Museum Alexander Koenig, Bonn, Germany, Inventory no.: ZFMK-TRI000835 (ex coll. Patrick Müller)

Preservation. The fossil male is well preserved in amber. Antennae present, as well as the six-segmented maxillary palps. The labial palps are not recognizable. The forewings show the venation well, the hind wings are unfortunately hidden. The inferior appendages of the male genital are visible from ventral.

Etymology. The extinct *Paduniella* species is named after its geological age of the Cretaceous period (latin: Cretaceum).

Description. Male, forewings ca. 2.8 mm long, antennae about half as long as forewing. Maxillary palps each six-segmented (Fig. 2C); labial palps not visible. Each forewing has forks II, III, IV, and V; hindwings not visible. In ventral view (Fig. 2B), male genitalia have deep, bifurcated inferior processes each with an elongate, overhanging process that is tapered and slightly curved apically. In lateral view (Fig. 2D), a slender median process is visible.

Diagnosis. The male genitalia of *Paduniella* species have inferior appendages whose apices are either bifurcated or unbifurcated (Li and Morse 1997). The extinct *Paduniella cretacea* sp. nov. has similar to the extant species, *P. tanidai* from Japan (Nishimoto, 2011) and *P. burmana* (Johanson & Olah, 2010) from the Oriental region, bifurcated inferior appendages that are deeply incised at the apex rather than superficially notched. The fossil species is distinguished from the two extant species in the deeply bifurcated apex of the inferior appendage because one fork is needle-like, acuminate, and curved and clearly protrudes above the other stouter fork. *Paduniella cretacea* **sp. nov.** is the first fossil *Paduniella* species found in ca. 100 million year old mid-Cretaceous Burmese amber.

Family Xiphocentronidae Ross, 1949

Subfamily †Palerasnitsyninae stat. nov.

Type genus. *Palerasnitsynus* Wichard, Ross & Ross, 2011. **Subfamily diagnosis.** The species of the extinct subfamily Palerasnitsyninae are characterised by the combination of the fore and hind wings' characters: in forewings by the presence of forks II, IV, V and by the absence of forks I and III and in hind wings by the presence of forks II and V and by the absence of the forks I, III, IV (Fig. 3B). Furthermore, the adults possess latero-apical dark sporns at the 3rd maxillary palp segments (Fig. 3A).

Systematic position. The extinct genus Palerasnitsynus, with currently 10 species, was found in the Oriental mid-Cretaceous Burmese amber and initially placed in the family Psychomyiidae (Wichard et al. 2011). The wing venation is remarkably reduced, with only apical forks II, IV, V in forewings and apical forks II and V in hind wings. The reduction of forewing venations with the absence of forks I and III and of hind wing venations with the absence of forks I, III and IV is observed in some Xiphocentronids, but not in the family Psychomyiidae. This combination of reduced forewings and hindwings is significantly true for the family Xiphocentronidae. Based on these characters, it is proposed to transfer the genus Palerasnitsynus to family Xiphocentronidae and to establish an extinct subfamily Palerasnitsyninae stat. nov. Its species are the oldest Xiphocentronids with a geological age of about 100 million years.

In Burmese amber, the species of the subfamily Palerasnitsyninae are certainly among the smallest caddisflies, reaching forewing lengths of only 1.8–2.6 mm (Wichard et al. 2011, 2018b). With an abundance of almost 40% of caddisflies, *Palerasnitsynus* species dominate the picture of caddisflies in the Burmese amber. They obviously have a tendency to swarm, as up to 100 embedded individuals have been counted in some ambers (Fig. 3C).

Genus Palerasnitsynus Wichard, Ross & Ross, 2011

Type species. *Palerasnitsynus ohlhoffi* Wichard, Ross & Ross, 2011.

Genus diagnosis. The extinct genus *Palerasnitsynus* is characterised by the combination of the fore and hind wings' characters: in forewings by the presence of forks II, IV, V and by the absence of forks I and III and in hind wings by the presence of forks II and V and by the absence of the forks I, III, IV (Fig. 3B). Furthermore the adults possess latero-apical dark sporns at the 3rd maxillary palp segments (Fig. 3A).

Palerasnitsynus vilarinoi sp. nov.

https://zoobank.org/53E80DF4-3FAB-4AEA-9418-B5796D02F00D Fig. 4

Types. A small Burmese amber contains three males, which are the holotype and two paratypes attesting to the new species. The amber is deposited Zoological Research Museum Alexander Koenig, Bonn, Germany, inventory no.: ZFMK-TRI000836 (BUB 3585 ex coll. Patrick Müller).

Preservation. Two males, preserved in varying condition, are embedded in a single amber (Fig. 4A). The



Figure 2. *Paduniella cretacea* sp. nov. in mid-Cretaceous Burmese amber, male holotype (Inventory no.: ZFMK-TRI0008359. **A.** Male in ventral view; **B.** Paired bifurcated inferior appendages in ventral view; **C.** Paired 6-segmented maxillary palps; **D.** Drawing of bifurcated inferior appendages (inf app) and median process (med pro) in lateral view.

holotype is completely preserved, from dorsal and ventral view. From ventral the pair of the genital inferior appendages is clearly visible. The paratype shows the genital laterally, although not absolutely clear. Forewings and hind wings are spread apart in the paratype.

Etymology. The new species is dedicated to Albane Vilarino, Brasil, who pointed out that the extinct genus *Palerasnitsynus* may belong to the family Xiphocentronidae.

Description. Male, adult with general characters of the genus, forewing length ca. 2 mm, antennae with about 20 cylindrical flagellomeres, plus scapus and pedicellus.

Genitalia: In ventral view (Fig. 4B), the inferior appendages are almost parallel, the separation of the basal coxopodite and the attached harpago is only weakly indicated, possibly they are fused. The inside flanks of the inferior appendage are loosely equipped with long and strong setae. The basal coxopodite is somewhat elongated and



Figure 3. Genus *Palerasnitsynus* in mid–Cretaceous Burmese amber. **A.** Typical dark sporns at the 3rd maxillary palp segment; **B.** Typical presence of forks II, IV, V in forewings and of forks II and V in hind wings; **C.** *Palerasnitsynus* swarming activities.

baso-laterally rounded, the harpago probably only slightly longer than the coxopodite. The apex of each harpago bears a dark spiny head with densified spines of nearly equal length. The needled heads of the adjacent harpagoes face each other (Fig. 4B). In lateral view (Fig. 4C), the male genital shows a pair of parallel intermediate processes (int pro) which in length barely reach the level of the medio-lateral process (med pro) which originated from the coxopodite is elongate-round and bears a bush of long setae. The intermediate process is needle-shaped, somewhat broadened in the middle, and tapers toward the apex to a ventrally inclined dark tip. The pair of short preanal appendages and a short needle-like process with dark tip, dorsally inclined, are not clearly visible.

Diagnosis.: *Palerasnitsynus vilarinoi* sp. nov. is exclusively characterized by the latero-apical dark sporns at the 3rd maxillary palp segments and by the presence of forks II, IV, V in forewings and by the presence of forks II and

V in hind wings. The new species is distinguished from all other species of *Palerasnitsynus* by the typical dark spiny head of each harpago apex facing each other with their spiny heads (Fig. 4B). The elongate needle-like intermediate processes terminate in a ventrally inclined tip. However, they do not extend longitudinally beyond the medio-lateral process of the coxopodite (Fig. 4C).

Suborder Integripalpia Martynov, 1924 Superfamily Leptoceroidea Leach, 1815 Family Odontoceridae Wallengren, 1891

Bipectinata orientalis Wichard et al., 2020b, comb. nov. Fig. 5

Systematic position. *Bipectinata orientalis* was not originally placed in the family Odontoceridae, but was



Figure 4. *Palerasnitsynus vilarinoi* sp. nov. in mid–Cretaceous Burmese amber (inventory no.: ZFMK-TRI000836). **A.** Holotype (left) and paratype (right); **B.** Holotype inferior appendages in ventral view; **C.** Paratype male genitalia in lateral view.

initially assigned to Calamoceratidae (Wichard et al. 2020), because the presence of wing fork IV on the forewing is not common to extant Odontoceridae. Nevertheless, characteristic features of the family Odontoceridae are synapomorphically present in the genus *Bipectinata*, such as five-segmented maxillary palps with a terminal segment not flexible or annulated, lack of ocelli, tibial spur formula 2/4/4; in forewings fork I present, discoidal cell closed and median cell absent. In addition, genus *Bipectinata* is closely related to the odontocerid genus



Figure 5. *Bipectinata orientalis* Wichard et al., 2020b. A. Male in ventral view; B. Bipectinate antenna; C. Plesiomorphic forewing venation with five apical forks.

Palaeopsilotreta, whose common synapomorphies involve a variable forewing media and the bipectinate antennae (Fig. 5B).

In trichopteran adults a complete set of five apical forks on the forewings is clearly a plesiomorphic character (Comstock 1918; Holzenthal et al. 2007). A reduction of the original wing venation and the reduction of apical forks are derived in many adults, especially within the superfamily Leptoceroidea. However, the extinct *Bipectinata orientalis* from the middle Cretaceous is characterized by the original arrangement of five apical forks in the forewing venation (Fig. 5C). This feature distinguishes it from the closely related species of the genus *Palaeopsilotreta*.

Family †Lepidochlamidae Wang et al., 2022, stat. nov.

Type genus + species. *Lepidochlamus nodosa* Wang et al., 2022.

Family diagnosis (based on Wang et al. 2022). Ocelli absent; antenna longer than forewing, at least 71 flagellomeres preserved; maxillary palpus five-segmented, terminal palpomere V longest, not annulated, shorter than combined lengths of preceding four palpomeres. Male forewing with forks I, III, and V; all crossveins absent. Hind wing covered with a single layer of angustifoliate scales; wing venation unknown. Tibial spur formula 2/4/4.

Systematic position. The monobasic family Lepidochlamidae Wang et al., 2022 is transferred to the Integripalpian infraorder Brevitentoria and to its superfamily Leptoceroidea: *Lepidochlamus nodosa* is characterized by the absence of ocelli and by the presence of the five-segmented maxillary palps with terminal segment not annulated, the antennae longer than forewings and the mid-tibia with preapical and apical pairs of spurs. The combination of these morphological features clearly points to the superfamily Leptoceroidea (Ross 1967; Weaver 1983, 1984; Frania and Wiggins 1997; Morse 1997).

According to Wang et al. (2022), the "Lepidochlamidae, whose hindwings are covered with scales, are the sister group of all other caddisflies (Eutrichoptera)." But as a member of the monobasic family Lepidochlamidae of the superfamily Leptoceroidea it cannot be the oldest representative of the order Trichoptera, forming a sister group relationship to all other Trichoptera, as suggested.

The establishment of the new taxa by Wang et al. (2022) is an overestimation of the scales on its hind wings. Scales on the hind wings of Brevitentorian species do not enforce the exceptional position of the "early caddisfly evolution" The scales on the hind wings are not a unique feature. Single-layered scales also cover other hindwings, e.g. Helicopsyche kariona Ross, 1975, Helicopsyche boularia Ross, 1975 (see Johanson 1998) and both wings together e.g. Helicopsyche megalochari Malicky, 1974 as well as forewings, e.g. Lepidostoma americanum Flint and Wiggins, 1961, Oecetis pechana Mosely & Kimmins, 1953. In xiphocentronid genera Drepanocentron and Abaria scales are found on fore- and hindwings as well on hindwings only (Schmid 1982). Scales on the wings (also called scalelike setae or scaloid setae) are well known in Trichoptera families and are probably derived from primitive setae.

In addition, the wing vein reduction in general and here the forewing venation reduction of Lepidochlamus nodosa is clearly phylogenetically derived. The plesiomorphic complete set of five apical forks is reduced by the loss of forks II and IV, which is common in some families of Brevitentoria as well as in its family Lepidochlamidae. Moreover, the nygmata in fork II and occasionally in the thyridial cell may be absent, and also the crossveins that would occlude the discoidal, medial, and thyridial cells, are not discernable most likely due to the embedding condition in amber. In my experience, the crossveins and nygmata are very difficult or impossible to recognize in amber. Only a comparison of several specimens of the same species allows a reliable statement about the presence or absence of the nygmata and the cross veins. The fossil Lepidochlamus nodosa is based on a single specimen, moderately preserved in mid-Cretaceous Burmese amber, kept as holotype no. CNU-TRI-MA-2015501, in Capital Normal University, Beijing, China.

Family †Palleptoceridae Wichard & Müller, 2022

Type genus and species. *Palleptocerus grimaldii* Wichard & Müller, 2022.

Family diagnosis (based on Wichard and Müller 2022). Tibial spur formula 2/4/4. Antennae longer than

forewings, flagellomeres cylindrical, elongate. Maxillary palps are five-segmented, terminal segment not annulated. Male forewing fork I and V present, discoidal cell absent. Hind wing exclusively fork V present.

Palleptocerus kuranishii sp. nov.

https://zoobank.org/52C081CE-D53F-410A-81C6-C88806ED9838 Fig. 6

Holotype. Mail-specimen deposited in the Systematic Entomology Collection of Hokkaido University Museum, Japan, inventory number: SEHU-54040 (ex coll. Ryoichi B. Kuranishi: Burmite 2).

Preservation. The fossil is embedded in an oval and polished piece of amber. The adult insect is completely preserved and clearly visible in dorso-ventral aspect. The wings are folded over the body like a saddle roof, making it difficult to see the hind wings. Head is visible, long antennae incomplete in length. Legs also present. In male genitalia, only the anterior paired inferior appendages are often clearly visible ventrally, further genital structures implied visible.

Etymology. The new species is dedicated in honor of the Japanese entomologist and scientist Ryoichi B. Kuranishi. I got to know and appreciate Ryoichi on the occasion of the 17th International Symposium on Trichoptera in Lunz, Austria, September 2022.

Description. *Head*: Laterally protruded compound eyes. Ocelli not present. Filiform antennae longer than forewings – incomplete in length, probably 30–40 flagellomeres; scapus conically thickened, pedipalpus small and globular; flagellomeres cylindrical, elongate (Fig. 6B). Five-segmented maxillary palps present, 5th segment not annulated. Labial palps three-segmented, terminal segment longest.

Wings (Fig. 6C): The forewings about 6 mm long. Venation Sc and R1 straight running parallel to the wing margin. R2 + R3 forming fork I present and R4+5 simple, discoidal cells absent. Media two-branched in M1+2 and M3+4. Cu1 two-branched in Cu1a and Cu1b, forming fork V. Crossvein m-cu closing the long thyridial cell. Venation Cu2 running into A at crossvein between Cu1b and Cu2, then A with Cu2 reaching wing margin.

Tibial spurs: 2/4/4.

Genitalia (Fig. 6D, E): In ventral view, the male genitalia equipped with a pair of inferior appendages, each consisting of a dark coxopodite, narrow at the base, distad slightly broadened, overall slightly conical bent, and attached an apically attached harpago slender than the coxopodite, tapers and curves slightly toward the genital middle.

Diagnosis. Ocelli absent. Antennae longer than forewings, flagellomeres elongate. Maxillary palps five-segmented, terminal segment not annulated. Forewings light brown, slightly narrow and apically rounded. In male forewing venation with forks I and V present, discoidal cells open and thyridial cells closed, long. Tibial spurs: 2/4/4. *Palleptocerus kuranishii* sp. nov. is distinguished



Figure 6. *Palleptocerus kuranishii* sp. nov. in mid–Cretaceous Burmese amber, male holotype (inventory number: SEHU-54040). **A.** Male in lateral view; **B.** Excerpt of the antenna with cylindrical, elongate flagellomeres; **C.** Drawing of the forewing venation; **D, E.** Male genitalia with paired inferior appendages in ventral view.

from *P. grimaldii* by a slender body shape and size of 6 mm forewing length and by the form and structure of the conical inferior appendages of the male genitalia.

Discussion

It is generally accepted that a great part of Southeast Asia consists of continental blocks that came from the supercontinent Gondwana, gradually moved northward, and docked with Southeast Asia during the Mesozoic. Gondwana had already broken off into two blocks about 130 million years ago: West Gondwana (Africa and South America) and east-Gondwana (India, Madagascar, Australia, Antarctica, and New Zealand) (McIntyre et al. 2017). The so-called West Burma Block is one of the continental blocks that split off from east-Gondwana at least 120–125 million years ago (Metcalfe 1996; Scotese 2014; Westerweel et al. 2019), gradually moved northward as an island in the Tethys Ocean over the course of more than 20 million years (Heine et al. 2004; Seton et al. 2012), and is finally placed in the form of a broad band from south to north in Myanmar (Fig. 7). In northern Myanmar, the Hukawng Valley is located in the northern area of the elongated West Myanmar Block, with an area of about 5.5 square miles, surrounded by low mountains. In line with the hypothesis of Gondwanan origin, the West Burma Block and consequently the Hukawng valley are located as a Gondwanan island within the Laurasian region of Myanmar.

The Burmese amber is from the Hukawng Valley in the West Burma Block (Poinar 2018) and is dated to the mid-Cretaceous (Cenomanian) about 98.8 ± 0.6 Ma ago (Shi et al. 2012). The amber is rich in plant and animal inclusions. The predominant arthropods are currently recorded in 8 classes, 66 orders, 624 families, 1,491 genera



Figure 7. Drawing of Myanmar, with the West Burma Block in the center inside the dashed line and the Hukawng Valley in the north.

with 2,349 species (Ross 2023). The numerous finds in Burmese amber raise the question of the origin of the organisms, whether they are from Gondwana and/or Laurasia. If the hypothesis of the Gondwanan origin of the Burmese amber and its paleobiota is correct, Gondwanan organisms and their descended lineages of the ca. 20 million years of oceanic island life are to be expected first. Only after that, Laurasian organisms could have joined.

Fossil caddisflies embedded in Burmese amber have an age of about 100 million years (mid-Cretaceous) and are most likely derived from Trichoptera of Gondwana, assuming that the West-Burma-Block containing the Hukawng amber is from Gondwana. However, it cannot be excluded that caddisflies from Laurasia may appear in addition to caddisflies from Gondwana, even though the majority probably originated from Gondwana.

The current Trichoptera checklist (Table 1) includes 14 families, of which four Integripalpian families, Burmapsychidae, Cretapsychidae, Lepidochlamidae and Palleptoceridae, are extinct and only found in Burmese amber. Very probably these extinct families have Gondwanan origin. They belong to the infraorder Brevitentoria and are assigned to the superfamilies Sericostomatoidea (Burmapsychidae, Cretapsychidae) and Lepidostomatoidea (Lepidochlamidae, Palleptoceridae). Both superfamilies have their origin or at least basal stem forms in the southern hemisphere (Thomas et al. 2020). The Sericostomatoidea separated from the Leptoceroidea around 135 Ma of early Cretaceous (Malm et al. 2013), about the same time when Gondwana split into west- and east-Gondwana (McIntyre et al. 2017). After that the West Burma Block separated from the east-Gondwana. In line with the hypothesis of Gondwanan origin of the West Burma Block and the Burmese amber their extinct Trichoptera families may be descended from the Gondwanan and Neotropical infraorder Brevitentoria.

The Burmese amber continues to reveal more information about the evolutionary history of Trichoptera. For example, the family Xiphocentronidae is distributed in the Neotropic and Oriental regions. The Burmese amber is located in the Oriental region in northern Myanmar and preserves extinct species of the genus *Palerasnitsynus*, which establish the subfamily Palerasnitsyninae stat. nov. of Xiphocentronidae. However, *Palerasnitsynus* is not closely related to the Xiphocentronidae genera, which occur in the Oriental region. The extinct genus of middle Cretaceous may be descended from a Neotropical lineage of Xiphocentronidae, on the basis that the hypothesis of a Gondwanan origin of the Burmese amber is correct.

Acknowledgements

The author is especially grateful to Wolfram Mey, André Nel and Albane Vilarino for their welcome discussions and comments. The description of the Burmese Trichoptera species would not have been possible without Patrick Müller and Bo Wang, who provided the Burmese amber with embedded caddisflies.

References

- Botosaneanu L (1981) On a false and a genuine caddisfly from Burmese amber (Insecta: Trichoptera, Homoptera). Bulletin Zoologisch Museum, Universiteit Van Amsterdam 8: 73–78.
- Comstock J (1918) The Wings of Insects. Ithaca, NY: Comstock Publishing Company, 430 pp.
- Cruickshank RD, Ko K (2003) Geology of an amber locality in the Hukawng Valley, Northern Myanmar. Journal of Asian Earth Sciences 21: 441–455. https://doi.org/10.1016/S1367-9120(02)00044-5
- Flint OS, Wiggins GB (2012) Records and descriptions of North American species in the genus *Lepidostoma*, with a revision of the *vernalis* group (Trichoptera: Lepidostomatidae). The Canadian Entomologist 93(4): 279–297. https://doi.org/10.4039/Ent93279-4
- Frania HE, Wiggins GB (1997) Analysis of morphological and behavioural evidence for the phylogeny and higher classification of Trichoptera (Insecta). Royal Ontario Museum, Life Sciences Contributions 160: 1–68. https://doi.org/10.5962/bhl.title.53487
- Heine C, Müller RD, Gaina C (2004) Reconstructing the lost eastern Tethys ocean basin: convergence history of the SE Asian margin and marine gateways. Geophysical Monograph Series 149: 1–18. https:// doi.org/10.1029/149GM03
- Holzenthal RW, Roger J, Blahnik RJ, Prather AL, Kjer KM (2007) Order Trichoptera Kirby, 1813 (Insecta), Caddisflies. Zootaxa 1668: 639–698. https://doi.org/10.11646/zootaxa.1668.1.29
- Johanson KA (1998) Phylogenetic and biogeographic analysis of the family Helicopychidae (Insecta: Trichoptera). Entomologica scandinavica Suplement 53: 1–172.
- Johanson KA, Oláh J (2010) Description of six new species of Oriental Paduniella (Insecta: Trichoptera: Psychomyiidae). Zootaxa 2548: 43–56. https://doi.org/10.11646/zootaxa.2548.1.3
- Kirby W (1813) Strepsiptera, a new order of insects proposed, and the characters of the order, with those of its genera. Transactions of the Linnean Society of London, Zoology 11: 86–122. https://doi. org/10.1111/j.1096-3642.1813.tb00040.x
- Leach WE (1815) Entomology. In: Brewster D (Ed.) The Edinburgh Encyclopedia, vol. 9: 57–172, William Blackburn, Edinburgh.
- Li YJ, Morse JC (1997) The *Paduniella* (Trichoptera: Psychomyiidae) of China, with a phylogeny of the World species. Insecta Mundi 276. https://digitalcommons.unl.edu/insectamundi/276
- Malicky H (1991) Some unusual caddisflies (Trichoptera) from southeastern Asia (Studies on caddisflies of Thailand, No. 5). In: Proceedings of the 6th International Symposium on Trichoptera, 381–384.
- Malm T, Johanson KA, Wahlberg N (2013) The evolutionary history of Trichoptera (Insecta): a case of successful adaptation to life in freshwater. Systematic Entomology 38: 459–473. https://doi.org/10.1111/ syen.12016
- Martynov AV (1924) Rucheiniki (caddisflies). Prakticheskaya Entomologiya 5: 1–384.
- McIntyre SRN, Lineweaver CH, Groves CP, Chopra A (2017) Global biogeography since Pangaea. Proceedings of the Royal Society B, Biological Sciences 284: 20170716. https://doi.org/10.1098/ rspb.2017.0716
- Metcalfe I (1996) Pre–Cretaceous evolution of SE Asian terranes. In: Hall R, Blundell D (Eds) Tectonic evolution of Southeast Asia. Geological Society Special Publication 106: 97–122. https://doi. org/10.1144/GSL.SP.1996.106.01.09

- Morse JC (1997) Phylogeny of Trichoptera. Annual Review of Entomology 42: 427–450. https://doi.org/10.1146/annurev.ento.42.1.427
- Mosely ME, Kimmins DE (1953) The Trichoptera (Caddis Flies) of Australia and New Zealand. British Museum (Natural History), London, 550 pp. https://doi.org/10.5962/bhl.title.118696
- Nishimoto H (2011) The genus *Paduniella* (Trichoptera: Psychomyiidae) in Japan. Zoosymposia 5: 381–390. https://doi.org/10.11646/ zoosymposia.5.1.30
- Poinar GO (2018) Burmese amber: evidence of Gondwanan origin and Cretaceous dispersion. Historical Biology 31: 1304–1309. https:// doi.org/10.1080/08912963.2018.1446531
- Ross AJ (2023) Supplement to the Burmese (Myanmar) amber checklist and bibliography, 2022. Palaeoentomology 006: 022–040. https:// doi.org/10.11646/palaeoentomology.6.1.6
- Ross HH (1949) Xiphocentronidae, a new family of Trichoptera. Entomological News 60: 1–7.
- Ross HH (1967) The Evolution and Past Dispersal of the Trichoptera. Annual Review of Entomology 12: 169–206. https://doi. org/10.1146/annurev.en.12.010167.001125
- Schmid F (1982) La famille des Xiphocentronidae (Trichoptera: Annulipalpia). The Memoirs of the Entomological Society of Canada 114(S121): 3–127. https://doi.org/10.4039/entm114121fv
- Scotese CR (2014) Atlas of Early Cretaceous Paleogeographic Maps, PALEOMAP Atlas for ArcGIS 2: The Cretaceous, Maps 23–31.
- Seton M, Müller RD, Zahirovic S, Gaina C, Torsvik T, Shepard G, Talsma A, Gurnis M, Turner M, Maus S, Chandler M (2012) Global continental and ocean basin reconstructions since 200 Ma. Earth-Science Review 113: 212–270. https://doi.org/10.1016/j. earscirev.2012.03.002
- Shi G, Grimaldi DA, Harlow GE, Wang J, Yang M, Lei W, Li Q, Li X (2012) Age constraint on Burmese amber based on U-Pb dating of zircons. Cretaceous Research 37: 155–163. https://doi.org/10.1016/j. cretres.2012.03.014
- Stephens JF (1829) A Systematic Catalogue of British Insects. Part 1: Insecta. Baldwin and Cradock, London, 416 pp.
- Stephens JF (1836) Illustrations of British Entomology (Mandibulata), vol. 6. Baldwin and Cradock, London, 151–154.
- Thomas JA, Frandsen PB, Prendini E, Zhou X, Holzenthal RW (2020) A multigene phylogeny and timeline for Trichoptera (Insecta). Systematic Entomology 45: 670–686. https://doi.org/10.1111/syen.12422
- Ulmer G (1903) Über die Metamorphose der Trichopteren. Abhandlungen des Naturwissenschaftlichen Vereins in Hamburg 18: 1–154.
- Ulmer G (1906) Neuer Beitrag zur Kenntnis aussereuropäischer Trichopteren. Notes from the Leyden Museum 28: 1–116.
- Ulmer G (1912) Die Trichopteren des Baltischen Bernsteins. Beiträge zur Naturkunde Preussens 10: 1–380.
- Ulmer G (1951) Köcherfliegen (Trichopteren) von den Sunda-Inseln. Tell I. Archiv f
 ür Hydrobiologie 19: 1–528.
- Walker F (1852) Cataloque of the Specimens of Neuropterous Insects in the Collection of the British Museum. Part I. Phryganides-Perlides. London (British Museum), 658 pp.
- Wallengren HDJ (1891) Skandinaviens Neuroptera. Andra afdelningen. Svenska Vetenskaps-Akademiens Handlingar 24: 1–173.
- Wang J, ZhangW, Wang L, Ren D (2019) A new caddisfly (Trichoptera: Polycentropodidae) from Upper Cretaceous amber of Myanmar. Cretaceous Research 99: 347–351. https://doi.org/10.1016/j. cretres.2019.01.021

- Wang J, Zhang W, Shih C, Ren D (2021) Two new mid-Cretaceous caddisflies (Trichoptera, Philopotamidae) from amber of Northern Myanmar. Cretaceous Research 126: 104891. https://doi. org/10.1016/j.cretres.2021.104891
- Wang J, Zhang W, Engel MS, Sheng X, Shih C, Ren D (2022) Early evolution of wing scales prior to the rise of moths and butterflies, Current Biology 32: 1–7. https://doi.org/10.1016/j.cub.2022.06.086
- Weaver III JS (1983) The evolution and classification of Trichoptera, with a revision of the Lepidostomatidae and a North America synopsis of this family. Ph. D. Dissertation, Clemson University, Clemson, 411 pp.
- Weaver III JS (1984) The evolution and classification of Trichoptera, Part I: The groundplan of Trichoptera. In: Morse JC (Ed.) Proceedings of 5th International Symposium on Trichoptera, Clemson, 1983. The Hague: W. Junk, 413–419.
- Westerweel J, Roperch P, Licht A, Dupont-Nivet G, Win Z, Poblete F, Ruffet G, Swe HH, Thi MK, Aung DW (2019) Burma Terrane part of the Trans-Tethyan arc during collision with India according to palaeomagnetic data. Nature Geoscience 12: 863–868. https://doi. org/10.1038/s41561-019-0443-2
- Wichard W (2021) Overview of the caddisflies (Insecta, Trichoptera) in mid-Cretaceous Burmese amber. Cretaceous Research 119: 104707. https://doi.org/10.1016/j.cretres.2020.104707
- Wichard W, Espeland M (2022) The family Cretapsychidae (Insecta, Trichoptera) from mid-Cretaceous Burmese amber, with descriptions of two new species. European Journal of Taxonomy 833: 1–11. https://doi.org/10.5852/ejt.2022.833.1879
- Wichard W, Kuranishi, RB (2023) Burmapsyche wolframmeyi sp. nov., a new species of the extinct family Burmapsychidae (Insecta, Trichoptera) embedded in mid-Cretaceous Burmese Amber. Palaeodiversity 15: 1–6. https://doi.org/10.18476/pale.v16.a1
- Wichard W, Müller P (2022) Palleptoceridae fam. nov., an extinct leptoceroid family in mid-Cretaceous Burmese Amber (Insecta, Trichoptera). Palaeoentomology 005: 468–474. https://doi.org/10.11646/ palaeoentomology.5.5.8
- Wichard W, Neumann C (2019) A new bizarre dysoneurid species (Insecta, Trichoptera) in Burmese amber. Fossil Record 22: 51–56. https://doi.org/10.5194/fr-22-51-2019
- Wichard W, Poinar G (2005) Köcherfliegen aus dem Burma Bernstein der oberen Kreide von Myanmar (Insecta, Trichoptera). Mitteilungen aus dem Geologisch-Palaontologischen Institut der Universitat Hamburg 89: 129–136.

- Wichard W, Wang B (2016) New Cretaceous caddisflies from Burmese amber (Insecta, Trichoptera). Cretaceous Research 61: 129–135. https://doi.org/10.1016/j.cretres.2016.01.004
- Wichard W, Wang B (2017) A remarkable caddisfly with bipectinate antennae in Cretaceous Burmese amber (Insecta, Trichoptera). Cretaceous Research 69: 198–203. https://doi.org/10.1016/j. cretres.2016.09.012
- Wichard W, Wang B (2020) Family Kambaitipsychidae (Insecta, Trichoptera) in mid-Cretaceous Burmese amber. Cretaceous Research 107: 104272. https://doi.org/10.1016/j.cretres.2019.104272
- Wichard W, Xu C-P (2022) The family Polycentropodidae (Insecta, Trichoptera) in mid-Cretaceous Burmese Amber. ZooKeys 1134: 171–183. https://doi.org/10.3897/zookeys.1134.93999
- Wichard W, Ross E, Ross A (2011) Palerasnitsynus gen. n. (Trichoptera, Psychomyiidae) from Burmese amber. ZooKeys 130: 323–330. https://doi.org/10.3897/zookeys.130.1449
- Wichard W, Neumann C, Müller P, Wang B (2018a) Family Dysoneuridae (Insecta, Trichoptera) in Cretaceous Burmese amber. Cretaceous Research 82: 138–146. https://doi.org/10.1016/j.cretres.2017.10.008
- Wichard W, Müller P, Wang B (2018b) The psychomyiid genus Palerasnitsynus (Insecta, Trichoptera) in mid-Cretaceous Burmese amber. Palaeodiversity 11: 151–166. https://doi.org/10.18476/pale.11.a8
- Wichard W, Espeland M, Wang B (2018c) Caddisflies with unusual hair-fans on the legs in Cretaceous Burmese amber (Insecta, Trichoptera). Palaeodiversity 11: 21–28. https://doi.org/10.18476/ pale.11.a3
- Wichard W, Müller P, Wang B (2020a) The philopotamid genus Wormaldia (Insecta, Trichoptera) embedded in mid-Cretaceous Burmese amber. Palaeodiversity 13: 39–47. https://doi.org/10.18476/ pale.v13.a4
- Wichard W, Espeland M, Müller P, Wang B (2020b) Burmese amber (Insecta, Trichoptera: Odontoceratidae, Calamoceratidae). European Journal of Taxonomy 653: 1–17. https://doi.org/10.5852/ ejt.2020.653
- Wichard W, Müller P, Xu C-P (2022a) The odontocerid genera *Psilo-treta* and *Palaeopsilotreta* (Insecta, Trichoptera) embedded in mid-Cretaceous Burmese amber. Palaeoentomology 4: 331–338. https://doi.org/10.11646/palaeoentomology.4.4.8
- Wichard W, Müller P, Fischer T (2022b) Family Pseudoneureclipsidae (Insecta, Trichoptera) in mid-Cretaceous Burmese amber. Palaeodiversity 15: 1–12. https://doi.org/10.18476/pale.v15.a1

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Beiträge zur Entomologie = Contributions to Entomology

Jahr/Year: 2023

Band/Volume: 73

Autor(en)/Author(s): Wichard Wilfried

Artikel/Article: Fossil Trichoptera embedded in mid-Cretaceous Burmese amber 167-179