

Corrigenda

Corrigendum: Differentiation in the ultrastructure of pectiniform antennae in species groups of the genus *Ctenoceratoda* Varga, 1992 (Lepidoptera, Noctuidae). Contributions to Entomology 73(1): 95–107. <https://doi.org/10.3897/contrib.entomol.73.e104072>

Zoltán Varga¹, László Ronkay², László Rákosy³

¹ Department of Evolutionary Zoology and Human Biology, University of Debrecen, H-4032 Debrecen, Egyetem tér 1, Hungary

² Hungarian Natural History Museum, Budapest, Hungary

³ Faculty of Biology and Geology, Babeş-Bolyai University, 44 Republicii, RO-400015 Cluj-Napoca, Romania

<https://zoobank.org/D0AB8552-C5A8-4FFA-A3FA-87067729F11B>

Corresponding author: Zoltán Varga (varga.zoltan@science.unideb.hu)

Academic editor: Thomas Schmitt | Received 11 November 2023 | Accepted 15 November 2023 | Published 8 December 2023

In the publication Varga et al. “Differentiation in the ultrastructure of pectiniform antennae” in Contributions to Entomology 73 (1) 2023, 95–107 | DOI 10.3897/contrib.entomol.73.e104072 due to a technical error, the

Figure 14 was dropped out and the Figure 15 was reproduced two times. Therefore, we replace now the correct Figure 14 with the original explanation.




Figure 14. **A.** *C. mallopyga dyschroa* Varga, Gyulai, G. Ronkay and L. Ronkay, 2017, medial part of the antenna with short, asymmetric processes and with long sensilla chaetica in terminal position, with straight or terminally curved sensilla trichoidea, 60× magnification; **B.** *C. mallopyga dyschroa* Varga, Gyulai, G. Ronkay et L. Ronkay, 2017, stalk and dorsal processes of the antenna, with long, nearly straight sensilla chaetica in terminal position, and with straight or terminally curved sensilla trichoidea, 150× magnification; **C.** *C. mallopyga dyschroa* Varga, Gyulai, G. Ronkay et L. Ronkay, 2017, stalk and ventral processes of the antenna, with long, nearly straight sensilla chaetica in terminal position, and with straight or terminally curved sensilla trichoidea, 150× magnification; **D.** *C. mallopyga dyschroa* Varga, Gyulai, G. Ronkay et L. Ronkay, 2017, distal part of the dorsal processes with long, nearly straight sensilla chaetica in terminal position, and with nearly straight or terminally curved sensilla trichoidea, 300× magnification; **E.** *C. mallopyga dyschroa* Varga, Gyulai, G. Ronkay et L. Ronkay, 2017, short ventral processes of the antenna with straight sensilla chaetica in terminal position and terminally curved sensilla trichoidea, 300× magnification; **F.** *C. mallopyga dyschroa* Varga, Gyulai, G. Ronkay et L. Ronkay, 2017, part of the stalk with relatively short sensilla trichoidea, densely covered by narrow, acute sensilla coeloconica, 1000× magnification.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Beiträge zur Entomologie = Contributions to Entomology](#)

Jahr/Year: 2023

Band/Volume: [73](#)

Autor(en)/Author(s): Varga Zoltán, Ronkay Laszlo, Rakosy Laszlo

Artikel/Article: [Corrigendum: Differentiation in the ultrastructure of pectiniform antennae in species groups of the genus Ctenoceratoda Varga, 1992 \(Lepidoptera, Noctuidae\). Contributions to Entomology 73\(1\): 95–107.](#)
<https://doi.org/10.3897/contrib.entomol.73.e104072> 249-250