CONTRIBUTIONS TO ENTOMOLOGY

Research Article

Introduced greenhouse-invertebrates in Potsdam and Berlin with a focus on ants (Hymenoptera, Formicidae) with eight new records for Europe, Germany or the Berlin-Brandenburg region

Elias Freyhof¹, Emil Janke²

1 Senckenberg Deutsches Entomologisches Institut, Eberswalder Straße 90, 15374 Müncheberg, Germany

2 Heilbornring 2, 15234 Frankfurt (Oder), Germany

https://zoobank.org/6A6FA209-BD17-450C-B117-C5C890773DBB

Corresponding author: Elias Freyhof (elifreyhof@gmail.com)

Academic editor: Thomas Schmitt | Received 18 September 2024 | Accepted 19 November 2024 | Published 20 December 2024

Abstract

Heated greenhouses are a hotspot for introduced species from warmer climates. We studied 24 greenhouses for their invertebrate fauna with a special focus on ants. Identifications were initialised by iNaturalist and its community, followed by literature, COI barcoding and consulting experts in some cases. We report 32 introduced species including eight new records at the regional, national or continental scale, four of which are ants. *Technomyrmex difficilis* and *Solenopsis texana* are recorded for the first time in Germany. *Plagiolepis alluaudi* and *Technomyrmex vitiensis* are new for Berlin and Brandenburg. *Alepia* cf. *viatrix* (Diptera, Psychodidae) is first recorded for continental Eurasia. *Cryptotermes cavifrons* (Blattodea, Kalotermitidae) and *Geonemertes pelaensis* (Hoplonemertea, Prosorhochmidae) are recorded the first time for Europe. An unidentified species of the genus *Anisorhynchodemus* (Tricladida, Geoplanidae) is recorded the first time for Germany. Here, we present records for 37 species (five of the ant species are native), all associated with pictures in iNaturalist. For 33 individuals of 20 species, we also provide COI sequence data supporting their identification. Furthermore, the comparison of greenhouse metadata with species composition showed that the introduced ant species are dependent on a high minimum temperature in the greenhouse.

Key Words

Biological invasions, COI barcoding, greenhouse fauna, iNaturalist, introduced species, new records

Introduction

The introduction of non-native species is a side effect of globalisation and the associated international trade of goods (Westphal et al. 2008). A major new report by the Intergovernmental Platform on Biodiversity and Ecosystem Services (IPBES 2023) summarises the threats posed by non-native species to the economy, food security and human health and their role in global biodiversity change. IPBES (2023) lists 37,000 species that have been introduced outside their native range. However, this process is highly dynamic, has accelerated considerably in recent years and the number of species involved may be grossly underestimated. In Germany, 1,280 non-native species are listed, of which only 119 are insects and all have been found in the wild in outdoor habitats (https://neobiota.bfn.de/grundlagen/anzahl-gebietsfremder-arten.html). The German assessment largely (but not completely) ignores non-native species that have become established in anthropogenic indoor habitats, such as heated greenhouses, industrial buildings and living rooms. These indoor species may have been considered of little relevance, as they may remain confined to indoor habitats and may not be able to pose a threat to nature and the economy. Whether this will remain the case as temperatures rise and conditions change due to climate change remains a question to be investigated in the future.

Copyright Elias Freyhof & Emil Janke. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

To create the (sub)tropical environment typical of heated greenhouses, soil and plant material have been imported from a wide range of tropical and subtropical countries and plants have also been exchanged amongst greenhouses. This results in a high potential for species not only to be introduced, but also to spread and to become established over time. The stable climatic conditions allow tropical species to build up populations and become abundant in greenhouses (Blatrix et al. 2018), creating completely new ecosystems (Hobbs et al. 2009). These species can become pests that are difficult and costly to control (Kenis and Branco 2010; Blatrix et al. 2018) and their establishment carries the risk that they, or their diseases, can spread from the greenhouses to outdoor habitats. By this, greenhouses can act as global 'disease pools' infecting other greenhouses with alien species. It is also interesting to investigate whether there is a common set of global species that occur in greenhouses and form their own ecological communities. Invertebrates, in particular, can be rapidly introduced into greenhouses and often remain undetected for long periods due to their often small size and hidden lifestyle (Stohlgren and Schnase 2006; Mehta et al. 2007). Furthermore, they are difficult to identify as identification resources often just cover national faunas, rarely including exotic species, are restricted to outdoor ecosystems or are too old to include recently arrived species (Suvák 2011; Blatrix et al. 2018; Báthori et al. 2024).

The aim of this study is to document non-native invertebrates in heated greenhouses in Berlin and Potsdam, with a particular focus on ants. Ants are amongst the most successful and damaging invaders worldwide (Lowe et al. 2000). The Global Invasive Species Database contains 19 invasive ant species and five ant species that are amongst the 100 worst invasive species (GISD 2024). So-called tramp species, which have been spread globally by human activities, can arise. The high adaptability of ants to new habitats is often due to social traits, such as polygyny, polydomy and super-colonial colony structure (Tsutsui and Suarez 2003). Tramp species in greenhouses are not able to colonise outdoors due to current climatic conditions (Boer and Vierbergen 2008). However, such tramp species are often species that are invasive in warmer climates. For example, nearly all tramp species found in this study are classified as "highly invasive" in AntWiki (https://www.antwiki.org), which means that they are "species that are essentially or actually worldwide in distribution" (https://www.antwiki.org). The only found tramp species that does not have an active AntWiki page is also a cosmopolitan tramp species (Seifert 2013). Ants in greenhouses can sometimes be very dominant and indirectly damage plants through their symbiotic relationship with hemipterans (Blatrix et al. 2018). For these reasons, their study is particularly important and ants have been collected more intensively. However, little is known about introduced invertebrate species in greenhouses in Germany and especially the occurrence of tramp ants is poorly understood. This

study aims to better assess the diversity of introduced invertebrates in greenhouses in Berlin and Potsdam. In addition, the occurrence of ant species in combination with temperature in greenhouses is assessed to better understand their lower temperature limits. This is an important factor for the spread and distribution of species. Furthermore, we aim to test whether a combination of morphological identification by AI and experts as well as genetic identification allows us to properly identify species that come from anywhere in the world.

Methods

The study examined six facilities (Fig. 1) in Berlin and Potsdam. These were divided into 24 climatic units. A climatic unit is a contiguous greenhouse or contiguous greenhouse section where the same climatic conditions prevail. For example, the Berlin Botanical Garden has many greenhouse sections that are heated to different degrees or have different air humidity. Therefore, 14 climatic units (Table 1) are distinguished. The metadata for each climatic unit (size of the greenhouse in m², use of the greenhouse, minimum temperature in °C, average humidity in %, use of insecticides or biological pest control, when the greenhouse was surveyed and the exact location of the greenhouse) are given in Table 1. These metadata were compared with the ant species composition.

The Pearson-correlation analysis was carried out in Excel (2024) to find out whether the number of introduced and native ant species and those of native species correlate with the minimum temperatures of the greenhouses.

Each greenhouse was searched manually for invertebrates. Loose objects, such as wood and stones, were turned over and the soil underneath was inspected. We also scanned the lower foliage. Ants were additionally lured with a bait of sugar-water or a mixture of fish oil, rum and honey (as recommended by Bernhard Seifert, pers. com.). The baits were placed on small (25×25 mm) plastic plates and placed approximately every four metres along the walkways throughout the greenhouses. After 30–60 minutes, the baits were checked for ants. Table 2 lists all ant species found in the greenhouses.

All species were photographed (Nikon D600 with macro lens, distance rings and tong flash) and a few individuals of each species were transferred to 90% ethanol. Georeferenced records were uploaded to the global reporting platform iNaturalist (www.inaturalist.org). All records are available on the iNaturalist profiles eliasfreyhof36614's Profile iNaturalist and emilvus's Profile iNaturalist. iNaturalist's artificial intelligence (computer vision) suggested an initial taxonomic identification, which was then critically reviewed by us. The iNaturalist community either suggested alternative species identifications or confirmed ours, a process that was an additional step towards a valid identification. These identifications were made by both amateur and expert naturalists. At least two matching identifications or two-thirds of matching



Figure 1. Location of greenhouses studied in Berlin and Potsdam. © GeoBasis-DE/LGB (2024), dl-de/by-2-0

identifications were required to achieve 'research grade' at iNaturalist. Some species that were difficult to identify from a photograph were identified morphologically and/ or genetically. Morphology was examined using binoculars and specialist literature (Table 3). Species identification remained a challenge due to the introduction of species from all over the world and the fact that literature is usually based on specific regions. Accordingly, a search was made to determine which species of a genus had already been introduced elsewhere. Suitable identification keys or publications were then searched for the potentially introduced species. Ants of the genera Technomyrmex and Solenopsis were sent to Bernhard Seifert (Görlitz) for morphometric identification. Diplopoda, Gastropoda, Hoplonemertea and Turbellaria were re-determined by experts on the basis of photographs (Table 3). The collected specimens are deposited at the Senckenberg German Entomological Institute, Müncheberg (Germany). Solenopsis spp., Technomyrmex spp., Oxidus gracilis, Leptogoniulus sorornus and Stenochrus portoricensis are deposited in the collection of the Senckenberg Museum für Naturkunde Görlitz (Germany).

DNA extraction and PCR

DNA was extracted from whole samples, leg or femur muscle tissue (stored in ethanol) using sterilised tweezers. After allowing the ethanol to evaporate, DNA was extracted using a Chelex extraction protocol (modified from Walsh et al. 1991). For each sample, 100 μ l Chelex solution (5%) and 5 μ l proteinase K (10 mg/ml) were added. All samples were then placed in a thermocycler (Labcycler Basic, SensoQuest, Göttingen, Germany) with the following program: 1 h at 55 °C, 15 min at 99 °C, 1 min at 37 °C. Extracts were stored at -20 °C.

The barcoding fragment of the mitochondrial COI gene was amplified using Qiagen DNA polymerase (QIAGEN Multiplex PCR Plus Kit, Qiagen, Hilden, Germany) at a concentration of 50% of the total reaction volume and a concentration of 33% of each primer. Primers LCO_1490 and HCO_2198 (Folmer et al. 1994) were used for the first attempt. Primers CLepFol R / CLepFol F, LCO1490-JJ2 / HCO2198-JJ2 (Astrin et al. 2016) and LCO1490-JJ / LCO1490-JJ (Astrin and Stüben 2008) were used for samples that did not work previously. The PCR programme included 38 cycles (5 min at 95 °C, $38 \times (30 \text{ s at})$ 95 °C, 90 s at 49 °C, 60 s at 72 °C), 30 min at 68 °C). The PCR products were then checked on a 1.3% agarose gel. Positive samples were purified using the ExoCleanUp FAST Kit (VWR International GmbH, Darmstadt, Germany) and sent to Macrogen Europe B.V. (Amsterdam, The Netherlands) for Sanger sequencing. Sequences were edited using Geneious (version R11) and then uploaded to BOLD Systems (v.4) (https://www.boldsystems.org/) under project ID GBAR (Table 4).

Greenhouse	Size in m ²	Usage	Minimum temperature in °C	Average air humidity in %	Insecticides or biological pest control use	Date of visit	Coordinates	Abbreviation of climatic units
Gärten der Welt (GW.), Balinese Garden ("Balinesischer Garten")	1200	plant cultivation	18	80	Biological pest control: Ichneumonidae wasps two times a month	13.11.2023	52°32.37'N, 13°34.656' E	/
FEZ., Eco Island ("Ökoinsel")	300	plant cultivation	10	45	insecticides: Ant bait boxes if needed Neudorff Loxiran https://www.neudorff.de/produkte/loxiran- ameisenkoederdose.html	24.11.2023	52°27.7533'N, 13°33.075'E	/
Botanical Garden Berlin (BGB.), house A	1900	plant cultivation	19	55	Biological pest control: Cryptolaemus montrouzieri, Anolis sagrei, Ichneumonidae, Eleutherodactylus coqui and more Insecticides: In winter, individual plants are sprayed with insecticides. If there are too many ants, they are controlled with gels.	22.11.2023	52°27.395'N, 13°18.447'E	/
BGB., house B	170	plant cultivation	18	70	as BGB.	27.11.2023, 14.05.2024	52°27.375'N, 13°18.467'E	В
BGB., house C	500	plant cultivation	17	70	as BGB.	27.11.2023, 14.05.2024	52°27.365'N, 13°18.478'E	С
BGB., house D	135	plant cultivation	15	70	as BGB.	27.11.2023, 14.05.2024	52°27.357'N, 13°18.462'E	D
BGB., house E	200	plant cultivation	17	58	as BGB.	27.11.2023, 14.05.2024	52°27.350'N, 13°18.445'E	Е
BGB., house F	370	plant cultivation	17	70	as BGB.	27.11.2023, 14.05.2024	52°27.358'N, 13°18.437'E	F
BGB., house G	200	plant cultivation	17	50	as BGB.	27.11.2023, 14.05.2024	52°27.365'N, 13°18.425'E	G
BGB., house H/I	570	plant cultivation	12	45	as BGB.	29.12.2023, 14.05.2024	52°27.395'N, 13°18.395'E	H/I
BGB., house K	200	plant cultivation	6	60	as BGB.	29.12.2023, 14.05.2024	52°27.408'N, 13°18.378'E	K
BGB., house L	135	plant cultivation	10	50	as BGB.	29.12.2023, 14.05.2024	52°27.413'N, 13°18.395'E	L
BGB., house M	500	plant cultivation	10	45	as BGB.	29.12.2023, 14.05.2024	52°27.422'N, 13°18.413'E	М
BGB., house N	170	plant cultivation	8	60	as BGB.	29.12.2023, 14.05.2024	52°27.413'N, 13°18.422'E	Ν
BGB. Cultivation house GW02-Blü	275	plant cultivation	19	50	as BGB.	09.01.2024	52°27.315'N, 13°18.485'E	0
BGB. cultivation house 52°27.147'N, 13°18.677'E	220	plant cultivation	18	60	as BGB.	09.01.2024	52°27.147'N, 13°18.676'E	Р
Biosphere Potsdam (BP.)	5000	show greenhouse	19	94	Biological pest control: Cryptolaemus montrouzieri, Ichneumonidae, Nematoda, Chalcidoidea Insecticides: Neudorff Loxiran Ameisenköderdosen: https://www. neudorff.de/produkte/loxiran-ameisenkoederdose.html https://www.unkrautvernichter-shop.de/IMIDASECT- Schabengel.html https://www.unkrautvernichter-shop. de/imidasect-ants.html	16.01.2024, 17.01.24	52°25.112'N, 13°02.917'E	/
Botanical Garden Potsdam (BGP.), Palmenhaus	300	plant cultivation	14	60	Biological pest control: multiple species Insecticides: Spruzit Pyritrin	30.01.2024	52°24.219'N, 13°01.563'E	РН
BGP., Epiphytenhaus	220	plant cultivation	19	100	as BGP.	31.01.2024	52°24.219'N, 13°01.563'E	EH
BGP. Nutzpflanzenhaus	225	plant cultivation	18	70	as BGP.	01.02.2024	52°24.213'N, 13°01.540'E	NH
BGP., Haus der tropischen Vielfalt	225	plant cultivation	18	80	as BGP.	02.02.2024	52°24.215'N, 13°01.527'E	HTV
BGP., Kakteenhaus	190	plant cultivation	10	50	as BGP.	03.02.2024	52°24.215'N, 13°01.515'E	KH
BGP., Victoriahaus	315	plant cultivation	19	80	as BGP.	04.02.2024	52°24.227'N, 13°01.533'E	VH
BGP., Farnhaus	220	plant cultivation	16	80	as BGP.	05.02.2024	52°24'13.5"N, 13°01'33.2"E	FH

Table 1. Climatic units (see Material and methods for abbreviations).

Table 2. Ant records (see Material and methods for abbreviations).

Species	Recorded from	New record(s)	Native range	Invasiveness (ranked by antwiki.org)	Identification based on
Lasius emarginatus (Olivier, 1792)	BGB. (A, E, H/I, K, L, M, N), BGP. (VH, FH)	/	Native	/	Seifert (2018), barcoding ,iNat RG iNat AI
Lasius niger (Linnaeus, 1758)	BGB. (A, E, H/I, K, L, M, N), BGP. (PH, FH), FEZ.	/	Native	/	Seifert (2018), iNat AI
Ponera coarctata (Latreille, 1802)	BGB. (K)	/	Native	/	Seifert (2018), barcoding, iNat RG, iNat AI
Solenopsis fugax (Latreille, 1798)	BGP. (PH)	/	Native	/	Seifert (pers. com.), barcoding, Seifert (2018), iNat RG, iNat AI
Tetramorium caespitum complex	BGP. (PH, KH), FEZ.	/	Native	/	Seifert (2018), iNat RG, iNat AI
Hypoponera ergatandria (Forel, 1893)	BGP. (PH, VH), BP.	/	unknown	highly invasive (cosmopolitan)	Seifert (2018), barcoding
Plagiolepis alluaudi Emery, 1894	BGB. (A, B, C, D, E, F, G, O, P), BGP. (EH)	Berlin & Brandenburg	Africa (Wetterer 2014)	highly invasive (cosmopolitan)	Wetterer (2014), barcoding, iNat RG, iNat AI
Solenopsis texana Emery 1895	BP., GW.	Germany	the americans		Seifert (pers. com.), barcoding
Technomyrmex difficilis Forel, 1892	BP.	Germany	Madagascan region (Wetterer 2013)	highly invasive (cosmopolitan)	Seifert (pers. com.), Bolton (2007), barcoding, iNat RG
Technomyrmex vitiensis Mann, 1921	BGB. (A, B, C, D, O), BGP. (PH, EH, NH, HTV, VH, FH)	Berlin & Brandenburg	Southeast Asian region (Bolton 2007)	highly invasive (cosmopolitan)	Seifert (pers. com.), Bolton (2007), barcoding, iNat RG
Tetramorium bicarinatum (Nylander, 1846)	BGB. (A, B, C)	/	Oriental region (Wetterer 2009)	highly invasive (cosmopolitan)	Garcia and Fisher (2011), barcoding, iNat RG, iNat AI

Abbreviations

BGB: Botanical Garden Berlin, **BGP**: Botanical Garden Potsdam, **BP**: Biosphere Potsdam, **GW**: Gärten der Welt, **FEZ**: FEZ. Ökoinsel greenhouse.

Results

A total of 32 introduced species were found in the five institutions. Insects are the largest taxonomic group with 14 species, including six species of ants, one species of termites, two cockroaches, two Orthoptera, one earwig, one weevil and one drainfly species. Seven species belong to spiders and, except for one species of *Schitzomida*, all belong to the order Araneae. There were also six species of snails, two species of millipedes, one species of isopods, one species of amphipods, one species of terrestrial planaria and one species of terrestrial nemertine (Tables 2, 3).

From these, we report eight species for the first time at the regional, national or continental scale, i.e. we found no published or unpublished records:

- *Technomyrmex difficilis* (Hymenoptera, Formicidae) is reported the first time for Germany.
- One of two *Solenopsis* (Hymenoptera, Formicidae) species found by us should be *Solenopsis texana*. This species is recorded for the first time from Germany.
- *Plagiolepis alluaudi* (Hymenoptera, Formicidae) is new for Berlin and Brandenburg.
- *Technomyrmex vitiensis* (Hymenoptera, Formicidae) is new for Berlin and Brandenburg.
- *Alepia* cf. *viatrix* (Diptera, Psychodidae) is a new record for continental Eurasia.

- *Cryptotermes cavifrons* (Blattodea, Kalotermitidae) is new for Europe.
- Geonemertes pelaensis (Hoplonemertea, Prosorhochmidae) is new for Europe.
- One so-far unidentified species of the genus Anisorhynchodemus (Tricladida, Geoplanidae) is first recorded for Germany.

iNaturalist (based on AI and/or community identifications) correctly identified 31 out of a total of 37 native and non-native species (83.8%). Of the eight new records, iNaturalist was able to identify five species. Twenty of the species were successfully barcoded. The attempt to barcode three additional species failed and the rest of the species were only photographed. In 13 cases, specialist literature was used for morphological identification. Of the total 37 species, 14 species were identified by experts (Tables 2, 3).

Correlation of minimal greenhouse temperatures with ant species numbers supports that the number of introduced species is dependent on high minimum temperatures, with the Pearson-correlation being significantly positive (r = 0.856; p < 0.01). For the native species, the correlation was negative, but not significant (r = -0.581; p = 0.078). Thus, native species occurred at almost all minimum temperatures (nine of the ten possible), while introduced species were not found in climatic units with a minimum temperature of 13 °C or less (Fig. 2). This temperature dependence was observed quite well in the BGB. The ants can move between climatic units there because the climatic units are not physically isolated. In climatic unit A, the minimum temperature was the highest with 19 °C. Tetramorium bicarinatum, Technomyrmex vitiensis and Plagiolepis alluaudi were abundant. In the adjoining two

Class	Order	Species	Recorded for	New record(s)	Native range	Identification based on
Insecta	Orthoptera	Gryllodes sigillatus/supplicans	BGB.	/	South-western Asia (www.speciesfile.org 2024)	barcoding, iNat RG, iNat AI
Insecta	Orthoptera	Tachycines asynamorus (Adelung, 1902)	FEZ., GW.	/	East Asia (Boettger 1950)	barcoding, iNat RG, iNat AI
Insecta	Blattodea	Cryptotermes cavifrons Banks, 1906	BP.	Europe	Florida (Brammer and Scheffrahn 2002)	barcoding, Scheffrahn and Krecek (1999); Scheffrahn and Su (1994)
Insecta	Blattodea	Periplaneta australasiae (Fabricius, 1775)	BGB., BGP., BP.	/	Africa (Rehn 1945)	barcoding, iNat RG, iNat AI
Insecta	Blattodea	Pycnoscelus surinamensis (Linnaeus, 1758)	BP.	/	Indo-Malaysia (Gade and Parker 1997)	barcoding, iNat RG, iNat AI
Insecta	Dermaptera	Euborellia arcanum Matzke & Kočárek, 2015	BP.	/	unknown	barcoding, iNat RG, iNat AI
Insecta	Coleoptera	Sitophilus oryzae A.Hustache, 1930	BP.	/	likely Indian subcontinent (Corrêa et al. 2017)	barcoding, iNat AI
Insecta	Diptera	<i>Alepia</i> cf. <i>viatrix</i> Jaume-Schinkel, Kvifte, Weele & Mengual, 2022	BGP.	Continental Eurasia	Neotropics (Jaume- Schinkel et al. 2023)	barcoding
Arachnida	Araneae	Amaurobius ferox (Walckenaer, 1830)	BGB.	/	Mediterranean region (Bellmann 2001)	iNat RG, iNat AI
Arachnida	Araneae	Hasarius adansoni (Audouin, 1826)	BGB., BGP., BP., GW.	/	Africa (World Spider Catalogue 2017)	iNat RG, iNat AI
Arachnida	Araneae	Holocnemus pluchei (Scopoli, 1763)	BGP.	/	Mediterranean region (World Spider Catalogue 2016)	iNat RG
Arachnida	Araneae	Parasteatoda tepidariorum (C.L.Koch, 1841)	BGB., BGP., BP.	/	unknown	iNat RG, iNat AI
Arachnida	Araneae	Pholcus phalangioides (Fuesslin, 1775)	BP.	/	Western Asia (World Spider Catalog 2024)	iNat RG, iNat AI
Arachnida	Araneae	Uloborus plumipes Lucas, 1846	BGP., BP.	/	Old World (World Spider Catalog 2020)	iNat RG, iNat AI
Arachnida	Schizomida	Stenochrus portoricensis Chamberlin, 1922	BP.	/	Southeast Asia (Rowland and Reddell 1977)	barcoding, Lauterbach et al. (2020)
Diplopoda	Spirobolida	Leptogoniulus sorornus (Butler, 1876)	BP.	/	Southeast Asia (Decker et al. 2014)	Peter Decker (pers. com.), iNat RG, iNat AI
Diplopoda	Polydesmida	Oxidus gracilis (C.L.Koch, 1847)	BGB., BGP., BP.	/	East Asia (Decker et al. 2014)	barcoding, iNat RG, iNat AI
Malacostraca	Amphipoda	Talitroidea	BGP.	/		iNat: one expert opinion
Malacostraca	Isopoda	Porcellio dilatatus Brandt, 1831	BGP.	/	Southwest Europe (Allspach 1992)	Andreas Allspach (pers. com.), iNat RG, iNat AI
Gastropoda	Stylommatophora	Ambigolimax valentianus (A.Férussac, 1821)	BGB., BGP.	/	Southwest Europe (www.naturportal- suedwest.de 2024)	Carsten Renker (pers. com.), iNat RG, iNat AI
Gastropoda	Stylommatophora	Hygromia cinctella (Draparnaud, 1801)	BGB.	/	Italy and surrounding regions (Beckmann and Kobialka 2008)	Carsten Renker (pers. com.), iNat RG, iNat AI
Gastropoda	Sorbebeoconcha	Melanoides tuberculata (O.F.Müller, 1774)	BP.	/	unknown	Carsten Renker (pers. com.), iNat RG, iNat AI
Gastropoda	Pulmonata	Physella acuta (Draparnaud, 1805)	BGP.	/	North America (Semenchenko et al. 2008)	Carsten Renker (pers. com.), iNat RG, iNat AI
Gastropoda	Stylommatophora	Subulina cf. octona (Bruguière, 1789)	BGP., BP.	/	Caribbean (Deisler and Abbott 1984)	Carsten Renker (pers. com.)
Hoplonemertea	Monostilifera	Geonemertes pelaensis Semper, 1863	BP.	Europe	Indopacific region (Morffe et al. 2020)	Leigh Winsor (pers. com.), barcoding, iNat RG, iNat AI
Turbellaria	Tricladida	Anisorhynchodemus sp.	BP.	Germany	unknown	Leigh Winsor (pers. com.), iNat RG, iNat AI

Table 3. Other introduced greenhouse fauna (see Material and methods for abbreviations).

climatic units, B and C, with minimum temperatures of 18 and 17 °C, respectively, the species composition did not change at all. At 17 °C minimum, *Technomyrmex vitiensis* was more dominant than *Tetramorium bicarinatum*. In the following climatic unit D with 15 °C, *Tetramorium bicarinatum* was missing. The other two species were also much less abundant than in the warmer climatic units. In climatic unit H/I with a minimum temperature of 12 °C, none of the introduced species was found (Fig. 2).

Table 4. Genetic data of collected specimens.

Species	Sample ID	Collection Date	Recorded from	Coordinates (Lat, Lon)	Bold ID	Highest similarity	Possible species COI
Cryptotermes cavifrons	DEI-	16.01.2024	BP.	52.4185, 13.0486	GBAR001-24	99.85% to TMTER440-15	/
T	DEL CIEllan 5201	22 11 2022	DCD	52 45(4 12 207(CD 4 D002 24	(Cryptotermes cavifrons)	/
hicarinatum	DEI-GISHYIII3391	22.11.2023	DUD.	52.4504, 15.5070	GBAR002-24	(Tetramorium hicarinatum)	/
Tetramorium	DEI-GISHvm5392	22.11.2023	BGB.	52.4564, 13.3076	GBAR003-24	100% to ASANA722-06	/
bicarinatum	,			,		(Tetramorium bicarinatum)	
Stenochrus portoricensis	SMNG_24/67178/1	17.01.2024	BP.	52.4185, 13.0486	GBAR004-24	100% to GACO1420-19	/
						(Stenochrus portoricensis)	
Stenochrus portoricensis	SMNG_24/67178/2	17.01.2024	BP.	52.4185, 13.0486	GBAR005-24	100% to GACO1420-19	/
T I	DEL CIEllan, 5202	16 01 2024	DD	52 4195 12 0496	CD 4 D00(24	(Stenochrus portoricensis)	
Technomyrmex aijjicius	DEI-GISHYIII3393	10.01.2024	Dr.	52.4185, 15.0480	GBAR000-24	(Technomyrmer difficilis)	/
Sitophilus orvzae	SDEI-	16 01 2024	BP	52 4185 13 0486	GBAR007-24	100% to CICRP053-15	/
Shophilus ory240	Coleoptera304561	1010112021	211	2211100, 1010100	02111007 21	(Sitophilus oryzae)	
Technomyrmex vitiensis	DEI-GISHym5394	22.11.2023	BGB.	52.4564, 13.3076	GBAR008-24	100% to LEFIJ12092-20	/
						(Technomyrmex vitiensis)	
Technomyrmex vitiensis	DEI-GISHym5395	22.11.2023	BGB.	52.4564, 13.3076	GBAR009-24	100% to LEFIJ12092-20	/
						(Technomyrmex vitiensis)	
Plagiolepis alluaudi	DEI-GISHym5396	21.11.2023	BGB.	52.4564, 13.3076	GBAR010-24	99.84% to ASAMZ391-0/	P. alluaudi, Nylanderia
Hypopopera eractandria	DELGISHvm5397	30.01.2024	BGP	52 4038 13 0254	GBAR011-24	100% to NODRV624-15	H punctatissima
Hypoponera ergananaria	DEI GIOITYIII5577	50.01.2021	DGI.	52.1050, 15.0251	GD/HOIT 21	(Hypoponera ergatandria)	H. ergatandria
Hypoponera ergatandria	DEI-GISHym5398	30.01.2024	BGP.	52.4038, 13.0254	GBAR012-24	100% to NODRY624-15	H. punctatissima,
	-					(Hypoponera ergatandria)	H. ergatandria
Ponera coarctata	DEI-GISHym5399	29.12.2023	BGB.	52.4568, 13.3063	GBAR013-24	100% to ANTBG104-11	P. coarctata, P. testacea
						(Ponera coarctata)	
Lasius emarginatus	DEI-GISHym5400	30.01.2024	BGP.	52.4037, 13.0255	GBAR014-24	100% to NOANT006-12	L. niger, L. platythorax,
						(Lasius niger)	L. brunneus, L. paralienus,
Lasius emarginatus	DELGISHvm5401	30.01.2024	BGP	52 4037 13 0255	GBAR015-24	100% to NOANT006-12	I niger I platythoray
Lusius emarginatus	DEI-GISHIyill5401	50.01.2024	DOI:	52.4057, 15.0255	GB/11015-24	(Lasius niger)	L. brunneus, L. paralienus,
						()	L. emarginatus
Tachycines asynamorus	DEI-	14.11.2023	GW.	52.5395, 13.5777	GBAR016-24	100% to GBORT531-14	/
	Hemimetabola100425					(Tachycines asynamorus)	
Periplaneta australasiae	DEI-	22.11.2023	BGB.	52.4564, 13.3076	GBAR017-24	100% to VAQT477-09	/
D	Hemimetabola100426	16 01 2024	DD	52 4195 12 0496	CD 4 D 019 24	(Periplaneta australasiae)	/
r ycnosceius surinamensis	DEI- Hemimetabola100427	10.01.2024	DP.	52.4165, 15.0460	UDAR018-24	(Pvcnoscelus surinamensis)	1
Euborellia arcanum	DEI-	17.01.2023	BP.	52.4185, 13.0486	GBAR019-24	100% to GBMND6588-21	/
	Hemimetabola100428			,		(Euborellia arcanum)	
Gryllodes sigillatus/	DEI-	14.05.2024	BGB.	52.4565, 13.3066	GBAR020-24	99.08% to MAORT859-12	G. sigillatus, G. supplicans
supplicans	Hemimetabola100429					(Gryllodes supplicans)	
Oxidus gracilis	VNR_021924	29.12.2023	BGB.	52.4568, 13.3063	GBAR021-24	100% to MYRUS048-09	/
Coon out ou too n ol a ou aia	DEI	16 01 2024	חח	52 4195 12 0496	CDAD022.24	(Oxidus gracilis)	/
Geonemertes pelaensis	DEI- Hemimetabola100430	16.01.2024	BP.	52.4185, 15.0480	GBAR022-24	(Geonemertes pelgensis)	/
Plagiolepis alluaudi	DEI-GISHvm5402	09.01.2024	BGB.	52,4524, 13,3113	GBAR023-24	100% to ASAMZ391-07	P. alluaudi, Nvlanderia
						(Plagiolepis alluaudi)	madagascarensis
Alepia cf. viatrix	DEI-	30.01.2024	BGP.	52.4035, 13.0256	GBAR024-24	100% to ASMII2842-22	/
	Hemimetabola100431					(Alepia viatrix)	
Solenopsis texana	DEI-GISHym5403	16.01.2024	BP.	52.4185, 13.0486	GBAR025-24	100% to ASPNA1425-10	/
<u> </u>	DEL CICILI 5404	16.01.0004	DD	50 4105 10 0400	CD 4 D026 24	(Solenopsis saudiensis)	1
Solenopsis texana	DEI-GISHym5404	16.01.2024	BP.	52.4185, 13.0486	GBAR026-24	100% to ASPNA1425-10 (Solanopsis saudiansis)	/
Solenopsis texana	DEI-GISHvm5405	16.01.2024	BP.	52.4185. 13.0486	GBAR027-24	100% to ASPNA1425-10	/
borenopolo reatina	DEI OIDIIJIID IOU	1010112021	211	2211100, 1010100	0011102/21	(Solenopsis saudiensis)	
Solenopsis texana	DEI-GISHym5406	14.11.2023	GW.	52.5395, 13.5777	GBAR028-24	100% to ASPNA1425-10	/
			-			(Solenopsis saudiensis)	
Solenopsis texana	DEI-GISHym5407	14.11.2023	GW.	52.5395, 13.5777	GBAR029-24	100% to ASPNA1425-10	/
<u> </u>	DEL CIQUE 5400	14.11.0000	CIV	50 5005 10 5555	CD 4 D 020 04	(Solenopsis saudiensis)	
solenopsis texana	DEI-GISHym5408	14.11.2023	GW.	32.3395, 13.5777	GBAR030-24	100% to ASPNA1425-10	/
Technomyrmer difficilis	DFI-GISHvm5409	16 01 2024	BP	52 4185 13 0486	GBAR031-24	100% to ASANA512-06	/
	SEI GIOIIJIIO (0)	- 010112027	21.		55.11001 27	(Technomyrmex difficilis)	, ,
Solenopsis fugax	DEI-GISHym5410	30.01.2024	BGP.	52.4036, 13.0261	GBAR032-24	100% to ANTBG284-11	/
	-					(Solenopsis fugax)	
Technomyrmex vitiensis	DEI-GISHym5411	30.01.2024	BGP.	52.4036, 13.0261	GBAR033-24	100% to LEFIJ12092-20	/
<i></i>	DELCION	00.11.0000	DOD	52 4026 12 226	1	(Technomyrmex vitiensis)	1
Tetramorium caespitum	DEI-GISHym5412	22.11.2023	BGB.	52.4036, 13.0261	/	/	/
Lasius niger	DEL-GISHvm5413	22 11 2023	BGB	52 4564 13 3076	/	/	/
Leptogoniulus sorornus	VNR 021982	17.01.2024	BP.	52.4185, 13.0486	/	/	/

Discussion

We found no introduced ant species in climatic units with minimum temperatures lower than 14 °C and the number of introduced species correlates positively with higher minimum temperatures. Our study shows that minimum temperature limits the colonisation success of introduced species (Fig. 2). This dataset could be useful for modelling the potential distribution of invasive ant species under future climate change conditions. Furthermore, minimising temperatures, where possible, could be a possible measure for controlling introduced ants in greenhouses. However, further research is needed in this field.

iNaturalist provides an efficient, global method of species identification, with its AI trained on millions of images and currently covering 93,369 species. It excels at identifying common species with distinctive characters, but struggles with cryptic species such as ants. Community identifications are a good complement to AI because many biologists and professionals, together with amateur naturalists, can make a solid identification, if good photographs with the relevant features are provided. iNaturalist also connected us to international experts who helped in identifications allowing us to identify several taxa. For example, we got to know the experts Sean Birk, Bek Craig and Leigh Winsor through iNaturalist. However, taxonomic groups with small specialist communities often go unidentified. We faced this challenge with snails. While iNaturalist is a good starting point, it was not the only method used for difficult-to-identify species.

Where samples were collected, identification was possible through literature and/or COI barcoding. Both methods

ant species

Elias Freyhof & Emil Janke: Invertebrates in greenhouses

However, these methods can provide very accurate identifications in many cases. Morphology-based literature requires matching keys or descriptions, which can be difficult to find for introduced species that can come from anywhere in the world. COI barcoding is more universal, but has the problem that not every individual can be identified to species level. For some species, there are few or no reference sequences in the database, which can lead to errors. At least one of these two methods was preferred by us to obtain a reliable identification of the more difficult-to-identify species.

The next method used was expert identification, which was an additional step, if one of the other methods did not work, was not possible or was not accurate enough. This method can provide a very reliable identification without a lot of knowledge and material. However, it requires contact with experts and some time. We also asked experts when only photos were taken and iNaturalist did not provide reliable identifications.

The next part of the discussion deals with the species which, as far as we could find out, are new records at the regional or even continental scale (Tables 2, 3).

Solenopsis texana (Fig. 3.)

This species of thief ant is extremely small (about 1.2 mm) and golden yellow in colour. Its native range should be the Americas, where it has been found in Canada, the USA, Mexico, Nicaragua, Costa Rica, Panama, Colombia and Brazil. It has been introduced into greenhouses in Europe,



Figure 2. Records of native and non-native ant species in greenhouses with different minimum temperatures.



Figure 3. *Solenopsis texana* nest with workers and larvae: Gärten der Welt.

including Norway and Germany (https://www.antwiki. org; (GBIF 2023a): www.gbif.org; Seifert, pers. com.). Its hidden lifestyle in the soil and its small size makes it very difficult to detect. The distribution of this species is, therefore, probably underestimated. According to evaluation of the image of a worker type (specimen identifier CASENT0904625 in www.antbase.org), this Solenopsis species should be Solenopsis texana Emery 1895 (B. Seifert, pers. comm. 11 October 2024). To our knowledge, Solenopsis texana has never been found in Germany. The taxonomy of this species is unclear and needs to be revised (Seifert, pers. com.). Barcoding resulted in Solenopsis saudiensis (100% similarity on best matches), which is a junior synonym of S. abdita (Sharaf et al. 2020). However, not all species of the S. molesta complex are available in GenBank making it impossible to identify our materials only by COI barcode. Seifert (pers. com.) examined the morphometry of our specimens and found them to be conspecific with an Solenopsis texana found by Gjershaug et al. (2016) in a greenhouse in Norway (NOR: 58.82005°N, 9.07145°E, Kristiansand, 80 m, Dyreparken, in Tropenhaus, A.Staverlokk 2016.04.22).

Technomyrmex difficilis (Fig. 4)

This species is a 3.5–4.0 mm long, black ant native to the Madagascar region (Wetterer 2013). It has been introduced to Central and North America, where it is known as an indoor species in the north. In Europe, *T. difficilis* has been recorded for indoor habitats in France, Belgium, Sweden and, for the first time by us, Germany. It has also been introduced to the Arabian Peninsula, South Africa, Southeast Asia and Oceania (https://www.antwiki.org; (GBIF 2023a): www.gbif.org). Blatrix et al. (2018) already stated that "the actual distribution of this species is poorly understood, as it was only distinguished from *T. albipes* since 2007 (Bolton 2007)". We found this species mostly amongst plants and Bolton (2007) reported that nests were often found in trees, shrubs, in loose mulch, under debris, in leaf litter, both on and above ground.



Figure 4. *Technomyrmex difficilis* Photo from AntWeb (www. antweb.org); photographer: April Nobile.

The reproduction and caste system of *T. difficilis* is quite peculiar. After the death of the primary queen, ergatoid males and intercastes appear in the nest. The intercastes can make up almost half of the colony (Warner 2003). The ergatoid males and intercastes mate within the colony and budding may occur. Given its high polygyny and polydomy, this species is a very successful disperser, known to form super-colonies and is classified as 'highly invasive'. This species is also known to feed on and protect sap-sucking hemipterans (Warner 2003).

Technomyrmex vitiensis (Fig. 5)

This species is a 3.5–4.0 mm long, black ant, native to Southeast Asia (Bolton 2007). It has been introduced to California, French Guiana, Suriname, Brazil, the Madagascar region and Oceania. In Europe, *T. vitiensis* was found in greenhouses in the UK, France, the Netherlands, Belgium, Switzerland, Germany, Denmark, Austria, Czech Republic, Poland and Finland (https://www.antwiki.org; (GBIF 2023c): https://www.gbif.org). In fact,



Figure 5. *Technomyrmex vitiensis* Photo from AntWeb (www. antweb.org); photographer: April Nobile.

T. vitiensis is a common greenhouse species in Europe and was found at many greenhouse localities (Pospischil 2011). To our knowledge, this is the first record for Berlin and Brandenburg. *Technomyrmex vitiensis* is part of the *T. albipes* species complex and was only distinguished from *T. albipes* in 2007. In addition, the species of this complex are not easy to distinguish from each other. This also contributes to their poorly known distribution. In general, the biology of *T. vitiensis* is similar to that of *T. difficilis*.

Plagiolepis alluaudi (Fig. 6)

A very small (1.5 mm) yellow ant native to Africa (Wetterer 2014). The biology of this species has recently been compiled by Blatrix et al. (2018). From Africa, it has been introduced to the Caribbean, California, Asia and Oceania. In Europe, it was found in greenhouses in the UK, Switzerland, Ireland, Germany, Estonia, Poland, the Netherlands, Belgium, Sweden and France (https://www.antwiki.org; (GBIF 2023b): https://www.gbif.org). To our knowledge, this is the first time this species has been reported from Berlin and Brandenburg. It appears to be a relatively recent invader of greenhouses in Europe. In France, for example, there are no records prior to 1990 (Blatrix et al. 2018). Outdoors, it nests in a variety of very small spaces: under the bark of dead trees, in the holes made by wood-boring insects, under clumps of grass, to name just a few. Indoors, it nests in cavities formed between the stems, petioles or leaves and stems of plants. It prefers to nest above ground, but is often found under flower pots. Blatrix et al. (2018) stated: "The nests are opportunistic, and the ants do not usually build structures or dig burrows. Each nest contains multiple queens, which facilitates their dispersal when materials or plants are exchanged. These ants are in addition very small and pale yellow and can be hard to detect when they are present in small numbers". Plagiolepis alluaudi protects and disperses sap-feeding hemipterans and feeds on their honeydew (González-Hernández et al. 1999). This species is also polygynous and polydomous, which contribute to its status as a 'highly invasive' species (https://www.antwiki.org).



Figure 6. *Plagiolepis alluaudi* Photo from AntWeb (www.ant-web.org); photographer: April Nobile.



Figure 7. Phragmotic soldier of *Crytotermes cavifrons*: Biosphere Potsdam.

Cryptotermes cavifrons (Fig. 7)

This dry-wood termite species is native to Florida (Brammer and Scheffrahn 2002). It has been introduced to South Carolina and Nigeria (GBIF 2023d: https://www.gbif.org). To our knowledge it has never been recorded for Europe or even Eurasia. Cryptotermes cavifrons nests in dry wood and does not need a connection to the ground. We even found a colony in a wooden bench in the greenhouse. Inside the wood, the termites build a nest of interconnected tunnels. From the outside, piles of faecal pellets are the only visible sign of their presence. In C. cavifrons, 1-2% of the colony are soldiers, which have phragmotic heads to plug the nest entrances. In dry-wood termites, the workers are replaced by pseudergates. These perform normal worker tasks, but can develop into alates when needed. This species is also unique in that alates are always present in the colony and swarms occur throughout the year (Brammer and Scheffrahn 2002).

Alepia cf. *viatrix* (Fig. 8)

This species of drainflies (Diptera, Psychodidae) is expected to be native to the Neotropics as all other species of the genus Alepia occur there (Jaume-Schinkel et al. 2022). Furthermore, the larvae depend on water reservoirs inside bromeliad plants (phytotelmata), which are endemic to the Neotropics. The introduction of this species to other parts of the world seems to be connected to the bromeliad trade (Jaume-Schinkel et al. 2023). Alepia viatrix was only recently described as a new species from the Duque da Terceira botanical garden on the Azores, containing bromeliad plants (Jaume-Schinkel et al. 2022). A following record came from Brisbane in Australia, in an area where bromeliads are cultivated. It then was found in South Africa, whereby only barcodes are available (GBIF 2023e: https://www.gbif.org). Our finding of this species was also in a greenhouse part in the Botanical Garden Potsdam, which has lots of bromeliads. We identified this tiny fly by barcoding, which is not that reliable as Alepia is a large



Figure 8. Alepia cf. viatrix: Biosphere Potsdam.

genus with 57 described species. From these, barcodes are only available from *A. viatrix* and *A. valentia* on BOLD Systems. However, the barcode has a similarity of 100% with a sample from Brisbane. Thus, this is the first record of this species for continental Eurasia, but further research including morphological analyses are needed.

Geonemertes pelaensis (Fig. 9)

A species of terrestrial nemertines (Hoplonemertea, Prosorhochmidae), which is probably native to the Indopacific Region (Morffe et al. 2020). It was introduced to the USA, where it has been recorded once from a greenhouse in Denver. It also occurs outdoors in Florida, Central America, the Caribbean, western parts of South America, the Madagascar region and Japan (GBIF 2023f: https:// www.gbif.org). To our knowledge, this is the first record for Europe. Geonemertes pelaensis is a hermaphrodite, a characteristic that makes dispersal much easier (Moore et al. 2001). It extrudes its proboscis when catching prey and/or if stimulated from outside. The diet of G. pelaensis is quite enigmatic. On the Seychelles, the species is found to be a predator of small molluscs (Gerlach 1998). On the Ogasawara Islands, however, G. pelaensis is a predator of Arthropods, which has led to devastating declines in populations of Isopoda and Amphipoda on that island since the 1980s. Shinobe et al. (2017) assume that these differences in diet could be due to different cryptic species involved.

Anisorhynchodemus sp. (Fig. 10)

A species of terrestrial flatworms of which the native range is unknown. The genus has been introduced to Florida, the Caribbean, the Madagascar region, Sri Lanka, Southeast Asia and Oceania (GBIF 2023g: https://www.gbif. org). As far as we know, it has never been reported from Germany. The taxonomy within this genus is still under discussion, as incompletely described Rhynchodeminae species are placed in the collective genus *Anisorhynchodemus* (Kawakatsu et al. 2003). Land planaria expert and researcher Leigh Winsor identified this genus on iNaturalist and said: "This appears to be the invasive *Anisorhynchodemus* species found in Florida, the Caribbean and Pacific oceanic islands" (Leigh Winsor pers com.).

The many new records for Berlin, Brandenburg, Germany and even at the continental scale demonstrate how poorly the faunas of greenhouses are known. While most of the tropical species recorded are unlikely to survive in the wild in Germany, they might be transported from our greenhouses to others, potentially reaching suitable climatic conditions along this road. It is not the aim of this study to speculate about the effect of invasive species. However, our findings highlight the immense knowledge gap of the greenhouse invertebrates. Furthermore, this study also shows how many different introduced taxonomic groups can be identified by a combination of the iNaturalist AI, its expert community and barcodes available from BOLD. Both open-access resources enabled the identification of most, but not all, of our species. Obviously, as biodiversity remains imperfectly described, many species complexes are awaiting their taxonomic revisions, making it actually impossible to identify some to the species level.

Indeed, we highly encourage other students, citizen scientists and experts to study greenhouse faunas as well as to publish their records. During our work, we came into contact with several citizen scientists, who told us that they have already seen particular species in greenhous-



Figure 9. Geonemertes pelaensis: Biosphere Potsdam.



Figure 10. Anisorhynchodemus sp.: Biosphere Potsdam.

es, but never found it worthwhile to publish these data. Nowadays, it is quite simple to publish single-species records through iNaturalist and other smartphone-based identification tools. To encourage citizen scientists to record greenhouse species, we set up an iNaturalist Project called "Greenhouse fauna of Europe".

Acknowledgements

We thank the staff of the greenhouses who allowed us to collect, to take photos of the greenhouse fauna, gave us information about the greenhouses or even sent us ant samples: BGB (Henrike Wilke); BGP (Ingo Kallmeyer, Michael Burkart); BP (Rosita Meyer, Kristina Hasselmann); GW (Peter Moede); FEZ (Beate Butz). We would also like to thank the team and some students of the Deutsches Entomologisches Institut Senckenberg. Many thanks to Eva Kleibusch, who taught us the steps of barcoding and other genetic tasks. Lara-Sophie Dey complemented this study with her genetic expertise. Thanks also to Stephan Blank for solving all sorts of problems. Christian Kutzscher showed how to prepare ants in a sensible way. Arne Köhler showed how to create and edit figures. Emilia Kramm helped a lot with the lab work, technical issues and the map. Nura Alaya helped to collect a missing sample. We are also grateful to the experts who helped with species identification: ants (Bernhard Seifert, Senckenberg Natural History Museum Görlitz (Germany)); snails (Carsten Renker, Mainz Natural History Museum); centipedes (Peter Decker, Senckenberg Natural History Museum Görlitz (Germany)); land planarians and terrestrial nemertines (Leigh Winsor, James Cook University (Australia); Schitzomida (Sean Birk Bek Craig Aarhus University (Denmark)); Isopoda (Andreas Allspach, Senckenberg Research Institute and Natural History Museum Frankfurt (Germany)).

References

Allspach A (1992) Die Landasseln Hessens. Naturschutz Heute 12: 1–6. AntWeb (2024) Version 8.108. California Academy of Science. [Online at] https://www.antweb.org [accessed on 2024-09-08]

- Astrin JJ, Stüben PE (2008) Phylogeny in cryptic weevils: molecules, morphology and new genera of western Palaearctic Cryptorhynchinae (Coleoptera: Curculionidae). Invertebrate Systematics 22(5): 503–522. https://doi.org/10.1071/IS07057
- Astrin JJ, Höfer H, Spelda J, Holstein J, Bayer S, Hendrich L, Bernhard AH, Kielhorn K-H, Krammer H-J, Lemke M, Monje JC, Morinière J, Rulik B, Petersen M, Janssen H, Muster C (2016) Towards a DNA barcode reference database for spiders and harvestmen of Germany. PLOS ONE 11(9): e0162624. https://doi.org/10.1371/journal.pone.0162624
- Báthori F, Herczeg G, Vilizzi L, Jégh T, Kakas C, Petrovics M, Csősz S (2024) A survey and risk screening of non-native ant species colonising greenhouses in Hungary. Biological Invasions 26(4): 1033– 1044. https://doi.org/10.1007/s10530-023-03227-9

- Beckmann KH, Kobialka H (2008) Hygromia cinctella (Draparnaud, 1801) auf dem Eroberungszug durch Deutschland (Gastropoda: Hygromiidae). Club Conchylia Informationen 39(1/2): 34–41.
- Bellmann H (2001) Kosmos-Atlas Spinnentiere Europas. Franckh-Kosmos Verlags-GmbH & Co., Stuttgart, 304 pp.
- Blatrix R, Colin T, Wegnez P, Galkowski C, Geniez P (2018) Introduced ants (Hymenoptera: Formicidae) of mainland France and Belgium, with a focus on greenhouses. Annales de la Société Entomologique de France (NS) 54(4): 293–308. https://doi.org/10.1080/00379271. 2018.1490927
- Boettger CR (1950) Die Gewächshausheuschrecke (*Tachycines asyn-amorus* Adelung). Abhandlungen der Braunschweigischen Wissenschaftlichen Gesellschaft 2: 13–39.
- Boer P, Vierbergen B (2008) Exotic ants in the Netherlands (Hymenoptera: Formicidae). Entomologische Berichten 68(4): 121–129.
- Brammer AS, Scheffrahn RH (2002) Drywood Termite, Cryptotermes cavifrons Banks (Insecta: Isoptera: Kalotermitidae). University of Florida: EENY 279. https://doi.org/10.32473/edis-in236-1999
- Bolton B (2007) Taxonomy of the dolichoderine ant genus *Technomyrmex* Mayr (Hymenoptera: Formicidae) based on the worker caste. Contributions of the American Entomological Institute 35: 1–150.
- Corrêa AS, Vinson CC, Braga LS, Guedes RNC, De Oliveira LO (2017) Ancient origin and recent range expansion of the maize weevil *Sitophilus zeamais*, and its genealogical relationship to the rice weevil *S. oryzae*. Bulletin of Entomological Research 107(1): 9–20. https://doi.org/10.1017/S0007485316000687
- Decker MP, Reip HS, Voigtländer K (2014) Millipedes and centipedes in German greenhouses (Myriapoda: Diplopoda, Chilopoda). Biodiversity Data Journal 2: e1066. https://doi.org/10.3897/ BDJ.2.e1066
- Deisler JA, Abbott RT (1984) Range extensions of some introduced land mollusks in the Bahama Islands, with first reports for four species. The Nautilus (Philadelphia, PA) 98(1): 12–17.
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. Molecular Marine Biology and Biotechnology 3: 294–299.
- Gade B, Parker Jr ED (1997) The effect of life cycle stage and genotype on desiccation tolerance in the colonizing parthenogenetic cockroach *Pycnoscelus surinamensis* and its sexual ancestor *P. indicus*. Journal of Evolutionary Biology 10(4): 479–493. https://doi.org/10.1046/ j.1420-9101.1997.10040479.x
- GBIF (2023a) GBIF Backbone Taxonomy. Checklist dataset: Technomyrmex difficilis Forel, 1892. https://doi.org/10.15468/39omei [Accessed via GBIF.org on 2024-09-11]
- GBIF (2023b) GBIF Backbone Taxonomy. Checklist dataset: *Pla-giolepis alluaudi* Emery, 1894. https://doi.org/10.15468/390mei [Accessed via GBIF.org on 2024-09-11]
- GBIF (2023c) GBIF Backbone Taxonomy. Checklist dataset: *Technomyrmex vitiensis* Mann, 1921. https://doi.org/10.15468/39omei [Accessed via GBIF.org on 2024-09-11-09-11]
- GBIF (2023d) GBIF Backbone Taxonomy. Checklist dataset: Cryptotermes cavifrons Banks, 1906. https://doi.org/10.15468/39omei [Accessed via GBIF.org on 2024-09-11]
- GBIF (2023e) GBIF Backbone Taxonomy. Checklist dataset: Alepia viatrix Jaume-Schinkel, Kvifte, Weele & Mengual, 2022. https:// doi.org/10.15468/390mei [Accessed via GBIF.org on 2024-09-11]

- GBIF (2023f) GBIF Backbone Taxonomy. Checklist dataset: Geonemertes pelaensis Semper, 1863. https://doi.org/10.15468/39omei [Accessed via GBIF.org on 2024-09-11]
- GBIF (2023g) GBIF Backbone Taxonomy. Checklist dataset: Anisorhynchodemus Kawakatsu, Froehlich, Jones, Ogren & Sasaki, 2003. https://doi.org/10.15468/39omei [Accessed via GBIF.org on 2024-09-11]
- GISD (2024) Global Invasive Species Database (2024). https://www. iucngisd.org/gisd/ [Accessed on 2024-11-10]
- Gjershaug JO, Staverløkk A, Ødegaard F (2016) Funn av fremmede maurarter i Norge i 2015.
- González-Hernández H, Johnson MW, Reimer NJ (1999) Impact of *Pheidole megacephala* (F.) (Hymenoptera: Formicidae) on the biological control of *Dysmicoccus brevipes* (Cockerell) (Homoptera: Pseudococcidae). Biological Control 15(2): 145–152. https://doi. org/10.1006/bcon.1999.0714
- Hamer MT, Turner CR (2024) Solenopsis abdita Thompson (Formicidae: Myrmicinae) new to Britain. Entomologist's Monthly Magazine 160(2): 144–150. https://doi.org/10.31184/M00138908.1602.4231
- Hobbs RJ, Higgs E, Harris JA (2009) Novel ecosystems: implications for conservation and restoration. Trends in Ecology & Evolution 24(11): 599–605. https://doi.org/10.1016/j.tree.2009.05.012
- iNaturalist (2024) iNaturalist. https://www.inaturalist.org [Accessed on 2024-11-9]
- IPBES (2023) Summary for Policymakers of the Thematic Assessment Report on Invasive Alien Species and their Control of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. Roy HE, Pauchard A, Stoett P, Renard Truong T, Bacher S, Galil BS, Hulme PE, Ikeda T, Sankaran KV, McGeoch MA, Meyerson LA, Nuñez MA, Ordonez A, Rahlao SJ, Schwindt E, Seebens H, Sheppard AW, Vandvik V (Eds). IPBES Secretariat, Bonn, Germany.
- Jaume-Schinkel S, Kvifte GM, Mengual X (2022) Alepia viatrix sp. nov. (Diptera: Psychodidae), a new species of a Neotropical genus found on the Azores Archipelago (Portugal). Zootaxa 5128(3): 384– 396. https://doi.org/10.11646/zootaxa.5128.3.4
- Jaume-Schinkel S, Mengual X, Howe AG, Fagan-Jeffries EP (2023) The hitchhiker's guide to Australia: the 18,000-km-long journey of *Alepia viatrix* Jaume-Schinkel, Kvifte, Weele & Mengual, 2022 (Diptera, Psychodidae) discovered through citizen science. Check List 19(4): 589–597. https://doi.org/10.15560/19.4.589
- Kawakatsu M, Ogren RE, Froehlich EM, Sasaki GY (2002) Additions and corrections of the previous land planarian indices of the world (Turbellaria, Seriata, Tricladida, Terricola). Bulletin of the Fuji Women's College (Series 2) 40: 157–177.
- Kenis M, Branco M (2010) Impact of alien terrestrial arthropods in Europe. Chapter 5. BioRisk 4: 51–71. https://doi.org/10.3897/biorisk.4.42
- Lauterbach S, Hörren T, Bauer T (2020) Stenochrus portoricensis neu für Nordrhein-Westfalen, mit Anmerkungen zur Verbreitung und Habitaten weiterer eingeschleppter Zwerggeißelskorpionarten in Europa (Arachnida: Schizomida). Arachnologische Mitteilungen: Arachnology Letters 60(1): 50–54. https://doi.org/10.30963/aramit6010
- Lowe S, Browne M, Boudjelas S, De Poorter M (2000) 100 of the world's worst invasive alien species: a selection from the global invasive species database. Volume 12. Auckland: Invasive Species Specialist Group.

- Mehta SV, Haight RG, Homans FR, Polasky S, Venette RC (2007) Optimal detection and control strategies for invasive species management. Ecological Economics 61(2–3): 237–245. https://doi. org/10.1016/j.ecolecon.2006.10.024
- Morffe J, Garcia N, Breugelmans K (2020) First record of the terrestrial nemertean *Geonemertes pelaensis* Semper, 1863 (Hoplonemertea: Prosorhochmidae) for Cuba. BioInvasions Records 9(2): 399–402. https://doi.org/10.3391/bir.2020.9.2.26
- Naturportal Südwest (2024) Naturportal Südwest: Ambigolimax valentianus (A. Férussac, 1822) Gewächshausschnegel. https://naturportal-suedwest.de/de/weichtiere/systematik/art/nacktschnecken-gastropoda/limacidae-schnegel/ambigolimax/valentianus/ [Accessed on 2024-20-08]
- Orthoptera Species File (2024) Orthoptera Species File: Species *Gryllodes sigillatus* (Walker, 1869). https://orthoptera.archive. speciesfile.org/Common/basic/Taxa.aspx?TaxonNameID=1123306 [Accessed on 2024-08-20]
- Pospischil R (2011) Role of tropical greenhouses for introduction and establishment of foreign ant species (Hymenoptera: Formicidae) in Central Europe. In: Robinson WH, de Carvalho Campos AE (Eds) Proceedings of the 7th International Conference on Urban Pests, Ouro Preto, Brazil, 7–10, 59–66.
- Ratnasingham S, Hebert PD (2007) BOLD: The Barcode of Life Data System (http://www.barcodinglife.org). Molecular Ecology Notes 7(3): 355–364. https://doi.org/10.1111/j.1471-8286.2007.01678.x
- Rehn JA (1945) Man's uninvited fellow traveler—the cockroach. The Scientific Monthly: 265–276.
- Rowland JM, Reddell JR (1977) A review of the cavernicole Schizomida (Arachnida) of Mexico, Guatemala, and Belize. Association for Mexican Cave Studies Bulletin 6: 79–102.
- Scheffrahn RH, Su NY (1994) Keys to soldier and winged adult termites (Isoptera) of Florida. Florida Entomologist 77(4): 460–474. https:// doi.org/10.2307/3495700
- Scheffrahn RH, Krecek J (1999) Termites of the genus Cryptotermes Banks (Isoptera: Kalotermitidae) from the West Indies. Insecta Mundi 13: 111–117.
- Seifert B (2013) Hypoponera ergatandria (Forel, 1893)—a cosmopolitan tramp species different from H. punctatissima (Roger, 1859) (Hymenoptera: Formicidae). Soil Organisms 85(3): 189–201. https://www.senckenberg.de/wp-content/uploads/2019/08/05_artikel bernhard seifert.pdf
- Seifert B (2018) The ants of central and north Europe. Lutra Verlags-und Vertriebsgesellschaft.
- Semenchenko V, Laenko T, Razlutskij V (2008) A new record of the North American gastropod *Physella acuta* (Draparnaud 1805) from the Neman River Basin, Belarus. Aquatic Invasions 3(3): 359–360. https://doi.org/10.3391/ai.2008.3.3.14
- Sharaf MR, Gotzek D, Guénard B, Fisher BL, Aldawood AS, Al Dhafer HM, Mohamed AA (2020) Molecular phylogenetic analysis and morphological reassessments of thief ants identify a new potential case of biological invasions. Scientific Reports 10(1): 12040. https:// doi.org/10.1038/s41598-020-69029-4
- Shinobe S, Uchida S, Mori H, Okochi I, Chiba S (2017) Declining soil Crustacea in a World Heritage Site caused by land nemertean. Scientific Reports 7(1): 12400. https://doi.org/10.1038/s41598-017-12653-4

- Stohlgren TJ, Schnase JL (2006) Risk analysis for biological hazards: what we need to know about invasive species. Risk Analysis: An International Journal 26(1): 163–173. https://doi.org/10.1111/j.1539-6924.2006.00707.x
- Suvák M (2011) Predatory and parasitic insects in greenhouses of Botanical Garden of PJ Šafárik University in Košice, Slovakia. Thaiszia–Journal of Botany, Košice 21: 185–205.
- Tsuji K, Furukawa T, Kinomura K, Takamine H, Yamauchi K (1991) The caste system of the dolichoderine ant *Technomyrmex albipes* (Hymenoptera: Formicidae): morphological description of queens, workers and reproductively active intercastes. Insectes Sociaux 38: 413–422. https://doi.org/10.1007/BF01241875
- Tsutsui ND, Suarez AV (2003) The colony structure and population biology of invasive ants. Conservation Biology 17(1): 48–58. https://doi.org/10.1046/j.1523-1739.2003.02018.x
- Walsh PS, Metzger DA, Higuchi R (1991) Chelex 100 as a medium for simple extraction of DNA for PCR-based typing from forensic material. Biotechniques 10(4): 506–513. https://doi. org/10.2144/000114018
- Wang C, Zhang X, Pan X, Li Z, Zhu S (2015) Greenhouses: hotspots in the invasive network for alien species. Biodiversity and Conservation 24: 1825–1829. https://doi.org/10.1007/s10531-015-0876-x

- Warner JR (2003) Bait preferences and toxicity of insecticides to white-footed ants *Technomyrmex albipes* (Hymenoptera: Formicidae). Doctoral dissertation, University of Florida.
- Westphal MI, Browne M, MacKinnon K, Noble I (2008) The link between international trade and the global distribution of invasive alien species. Biological Invasions 10: 391–398. https://doi.org/10.1007/ s10530-007-9138-5
- Wetterer JK (2009) Worldwide spread of the penny ant, *Tetramorium bicarinatum* (Hymenoptera: Formicidae). Sociobiology 54(3): 811.
- Wetterer JK (2013) Worldwide spread of the difficult white-footed ant, *Technomyrmex difficilis* (Hymenoptera: Formicidae). Myrmecological News 18: 93–97. https://doi.org/10.1653/024.097.0231
- Wetterer JK (2014) Worldwide spread of Alluaud's little yellow ant, *Plagiolepis alluaudi* (Hymenoptera: Formicidae). Myrmecological News 19: 53–59.
- World Spider Catalog (2024) World Spider Catalog: Version 25.0. Natural History Museum Bern. Online at: http://wsc.nmbe.ch. https://doi.org/10.24436/2 [accessed on 2024-07-15]
- Yamauchi K, Furukawa T, Kinomura K, Takamine H, Tsuji K (1991) Secondary polygyny by inbred wingless sexuals in the dolichoderine ant *Technomyrmex albipes*. Behavioral Ecology and Sociobiology 29: 313–319. https://doi.org/10.1046/j.1523-1739.2003.02018.x

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Beiträge zur Entomologie = Contributions to Entomology

Jahr/Year: 2024

Band/Volume: 74

Autor(en)/Author(s): Freyhof Elias, Janke Emil

Artikel/Article: Introduced greenhouse-invertebrates in Potsdam and Berlin with a focus on ants (Hymenoptera, Formicidae) with eight new records for Europe, Germany or the Berlin-Brandenburg region 235-248