

Literatur:

- BELKA, Z. (1991): Potential conodont levels for intercontinental division of the Lower Carboniferous. – Cour. Forsch.-Inst. Senckenberg, **130**: 11-14.
- CONRAD, J. (1984) : Les séries Carbonifères du Sahara Central Algérien. Stratigraphie, sédimentation, évolution structurelle. – Thèse Doct. Etat ès-Sci. nat., Univ. Droit, d'Econ. Sci. d'Aix Marseille, 370 S.
- DEVUYST, F.-X., HANCE, L., HOU, H., WU, X., TIAN, S., COEN, M. & SEVASTOPULO, G. (2003): A proposed Global Stratotype Section and Point for the base of the Viséan Stage (Carboniferous): the Pengchong section, Guangxi, South China. – Episodes, **26** (2): 105-115.
- GORDON, M. jr. & MASON, C. E. (1985): Progradation of the Borden Formation in Kentucky, U.S.A., demonstrated by successive early Mississippian (Osagean) ammonoid faunas. – C. R. 10th Congr. Internat. Strat. Carb., Madrid 1983, **I**: 191-198.
- KORN, D., BOCKWINKEL, J., EBBIGHAUSEN, V. & KLUG, C. (2003): Palaeobiogeographic and evolutionary meaning of an early Late Tournaisian ammonoid fauna from the Tafilalt of Morocco. – Acta Palaeont. Pol., **48** (1): 71-92.
- LANE, R. H. & BRECKLE, P. L. (2001): Type Mississippian subdivisions and biostratigraphic succession. – In: HECKEL, P. H. (Ed.), Stratigraphy and Biostratigraphy of the Mississippian Subsystem (Carboniferous System) in its Type Region, the Mississippi River Valley of Illinois, Missouri, and Iowa, I.U.G.S. Subcommission on Carboniferous Stratigraphy, Guidebook for Field Conference, September 8-13, 2001, S. 93-108.
- RAMSBOTTOM, W. H. C. & SAUNDERS, W. B. (1985): Evolution and evolutionary biostratigraphy of Carboniferous ammonoids. – J. Paleont., **59** (1): 123-139.
- RILEY, N. J. (1991): A global review of mid-Dinantian ammonoid biostratigraphy. – Cour. Forsch.-Inst. Senckenberg, **130**: 133-143.
- SCHMIDT, H. (1925): Die carbonischen Goniatiten Deutschlands. – Jb. preuß. geol. Landesanst., **45**: 489-609.
- VOGES, A. 1959. Conodonten aus dem Unterkarbon desw Sauerlandes (*Gattendorfia*- und *Pericyclus*-Stufe). – Paläont. Z., **33**: 266-314.
- WEBSTER, G. D. & GROESSENS, E. (1991): Conodont subdivision of the Lower Carboniferous. – Cour. Forsch.-Inst. Senckenberg, **130**: 31-40.
- WEYER, D. (1972): Trilobiten und Ammonoideen aus der *Entogonites nasutus*-Zone (Unterkarbon) des Büchenberg-Sattels (Elbingeröder Komplex, Harz), Teil I. – Geologie, **21**: 166-184.

A NEW METHOD FOR QUALITATIVE ESTIMATION OF PRECIPITATION USING FOSSIL AMPHIBIANS AND REPTILES

Madelaine BÖHME¹ & August ILG²

¹ Department of Earth and Environmental Sciences, Section Palaeontology, Ludwig-Maximilians-University Munich, Richard-Wagner-Str. 10, D-80333 Munich

² Schumannstrasse 83, D-4023 Düsseldorf; e-mail: august.ilg@t-online.de

Precipitation is an important geodynamic control factor coupled to tectonics, erosion, continental run-off and weathering, and to oceanic circulation. But in practice this climate parameter is difficult to estimate and long term and high resolution continental records are mostly lacking. Here we introduce a new palaeobiological method of quantitative estimation of paleo-precipitation based on indexing of ecophysiological groups within herpetological communities. In recent communities this groups show a highly significant correlation to the annual precipitation. We apply this correlation to a high resolution ~11 million year continental sequence of the Calatayud-Daroca and Teruel-Alfambra sections (Northern Spain) in the western Mediterranean and to several Neogene North-South transects. The results show that the new method will be a powerful tool to reconstruct past temporally and spatially precipitation patterns and will therefore contribute to a better understanding and modelling of geodynamic processes. It complements palaeobotanical methods, because it is applicable on a different set of sedimentary and taphonomic facies.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte des Institutes für Geologie und Paläontologie der Karl-Franzens-Universität Graz](#)

Jahr/Year: 2005

Band/Volume: [10](#)

Autor(en)/Author(s): Böhme Madelaine, Ilg August

Artikel/Article: [A new method for qualitative estimation of precipitation using fossil amphibians and reptiles. 9](#)