

GEOMETRIC MORPHOMETRICS –

A contribution to the study of shape variability in Ostracods

Angel Baltanás

Department Ecología, Universidad Autónoma de Madrid (Edif. Biología), E-28049 Madrid
(E-Mail: angel.baltanas@uam.es).

We have told each other so often and with such force and such eloquence of the uses to which the study of ostracodes has been applied that we have overlooked one startling fact: almost no one uses ostracodes for anything.
R.L. Kaesler (1983)

Why shape?

Biodiversity is an issue of main concern not only for scientists but for the whole society as well. Taxonomic richness is but one of the many ways we use to express biological diversity. Morphological disparity – the amount of shape variability within a clade – is another. And given that both features do not necessarily correlate (Cherry et al. 1979, Foote 1993a), their comparison can provide further evidence about ecological and evolutionary processes involved in the production and maintenance of biodiversity.

Morphological disparity can be explored at a wide range of taxonomic levels. Indeed, there has been a growing interest in methods addressing morphological disparity in recent years (Foote 1997, McGhee 1999, Ciampaglio et al. 2001, Wills 2001, Zelditch et al. 2004). Some ‘classic’ studies concern the morphospace occupied by spiral (Raup 1967) or planar branch systems (McKinney 1981) aiming to understand macroevolutionary patterns at high-rank taxonomic levels. Studies in ecomorphology, however, commonly focus at lower taxonomic levels (mainly, closely related species) seeking for correlations between morphology and ecological requirements (Norberg 1994) or for the effects of competitive selective pressures assumed to occur between (Dayan et al. 1990). At the species level, between-populations disparity and its correlation with environmental conditions is used to evaluate adaptation to local conditions (Loik and Noble 1993); and at further detail, morphological variability within a population can be related to the niche concept and dynamics (Pulliam 1986) or to sexual selection (Møller 1994).

Finally, there is an increasing interest in exploring the potential for shape change of a given genotype – the phenotypic plasticity of morphological features –, as well as its adaptive value (Schlichting and Pigliucci 1998).

The amount of morphospace occupied by a set of clades has been used as indicator of ecological diversity (Warheit et al. 1999), evolutionary radiation (McGhee 1999), morphological convergence in distant communities (Ricklefs and Miles 1994), or selective extinctions (Roy and Foote 1997). Less frequent is its use as tracer of environmental conditions or dispersal routes of groups below the species level (populations, clones, ...).

Why Ostracods?

Most of the issues outlined above can be extensively addressed using ostracods. Indeed, this group of organisms can be labelled as ideal for a morphometric approach because of its high taxonomic richness and the diversity of habitats occupied. In addition, ostracods have an extensive fossil record, a feature that allows the examination of shape-environment relationships back into evolutionary time scale.

Morphometric study of ostracods mainly focuses on the analysis of carapace shape. Ostracod carapace has a marked functional meaning; it is the interface between the organism and its environment (Benson 1981). Hence, ostracod carapaces can be considered as engineering solutions, a compromise between design and materials, developed to match specific environmental conditions (Benson 1981). Consequently, it is assumed that ostracod carapace is subject to selection pressures (i.e. has adaptive value).

At the specific level ostracod carapaces include such a number of features (tubercles, ribs, nodes, spines, ...) and are conservative enough to be used for taxonomical identification in both neontological and, specially, paleontological studies. Carapaces, however, are not invariant morphological features at the specific level; indeed, valve shape variability has been extensively documented both within- and between-populations.

Methods for the study of shape change and variability

Form and shape

‘Form’ is an attribute of organisms that is made of two components: size and shape (Benson 1975, Bookstein 1989, Foote 1995, Baltanás et al. 2000). To discern between ‘form’

and ‘shape’ is not a trivial matter given that we frequently deal with information regarding ‘shape’ which, in fact, is related to ‘size’ (allometry). This is particularly the case when studying ontogenetic processes or in comparisons between individuals grown under different environmental conditions (Rohlf and Bookstein 1987).

Concerning methods and techniques available for the study of shape change and variability, several approaches exist but we will here concentrate in two: Traditional Morphometrics and Geometric Morphometrics.

Traditional Morphometrics

This approach, also named Multivariate Morphometry (Reyment 1985, Foote 1995) and Multivariate Biometry (Bookstein 1993), is an application of multivariate statistics to morphometric issues. Although widely used, these techniques have a main flaw: they do not recognize the geometric origin of the data under scrutiny. Variables used in this approach — distances, angles and ratios—are out of context both geometrically and biologically (Bookstein 1993). In other words, the set of variables used in these procedures preclude the reconstruction of the original shape out of their values. Such loss of information makes these methods of limited value.

Statistical techniques aimed to study relationships between morphological features (length, height, weight, ...) developed well before the term ‘biometry’ was coined (Galton 1869, 1889). Examples can be found in the works of Montbeillard, Quetelet and Galton; as well as in later contributions by Edgeworth, Pearson, Fisher and Wright.

The many multivariate techniques existing, which have been applied to numerous sets of meristic data derived from a plethora of organisms, emphasize the structure of the covariance matrix over other aspects of the measurements and lack any connection to the geometrical arrangement of such measurements, their biological meaning or the functional processes related to the organism development (Bookstein 1993). Such situation can be noticed in the first publications that use the term ‘morphometry’ in its current use (Blackith 1965, Blackith et al. 1971).

In addition, traditional morphometrics has some severe limitations (Lestrel 1997): (a) it is highly subjective; (b) it does not preserve information on, *i.e.* it is not possible to recover the original shape out of morphometric variables used (distances, angles and ratios); and (c) all variables used are but a small amount of all information about shape contained in a biological object.

Aware of such circumstances, several scientists (Jolicoeur 1963, Burnaby 1966, Mosimann 1970) tried to put additional emphasis on the biological foundations of morphometric data. Their attempt, however, was not successful enough. The actual turnover occurs at the beginning of the '80s with the rise of the so-called Geometric Morphometrics (Rohlf 1990a, Rohlf and Marcus 1993, Bookstein 1991, 1993).

Geometric Morphometrics

Geometric morphometrics inspire, partially at least, in the work of D'Arcy W. Thompson (1942) who approached the study of biological shape change as distortions occurring in a cartesian coordinate system which have been previously selected on the basis of its biological homology. Shape is a definite entity, a configuration of points that keep geometric relationships among them and cannot be split into isolated items (like length or height). Confronted with a biological shape, the morphometrician will attempt to describe it in terms of transformation from an original reference shape. Although the approach proposed by Thompson was very appealing and promising it was not accompanied by any analytical procedure. It was the arrival of the computer age, several decades later, that makes it possible to develop application for morphometric analysis based on Thompson's ideas feasible (Bookstein 1993).

Within geometric morphometrics, comparisons between organic forms are addressed by collecting information concerning the location of discrete points, called *landmarks*. A set of homologous points, *landmarks*, provides information of the biological form given they are distributed homogeneously on the organism and bear some biological meaning (Goodall 1983, Bookstein 1984, 1986, Chapman 1990, Rohlf and Slice 1990, Schweitzer and Lohmann 1990, Reilly 1990, Bookstein 1991, 1993, Reyment and Abe 1995, Foote 1995, Stone 1998). The analysis of configurations of *landmarks* allows the study of shape change without decomposing it into artificial variables. There are several landmark-based methods (fig. 1) and an updated review can be found in Zelditch et al. 2004. Concerning ostracods, there are several studies that apply landmark methods (Kaesler and Foster 1987, Reyment et al. 1988, Abe et al. 1988, Reyment and Bookstein 1993, Reyment 1995, 1997, Elewa 2004).

For a large number of ostracod species, however, it is not possible to identify *landmarks*, or, at least, a number of landmarks large enough to make that approach feasible. Under such circumstances there is an option: *Outline Analysis* (Rohlf 1990b). Outline analysis operates on the following basis: (1) when landmarks are not available one should record the

positions of a rather high number of points along the contour of the studied object; (2) a mathematical function must be fitted to such observations in order to (3) explore differences between shapes through the analysis of the mathematical descriptors fitted to them. This approach includes a variety of specific methods (fig. 1), among others ‘*Eigenshape*’ analysis (Lohmann 1983, Schweitzer et al. 1986, Lohmann and Schweitzer 1990), standard Fourier descriptors (Kaesler and Waters 1972), and Elliptic Fourier Analysis (Kuhl and Giardina 1982, Kaesler and Maddocks 1984, Rohlf and Archie 1984, Foote 1989, Rohlf 1995, Lestrel 1997, McLellan and Endler 1998, Baltanás and Geiger 1998).

Geometric Morphometrics




Figure 1: Sketch of relationships between some methods in the realm of Geometric Morphometrics.

Course Outline

Sessions in the course will offer a close view to some of the methods mentioned above together with exercises dealing with related aspects like ‘Data Acquisition Procedures’ and ‘Multivariate Analysis of Shape Descriptors’.

Selected References*

[*References here included are those mentioned in the text above and many others which have not been explicitly quoted but that might be of interest for those attending the course]

- Abe, K., Reyment, R. A., Bookstein, F. L., Honigstein, A., Almogi-Labin, A., Rosenfeld, A., and Hermelin, O. 1988. Microevolution in two species of ostracods from the Santonian (Cretaceous) of Israel. *Historical Biology*, 1: 303-322.
- Alcorlo, P., Baltanás, A., and Arqueros L. 1999. Intra-clonal shape variability in the non-marine ostracod *Heterocypris barbara* (Crustacea, Ostracoda). *Yerbilimeri (Geosound)*, 35: 1-11.
- Bachnou, A., Carbonnel, G., and Bouab, B. 1999. Morphométrie des Hemicysterinae (Ostracodes) par modélisation mathématique du profil latéral externe. Application systématique et phylogénétique. *C.R. Acad. Sci. Paris, Sect. Terre-Planetes. Paléontol.*, 328: 197-202.
- Baltanás, A., Alcorlo, P. and Danielopol, D. L. 2002. Morphological disparity in populations with and without sexual reproduction: a case study in *Eucypris virens* (Crustacea: Ostracoda). *Biological Journal of the Linnean Society*, 75: 9-19.
- Baltanás, A., Brauneis, W., Danielopol, D. L. and Linhart, J. 2003. Morphometric methods for applied ostracodology: tools for outline analysis of non-marine ostracodes. *The Paleontological Society Papers*, 9: 101-118
- Baltanás, A., Namiotko, T. and Danielopol, D. L. 2000a. Biogeography and disparity within the genus *Cryptocandona* (Crustacea, Ostracoda). *Vie et Milieu*, 50: 397-310.
- Baltanás, A., Otero, M., Arqueros, L., Rossetti, G. and Rossi, V. 2000b. Ontogenetic changes in the carapace shape of the non-marine ostracod *Eucypris virens* (Jurine). *Hydrobiologia*, 419: 65-72.
- Baltanás, A. and Geiger, W. 1998. Intraspecific morphological variability: morphometry of valve outlines. In: K. Martens (Ed.) *Sex and Parthenogenesis. Evolutionary Ecology of Reproductive Modes in Non-Marine Ostracods*. Leiden: Backhuys Publishers, 127-142.
- Benson, R. H., Chapman, R. E. and Siegel, S. 1982. On the measurement of morphology and its change. *Paleobiology*, 8(4): 328-339.
- Benson, R. H. 1975. Morphological Stability in Ostracoda. *Bulletin of American Paleontology*, 65: 13-46.

- Benson, R. H. 1976. The evolution of the ostracode *Costa* analyzed by "Theta-Rho" difference. *Abhandlungen und Verhandlungen des Naturwissenschaftlichen Vereins in Hamburg*, (NF), 18/19 (Suppl.): 127-139.
- Benson, R. H. 1981. Form, Function, and architecture of ostracode shells. *Ann. Rev. Earth Planet. Sci.*, 9: 59-80.
- Benson, R. H. 1982. Deformations, Da Vinci's concept of form, and the analysis of events in evolutionary history. In: E. Montanaro (Ed.) *Paleontology, essential of historical geology*. Istituto di Paleontologia, Università di Módena, Módena. pp. 241-277.
- Blackith, R. and Reyment, R. 1971. *Multivariate Morphometrics*. Academic Press, New York.
- Blackith, R. 1965. *Morphometrics*. In: T. H. Waterman and H. J. Morowitz (Eds.) *Theoretical and Mathematical Biology*. Blaisdell, New York.
- Bookstein, F. L. 1984. Tensor biometrics for changes in cranial shape. *Annals of Human Biology*, 11(5): 413-437.
- Bookstein, F. L. 1986. Size and shape spaces for landmark data in two dimensions. *Statistical Science*, 1: 181-242.
- Bookstein, F. L. 1989. "Size and shape": a comment on semantics. *Systematic Zoology*, 38: 173-180.
- Bookstein, F. L. 1991. *Morphometric tools for landmark data: Geometry and Biology*. New York: Cambridge University Press.
- Bookstein, F. L. 1993. A brief history of the morphometric synthesis. In: L. F. Marcus, E. Bello and A. García-Valdecasas (Eds.) *Contributions to Morphometrics*. Monografías. Museo Nacional de Ciencias Naturales. Consejo Superior de Investigaciones Científicas. pp. 15-40.
- Burke, C. D., Full, W. E. and Gernant, R. E. 1987. Recognition of fossil fresh water ostracodes: Fourier shape analysis. *Lethaia*, 20: 307-314.
- Burnaby, T. P. 1966. Growth-invariant discriminant functions and generalized distances. *Biometrics*, 22(1): 96-110.
- Burstin, J. and Charcosset, A. 1997. Relationship between phenotypic and marker distances: theoretical and experimental investigations. *Heredity*, 79: 477-483.
- Chapman, R. E. 1990. Conventional Procrustes Approaches. In: F. J. Rohlf and F. L. Bookstein (Eds.) *Proceedings of the Michigan Morphometrics Workshop Special Publication*. The Natural Museum of Natural History. The Smithsonian Institution. Washington, D. C. 2 (12): 251-267.

- Cherry, L. M., Case, S. M., Kunkel, J. G. and Wilson, A. C. 1979. Comparisons of frogs, humans and chimpanzees. *Science*, 204: 435.
- Cherry, L. M., Case, S. M., Kunkel, J. G., Wyles, J. S. and Wilson, A. C. 1982. Body shape metrics and organismal evolution. *Evolution*, 36(5): 914-933.
- Ciampaglio, C. N., Kemp, M. and McShea, D.W. 2001. Detecting changes in morphospace occupation patterns in the fossil record: characterisation and analysis of measure of disparity. *Paleobiology*, 27: 695-715.
- Danielopol, D. L., Ito, E., Wansard, G., Kamiya, T., Cronin, T. and Baltanás, A. 2002. Techniques for Collection and Study of Ostracoda. In: J. A. Holmes, A. R. Chivas (Eds.) *The Ostracoda: Applications in Quaternary Research*. Washington DC: The American Geophysical Union. pp. 65-97.
- Dayan, T., Simberloff, D., Tchernov, E. and Yom-Tov, Y. 1990. Feline canines: Community-wide character displacement among the small cats of Israel. *American Naturalist*, 136: 39-60
- Digby, P. G. N. and Kempton, R. A. 1987. *Multivariate Analysis of Ecological Communities*. Chapman and Hall, London.
- Dryden, I. L. and Mardia, K. V. 1998. *Statistical Shape Analysis*. John Wiley and Son, Chichester.
- Elewa, A.M. T. 2004. Application of geometric morphometrics to the study of shape polymorphism in Eocene ostracodes from Egypt and Spain. In: Elewa, A. M. T (Ed.) *Morphometrics. Application in Biology and Paleontology*. Springer-Verlag, Berlin. pp. 7-28.
- Person, S, Rohlf, F. J. and Koehn, R. K. 1985. Measuring shape variation among two-dimensional outlines. *Systematic Zoology*, 34(4): 59-68.
- Fink, W. L. 1990. Data acquisition for morphometric analysis in systematic biology. In: F. J. Rohlf and F. L. Bookstein (Eds.) *Proceedings of the Michigan Morphometrics Workshop*. The University of Michigan, Museum of Zoology, Ann Arbor. pp. 9-19.
- Foote, M. 1989. Perimeter-based Fourier analysis: a new morphometric method applied to the trilobite cranidium. *Journal of Paleontology*, 63: 80-885.
- Foote, M. 1992. Paleozoic record of morphological diversity in blastozoan echinoderms. *Proc. Natl. Acad. Sci. USA*, 89: 7325-7329.
- Foote, M. 1993a. Discordance and concordance between morphological and taxonomic diversity. *Paleobiology*, 19(2): 185-204.

- Foote, M. 1993b. Contributions of individual taxa to overall morphological disparity. *Paleobiology*, 19(4): 403-419.
- Foote, M. 1995. Analysis of Morphological Data. In: N. L. Gilinsky and P. W. Signor (Eds.) *Analytical Paleobiology. Short Courses in Paleontology*. University of Tennessee and the Paleontological Society. Knoxville, 14: 59-86.
- Foote, M. 1997. The evolution of morphological diversity. *Annual Review of Ecology and Systematics*, 28: 129-152.
- Foster, D. W. and Kaesler, R. L. 1988. Shape analysis: Ideas from Ostracoda. In: M. L. McKinney (Ed.) *Heterochrony in Evolution*. Plenum Press. New York. pp. 53-69.
- Galton, F. 1869. *Hereditary genius: an inquiry into its laws and consequences*. London: Macmillan.
- Galton, F. 1889. *Natural Inheritance*. London: Macmillan.
- Goodall, C. R. 1983. The statistical analysis of growth in two dimensions. Doctoral dissertation, Dept. Statistics, Harvard University.
- Haines, J. A. and Crampton, J. S. 2000. Improvements to the method of Fourier shape analysis as applied in morphometric studies. *Paleontology*, 43: 765-783.
- Humphries, J. M, Bookstein, F. L, Chernoff, B., Smith, G. R., Elder, R. L. and Poss, S. G. 1981. Multivariate discrimination by shape in relation size. *Systematic Zoology*, 30 (3): 291-308.
- Irizuki, T. and Sasaki, O. 1993. Analysis of morphological changes through ontogeny: genera *Baffinicythere* and *Elofsonella* (Hemicytherinae). In: K. G. McKenzie and P. J. Jones (Eds.) *Ostracoda in the earth and life sciences*. A.A. Balkema, Rotterdam. pp.: 335-350.
- James, F. C. and McCulloch, C. E. 1990. Multivariate Analysis in Ecology and Systematics: Panacea or Pandora's box? *Annual Review of Ecology and Systematics*, 21: 129-166.
- Jolicoeur, P. 1963. The multivariate generalization of the allometry equation. *Biometrics*, 19: 497-499.
- Kaesler, R. L. and Foster, D. W. 1987. Ontogeny of *Bradleya normani* (Brady): Shape analysis of *landmarks*. In: Hanai, T., Ikeya, N. and Ishizaki, K. (Eds.) *Evolutionary Biology of Ostracoda*. Elsevier Kodansha. pp. 207-218.
- Kaesler, R. L. and Maddocks, R. F. 1984. Preliminary harmonic analysis of outlines of recent Macrocypridid Ostracoda. In: N. Krstic (Ed.) *The Taxonomy, Biostratigraphy and Distribution of Ostracodes*. Belgrade: Serbian Geological Society, 169-174.
- Kaesler, R. L. and Waters, J. A. 1972. Fourier analysis of the ostracode margin. *Geological Society of America Bulletin*, 83: 1169-1178.

- Kaesler, R. L. 1997. Phase angles, harmonic distance, and the analysis of form. In: P.E. Lestrel (Ed.) *Fourier descriptors and their applications in Biology*. Cambridge University Press, Cambridge. pp. 106-125.
- Koehl, M. A. R. 1996. When does morphology matter? *Annual Review of Ecology and Systematics*, 27: 501-542.
- Kuhl, F. P and Giardina, C. R. 1982. Elliptic Fourier features of a closed contour. *Computer Graphics and Image Processing*, 9: 236-258.
- Legendre, P. and Legendre, L. 1998. *Numerical Ecology*. (2nd ed.). Development in Environmental Modelling, 20. Elsevier, Amsterdam. pp: 853.
- Lestrel, P. E. 1997. Introduction and overview of Fourier descriptors. In: Lestrel, P. E. (Ed.) *Fourier descriptors and their applications in Biology*. 4. Cambridge U.P., Cambridge. pp. 22-24.
- Lohman, G. P. 1983. Eigenshape analysis of microfossils: A general morphometric procedure for describing changes in shape. *Mathematical Geology*, 15: 659-672.
- Lohmann, G. P. and Schweitzer, P. N. 1990. On Eigenshape Analysis. In: F. J. Rohlf and F. J. Bookstein (Eds.) *Proceedings of the Michigan Morphometrics Workshop*. Ann Arbor, Michigan: The University of Michigan Museum of Zoology. pp. 147-166.
- Loik, M. E. and Noble, P. S. 1993. Freezing tolerance and water relations of *Opuntia fragilis* from Canada and the United States. *Ecology*, 74: 1722-1732.
- MacLeod, N. 1999. Generalizing and extending the eigenshape method of shape space visualization and analysis. *Paleobiology*, 25: 107-138.
- Majoran, S. 1990. Ontogenetic changes in the ostracod *Cytherella* cf. *ovata* Roemer from the Cenomanian of Algeria. *J. Micropal.*, 9: 37-44.
- Maness, T. R. and Kaesler, R. L. 1987. Ontogenetic changes in the carapace of *Tyrrhenocythere amnicola* (Sars) a hemicytherid ostracod. *Univ. Kansas Paleontol. Contrib.* 118: 1-15.
- Marcus, V. and Weeks, S. C. 1997. The effects of pond duration on the life history traits of an ephemeral pond crustacean, *Eulimnadia texana*. *Hydrobiologia*, 359: 213-21.
- Marcus, L. F. 1993. Some aspects of multivariate statistics for morphometrics. In: L. F. Marcus, E. Bello, E. and A. García-Valdecasas (Eds.) *Contributions to Morphometrics*. Monografías del Museo Nacional de Ciencias Naturales 8, Madrid. pp. 95-130.
- McGhee, G. R. 1999. *Theoretical Morphology*. Columbia University Press, New York.
- McKinney, F. K. 1981. Planar branch systems in colonial suspension feeders. *Paleobiology*, 7: 344-354.

- McLellan, T. and Endler, J. A. 1998. The relative success of some methods for measuring the shape of complex objects. *Systematic Zoology*, 47(2): 264-281.
- Møller, A. P. 1994. *Sexual selection and the Barn Swallow*. Oxford University Press, Oxford.
- Mosimann, J. E. 1970. Size allometry: Size and Shape Variables with Characterizations of the Lognormal and Generalized Gamma Distributions. *Journal of the American Statistical Association*, 65(330): 930-945.
- Norberg, U. 1994. Wing design, flight performance, and habitat use in bats. In: P. C. Wainwright and S. M. Reilly (Eds.) *Ecological Morphology*. University of Chicago Press, Chicago. pp. 205-239.
- Pielou, E. C. 1977. *Mathematical Ecology*. Wiley-Interscience Publication. John Wiley and Sons. New York.
- Pielou, E. C. 1984. *The Interpretation of Ecological Data: A Primer on Classification and Ordination*. John Wiley and Sons, Inc. USA.
- Raup, D. M. 1967. Geometric analysis of shell coiling: Coiling in ammonoids. *Journal of Paleontology*, 41:43-65.
- Ray, T. S. 1990. Application of eigenshape analysis to second order leaf shape ontogeny in *Syngonium podophyllum* (Araceae). In: F. J. Rohlf and F. J. Bookstein (Eds.) *Proceedings of the Michigan Morphometrics Workshop*. The University of Michigan, Museum of Zoology, Ann Arbor. pp. 201-213.
- Reilly, S. M. 1990. Comparative Ontogeny of cranial shape in Salamanders using Resistant Fit Theta Rho Analysis. In: F. J. Rohlf and F. L. Bookstein (Eds.). *Proceedings of the Michigan Morphometrics Workshop Special Publication*. No: 2. The Natural Museum of Natural History. The Smithsonian Institution. Washington. Chapter 16. pp. 311-321.
- Reyment, R. A and Abe, K. 1995. Morphometrics of *Vargula hilgendorffii* (Müller), (Ostracoda, Crustacea). *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut*, 92: 325-336.
- Reyment, R. A. and Bookstein, F. L. 1993. Intraspecific variability in shape in *Neobuntonia airella*: an exposition of geometric morphometry. In: K. G. McKenzie and P. J. Jones (Eds.) *Ostracoda in the Earth and Life Sciences*. Rotterdam: AA Balkema. pp. 291-314.
- Reyment, R. A. 1985. *Multivariate Morphometrics and Analysis of Shape*. *Mathematical Geology*, 17(6): 591-609.
- Reyment, R. A., Bookstein, F. L., McKenzie, K. G. and Majoran, S. 1988. Ecophenotypic variation in *Mutilus pumilus* (Ostracoda) from Australia, studied by canonical variate analysis and tensor biometrics. *Journal of Micropalaeontology*, 7: 11-20.

- Reyment, R. A. 1991. *Multidimensional Paleobiology*. Pergamon Press, Oxford.
- Reyment, R. A. 1995. On multivariate morphometrics applied to Ostracoda. In: J. Riha (Ed.) *Ostracods and Biostratigraphy*. Rotterdam: AA Balkema. pp. 43-48.
- Reyment, R. A. 1997. Evolution of shape in Oligocen and Miocene *Notocarinovalva* (Ostracoda, Crustacea): a multivariate statistical study. *Bulletin of Mathematical Biology*, 59: 63-87.
- Rohlf, F. J. and Archie, J. 1984. A comparison of Fourier methods for the description of wing shape in mosquitos (Diptera: Culicidae). *Systematic Zoology*, 33(3): 302-317.
- Rohlf, F. J. and Bookstein, F. L. 1987. A comment on shearing as a method for "size correction". *Systematic Zoology*, 36(4): 356-367.
- Rohlf, F. J. and Marcus, L. F. 1993. A revolution in morphometrics. *Trends in Ecology and Evolution*, 8: 129-132.
- Rohlf, F. J. and Slice, D. 1990. Extensions of the procrustes method for the optimal superimposition of landmarks. *Systematic Zoology*, 39(1): 40-59.
- Rohlf, F. J. 1986. Relationship among eigenshape analysis, Fourier analysis, and analysis of coordinates. *Mathematical Geology*, 18: 845-854.
- Rohlf, F. J. 1990a. Morphometrics. *Annual Review of Ecology and Systematics*, 21: 299-316.
- Rohlf, F. J. 1990b. Fitting curves to outlines. In: F. J. Rohlf and F. J. Bookstein (Eds.) *Proceedings of the Michigan Morphometrics Workshop*. Ann Arbor, Michigan. The University of Michigan. Museum of Zoology. pp. 167-177.
- Rohlf, F. J. 2004. tpsDig, digitize landmarks and outlines, version 2.0. Department of Ecology and Evolution, State University of New York at Stony Brook.
- Roy, K. and Foote, M. 1997. Morphological diversity as a biodiversity metric. *Trends in Ecology and Evolution*, 12.
- Sampson, P. D., Bookstein, F. L., Sheehan, F. H. and Bolson, E. L. 1996. Eigenshape analysis of left ventricular outlines from contrast ventriculograms. In: L. F. Marcus (Ed.) *Advances in Morphometrics*. Plenum Press, New York. pp. 211-234.
- Schlichting, C. D. and Pigliucci, M. 1998. *Phenotypic Evolution – a Reaction Norm perspective*. Sinauer Associates, Inc.
- Schweitzer, P. N. and Lohmann, G. P. 1990. Life-history and the evolution of ontogeny in the ostracode genus *Cyprideis*. *Paleobiology*, 16: 107-125.
- Schweitzer, P. N., Kaesler, R.L. and Lohmann, G. P. 1986. Ontogeny and heterochrony in the ostracode *Cavellina* Coryell from Lower Permian rocks in Kansas. *Paleobiology*, 12: 290-301.

- Siegel, A. F and Benson, R. H. 1982. A robust comparison of biological shapes. *Biometrics*, 38(2): 341-350.
- Smith, L. H. and Bunje, P. M. 1999. Morphologic diversity of inarticulate brachiopods through the Phanerozoic. *Paleobiology*, 25(3): 396-408.
- Sneath, P. H. A. Trend-surface analysis of transformation grids. *J. Zool.* 1967, 151: 65-122.
- Stone, J. R. 1998. Landmark-Based Thin-Plate Spline Relative Warp analysis of Gastropod Shells. *Systematic Biology*, 47(2): 254-263.
- Thompson, D. W. 1942. On growth and form. Cambridge University Press. Second Edition.
- Wayne, R. K. and O'Brien, S. J. 1986. Empirical demonstration that structural genes and morphometric variation of mandible traits are uncoupled between mouse strains. *Journal of Mammalogy*, 67: 441-449.
- Weider, L. J., Beaton, M. J. and Hebert, P. D. N. 1987. Clonal diversity in high-arctic populations of *Daphnia pulex*, a polyploid apomictic complex. *Evolution*, 41: 1335-1346.
- Wills, M. A. 2001. Morphological disparity: a primer. In: J. M. Adrian, G. D. Edgecombe and B. S. Lieberman (Eds.) *Fossils, Phylogeny and Form: An Analytical approach*. Kluwer Academic/Plenum Publishers. pp. 55-144.
- Zahn, C. T. and Roskies, R. Z. 1972. Fourier descriptors for plane closed curves. *IEEE Trans. Comp.*, C-21: 269-281.
- Zelditch, M. L., Swiderski, D. L., Sheets, H. D. and Fink, W. L. 2004. *Geometric Morphometrics for Biologists*. Elsevier Academic Press, Amsterdam.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte des Institutes für Geologie und Paläontologie der Karl-Franzens-Universität Graz](#)

Jahr/Year: 2008

Band/Volume: [13](#)

Autor(en)/Author(s): Baltanás Angel

Artikel/Article: [Geometric Morphometrics - A contribution to the study of shape variability in Ostracods. 3-15](#)