

Ber. Inst. Erdwiss. K.-F.-Univ. Graz	ISSN 1608-8166	Band 14	Graz 2009
<i>Paleozoic Seas Symposium</i>		Graz, 14-18 th September 2009	

Sedimentary successions on Baltoscandia support an early start of the Ordovician-Silurian Icehouse period

CALNER, M.¹, LEHNERT, O.^{2,3} & NÖLVAK, J.⁴

(1) GeoBiosphere Science Centre, Lund University, Sölvegatan 12, SE-223 62 Lund, Sweden; mikael.calner@geol.lu.se

(2) Universität Erlangen, Geozentrum Nordbayern, Schlossgarten 5, D-91054 Erlangen, Germany; lehnert@geol.uni-erlangen.de

(3) Czech Geological Survey, Klárov 3/131, 118 21 Prague 1, Czech Republic

(4) Tallinn University of Technology, Institute of Geology, Ehitajate tee 5, 19086 Tallinn, Estonia; nolvak@gi.ee

The timing and controls for a shift from Greenhouse to Icehouse conditions in the Late Ordovician has been debated for many years. The terminal Ordovician (Hirnantian) glaciation was early recognised as a short-lived cooling of global climate in a longer-term Greenhouse cycle (FISHER 1981). The same view has later been supported by e.g. BRENCHLEY *et al.* (1994, 2003). A growing body of sedimentary and stable isotopic evidence suggests that cooling started several millions of year's earlier, in the early-middle Katian (FRAKES *et al.* 1992, SALTZMAN & YOUNG 2005, CALNER *et al.* 2009).

We provide sedimentary evidence from Baltoscandia for a major, and inferably climatically driven, sea-level fall and subsequent transgression close to the *Amorphognathus superbis* and *A. ordovicicus* zonal boundary in the middle Katian. Based on our high-resolution stable isotopic data (LEHNERT *et al.* 2008) this sea-level cycle overlaps with the Waynesville carbon isotope excursion, starting near the boundary between the Ka2 and Ka3 stage slices (see BERGSTRÖM *et al.* 2009). The combined evidence for sea-level change includes substantial, regional facies changes, preservation of erosional channels, and an associated palaeokarst horizon that can be traced several 100's of kilometres and across different depth-related facies belts of the basin. Contemporaneous sedimentary changes in the tropics (Laurentia), as well as in high latitude peri-Gondwana basins (Prague Basin), together with marked isotopic and faunal changes during this time interval suggest that this is an environmental perturbation that deserves global recognition.

In this presentation we focus on the recently documented palaeokarst in Baltoscandia. The palaeokarst horizon has an unusually wide spatial extent and can be followed more or less from the Caledonian foreland in the west, across Sweden and into the East Baltic area in the east. The horizon is several metres thick and related to an unconformity cut into fine-grained and sparsely fossiliferous carbonates formed at temperate latitudes: the Slandrom Limestone in Sweden and the time-equivalent Saunja Formation in the East Baltic area. The lithology of these formations and the morphology of solution cavities in them look unexpectedly similar in remote sections, such as in southern and central Sweden and in Estonia and Latvia. The combined evidence for a karst origin includes 1) frequent karren-like morphologies interpreted as 'Swiss-cheese' karst [*sensu* BACETA *et al.* 2001], 2) local occurrences of solution/collapse breccia, 3) presence of bladed pseudospars crystals in solution cavities (comparable to pseudospars illustrated by SCHOLLE & ULMER-SCHOLLE (2003), and interpreted as formed through early meteoric flushing), and 4) carbon isotope values indicating meteoric influence to the succession. These findings are put in context with previously reported, large-scale erosional channels that cut down several tens of metres below the Fjäckå Shale in the subsurface Baltic Sea area (TUULING & FLODÉN 2000), and with regional, anomalous thickness variations in the Slandrom Limestone and Saunja Formation, all together forming strong support for regional exposure of the Baltoscandian continent in the middle Katian. The surface of the palaeokarst horizon is overlain by the Fjäckå Shale (or lateral coeval shale formations) throughout large parts of Baltoscandia. This excellent marker bed constitutes dark brown to black, organic-rich shale and marks a swift transgression across the area in the early *A. ordovicicus* Zone. Based on chitinozoan

Ber. Inst. Erdwiss. K.-F.-Univ. Graz	ISSN 1608-8166	Band 14	Graz 2009
<i>Paleozoic Seas Symposium</i>		Graz, 14-18 th September 2009	

biostratigraphy, there is a notable hiatus between the time interval of palaeokarst formation and the flooding of the basin during the deposition of the Fjäcka Shale.

The results from Baltoscandia presented here and in a recent, related paper of our research group (CALNER *et al.* 2009) add to the series of publications that more recently have provided new evidence for an early start of the Ordovician-Silurian Icehouse period.

References

- BACETA, J.I., WRIGHT, V.P. & PUJALTE, V. (2001): Palaeo-mixing zone karst features from Palaeocene carbonates of north Spain: criteria for recognizing a potentially widespread but rarely documented diagenetic system. – *Sedimentary Geology*, 139: 205-216.
- BERGSTRÖM, S.M., CHEN, X., GUTIÉRREZ-MARCO, J.C. & DRONOV, A. (2009): The new chronostratigraphic classification of the Ordovician System and its relations to major regional series and stages and to $\delta^{13}\text{C}$ chemostratigraphy. – *Lethaia*, 42: 1-11.
- BRENCHLEY, P.J., CARDEN, G.A., MARSHALL, J.D., ROBERTSON, D.B.R., LONG, D.G.F., MEIDLA, T., HINTS, L. & ANDERSON, T.F. (1994): Bathymetric and isotopic evidence for a short-lived Late Ordovician glaciation in a greenhouse period. – *Geology*, 22: 295-298.
- BRENCHLEY, P.J., CARDEN, G.A., HINTS, L., KALJO, D., MARSHALL, J.D., MARTMA, T., MEIDLA, T. & NÖLVAK, J. (2003): High-resolution stable isotope stratigraphy of Upper Ordovician sequences: Constraints on the timing of bioevents and environmental changes associated with mass extinction and glaciation. – *Geological Society of America Bulletin*, 115: 89-104.
- CALNER, M., LEHNERT, O. & JOACHIMSKI, M. (2009): Carbonate mud mounds, conglomerates and sea-level history in the middle Katian (Upper Ordovician) of central Sweden. – *Facies*, published online 090515 [doi: 10.1007/s10347-009-0192-6].
- FISHER, A.G. (1981): Climatic oscillations in the biosphere. – *In*: NITECKI, M. (ed.): *Biotic crises in ecological and evolutionary times*. New York Academy press: 103-131.
- FRAKES, L.A., FRANCIS, J.E. & SYKTUS, J.I. (1992): *Climatic modes of the Phanerozoic*. – Cambridge University Press, Cambridge, 274 pp.
- LEHNERT, O., CALNER, M., JOACHIMSKI, M., BUGGISCH, W. & NÖLVAK, J. (2008): Intra-Baltic and Trans-lapetus correlation of Upper Ordovician $\delta^{13}\text{C}$ data from the Borensult-1 core (Östergötland, Sweden). – *In*: KRÖGER, B. & SERVAIS, T. (eds): *Palaeozoic Climates. Abstracts, International Congress, Lille, August, 22-31, 2008, Lille, France*: 61.
- SALTZMAN, M.R. & YOUNG, S.A. (2005): Long-lived glaciation in the Late Ordovician? Isotopic and sequence-stratigraphic evidence from Laurentia. – *Geology*, 33: 109-112.
- SCHOLLE, P.A. & ULMER-SCHOLLE, D.S. (2003): *A Color Guide to the Petrography of Carbonate Rocks: Grains, textures, porosity, diagenesis*. – *American Association of Petroleum Geologists Memoir*, 77: 474 pp.
- TUULING, I. & FLODÉN, T. (2000): Late Ordovician carbonate buildups and erosional features northeast of Gotland, northern Baltic Sea. – *GFF*, 122: 237-249.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte des Institutes für Geologie und Paläontologie der Karl-Franzens-Universität Graz](#)

Jahr/Year: 2009

Band/Volume: [14](#)

Autor(en)/Author(s): Calner Mikael, Lehnert O., Nolvak J.

Artikel/Article: [Sedimentary successions on Baltoscandia support an early start of the Ordovician-Silurian Icehouse period. 16-17](#)