

Ber. Inst. Erdwiss. K.-F.-Univ. Graz	ISSN 1608-8166	Band 14	Graz 2009
<i>Paleozoic Seas Symposium</i>		Graz, 14-18 th September 2009	

The Tremadocian through Darriwilian conodont succession of NE Spitsbergen: faunal affinities and intercontinental correlation

LEHNERT, O.¹, STOUGE, S.² & BRANDL, P.¹

(1) University of Erlangen-Nürnberg, Geozentrum Nordbayern, Abteilung Krustendynamik, Schloßgarten 5, D-91054 Erlangen, Germany; lehnert@geol.uni-erlangen.de, Philipp.Brandl@asamnet.de

(2) Geological Museum, University of Copenhagen, Øster Voldgade 5-7, DK-1350, Copenhagen K, Denmark; svends@snm.ku.dk

The strata of the Cambro–Ordovician Oslobreen Group, composed of the Tokammane (Cambrian), Kirtonryggen (Lower Ordovician) and Valhallfonna (Middle Ordovician) Formations are exposed along the coast of Ny Friesland in NE Spitsbergen. The Lower to Middle Ordovician conodont faunas of this region are very well preserved displayed also by the CAI 1 of the material in the collections.

A low yield is typical for the lower part of the Kirtonryggen Formation. The oldest Ordovician conodont faunas span the *Rossodus manitouensis* through *Macerodus dianae* zones (Tremadocian, Spora and Basisletta Member) followed by the Floian *Oepikodus communis* Zone recovered from the richer upper part of the formation (Nordporten Member). All the taxa are typical of the tropical North American Midcontinent faunal province (e.g., species of *Eucharodus*, *Ulrichodina*, *Aloxocoonus*, *Colaptoconus*, *Macerodus*, *Oneotodus*) and can well be correlated to the North American Standard Zonation (Ross *et al.* 1997). The different facies display deposition under warm conditions and large stromatolithes are recognized within the sub- to intertidal dolostones of the Basisletta Member (FORTEY & BRUTON 1973). Carbonates of the overlying Nordporten Member reflect more open marine conditions (mud- to wackestones with grainstone interbeds, water depth between FWB and SWB; BRANDL 2009).

The tropical Midcontinent-type assemblages dominated the region until open marine associations invaded the shallow shelf (i.e. *Evae* transgression). The pandemic *Oepikodus evae* becomes frequent and is succeeded by *O. intermedius* in abundance in the Olenidsletta Member, i.e. the lower member of the Valhallfonna Formation. In this part of the succession, the most common associated species are *Bergstroemognathus*, *Protoprioniodus*, *Protopanderodus*, *Oistodus*, *Wandelia*, *Phragmodus*, and *Semiacontiodus*. These faunas of mixed affinities display more open marine conditions which is supported by oxygen isotope data. The $\delta^{18}\text{O}$ values indicate that relatively cooler sea-water temperatures prevailed from the *O. evae* Zone and upwards. The Olenidsletta Member formed in deeper water environments and is dominated by dark platy, graptolite-bearing mudstones with thin marly intercalations (e.g., FORTEY & BRUTON 1973, FORTEY & COCKS 2003, BRANDL 2009). The upper strata (Profilbekken Member) are characterized by the *Periodon-Paroistodus* assemblage characteristic for outer shelf, shelf margin, and upper slope settings around Laurentia and may especially well be compared with the faunal succession of Newfoundland (e.g., STOUGE 1984, JOHNSTON & BARNES 1999).

The Valhallfonna Formation comprises the Floian and Dapingian stages and ends in the early Darriwilian *Lenodus variabilis* Zone. Palaeobiogeographically, the Ny Friesland faunas show close affinities to coeval assemblages from North-East Greenland and West Newfoundland. The occurrence of early Darriwilian *Phragmodus spicatus* and *Juanognathus leptosomatus* known from Australo-Asian peri-Gondwana allows us also to correlate precisely to these areas.

Our conodont zonation is closely tied to a recently established $\delta^{13}\text{C}$ isotope curve from Ny Friesland (BRANDL 2009) and a correlation to $\delta^{13}\text{C}$ curves from North East Greenland, Newfoundland (AZMY *et al.*, unpublished, pers comm. 2009) and Argentine Precordillera (BUGGISCH *et al.* 2004) is presented. Observed sea-level changes are compared with the detailed Ordovician sea-level curve established by NIELSEN (2004). Finally, the

Ber. Inst. Erdwiss. K.-F.-Univ. Graz	ISSN 1608-8166	Band 14	Graz 2009
<i>Paleozoic Seas Symposium</i>		Graz, 14-18 th September 2009	

palaeogeographical implications of characteristic gaps in the sedimentary successions of the different terranes along the northern Laurentian margin (e.g., HARLAND 1997, SMITH 2000, STOUGE *et al.* 2001, KNIGHT *et al.* 2001) are discussed.

References

- BRANDL, P. (2009): Carbon and oxygen isotopes, stratigraphy, and facies of the Oslobreen Group (northeast Ny Friesland, Svalbard). - Unpubl. Diploma thesis, Part B; Geozentrum Nordbayern, Universität Erlangen-Nürnberg, Germany, 114 pp.
- BUGGISCH, W., KELLER, M. & LEHNERT, O. (2003): Carbon isotope record of Late Cambrian to Early Ordovician carbonates of the Argentine Precordillera. – *Palaeogeography, Palaeoclimatology, Palaeoecology*, 195: 357-373.
- FORTEY, R.A. & BRUTON, D.L. (1973): Cambro-Ordovician rocks adjacent to Hinlopenstretet, north Ny Friesland, Spitsbergen. - *Geological Society of America Bulletin*, 84: 2227-2242.
- FORTEY, R.A. & COCKS, L.R.M. (2003): Paleontological evidence bearing on global Ordovician-Silurian continental reconstructions. – *Earth-Science Reviews*, 61: 245-307.
- HARLAND, W.B. (1997): Cambrian-Ordovician history. – *In*: HARLAND, W.B. (ed.): *The Geology of Svalbard*. – *Geological Society Memoir*, 17: 257-271.
- JOHNSTON, D.I. & BARNES, C.R. (1999): Early and Middle Ordovician (Arenig) conodonts from St. Pauls Inlet and Martin Point, Cow Head Group, Western Newfoundland, Canada. 1. Biostratigraphy and paleoecology. – *Geological et Palæontologica*, 33: 21–70.
- KNIGHT, I., BOYCE, W.D., STOUGE, S.S., HARPER, D.A.T. & CHRISTIANSEN, J.L. (2001): Lower Ordovician carbonate rocks of central-east Greenland: Details of stratigraphic sections and their comparison to coeval carbonate shelf rocks in western Newfoundland. - *Report of Activities 2001*, Government of Newfoundland and Labrador, Department of Mines and Energy: 16-19.
- NIELSEN, A.T. (2004): Ordovician sea level changes: A Baltoscandian Perspective. – *In*: WEBBY, B.D., PARIS, F., DROSER, M., PERCIVAL, I., (eds): *The Great Ordovician Diversification Event*. Columbia University Press: 84-93.
- ROSS, R.J. JR., HINTZE, L.F., ETHINGTON, R.L., MILLER, J.F., TAYLOR, M.E. & REPETSKI, J.E. (1997): The Ibexian, lowermost series in the North American Ordovician. - *United States Geological Survey Professional Paper*, 1579-A: 1-50.
- SMITH, P.M. (2000): Cambro–Ordovician stratigraphy of Bjørnøya and North Greenland: Constrains on tectonic models for the Arctic Caledonides and Tertiary opening of the Greenland Sea. – *Journal of the Geological Society*, London, 157: 459-470.
- STOUGE, S.S. (1984): Conodonts of the Middle Ordovician Table Head Formation, western Newfoundland. – *Fossils and Strata*, 16: 1-145.
- STOUGE, S., BOYCE, W.D., CHRISTIANSEN, J.L., HARPER, D.A.T. & KNIGHT, I. (2001): Lower–Middle Ordovician stratigraphy of North-East Greenland. – *Geology of Greenland Survey Bulletin*, 191: 117-125.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte des Institutes für Geologie und Paläontologie der Karl-Franzens-Universität Graz](#)

Jahr/Year: 2009

Band/Volume: [14](#)

Autor(en)/Author(s): Lehnert O., Stouge Svends, Brandl Philipp

Artikel/Article: [The Tremadocian through Darriwilian conodont succession of NE Spitsbergen: faunal affinities and intercontinental correlation. 45-46](#)