

Ber. Inst. Erdwiss. K.-F.-Univ. Graz	ISSN 1608-8166	Band 14	Graz 2009
<i>Paleozoic Seas Symposium</i>		Graz, 14-18 <sup>th</sup> September 2009	

## Stable carbon and oxygen isotope values from Silurian brachiopod shells from Gotland (Sweden): primary or diagenetically altered signals?

WACKER, U.<sup>1</sup>, JOACHIMSKI, M.M.<sup>2</sup> & MUNNECKE, A.<sup>1</sup>

(1) Geozentrum Nordbayern, Fachgruppe Paläoumwelt, Loewenichstr. 28, D-91054 Erlangen;  
wacker.ulrike@web.de, axel.munnecke@gzn.uni-erlangen.de

(2) Geozentrum Nordbayern, Fachgruppe Krustendynamik, Schloßgarten 5, D-91054 Erlangen;  
joachimski@geol.uni-erlangen.de

Fossil brachiopods are commonly used for stable isotope analysis especially of Palaeozoic and Mesozoic sediments. The  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  signals are used for both stratigraphic correlation and palaeoenvironmental reconstructions since it is assumed that brachiopods secrete shell calcite in near-isotopic equilibrium with ambient sea water.

The aim of this study is to reconstruct high-resolution isotope records from preserved shells. Brachiopod shells derive from the Silurian of Gotland where brachiopods are known to be well-preserved (Fig. 1). The investigated organisms lived in tropical shallow water in the Baltic Sea.

To determine the state of preservation, the shells were screened by cathodoluminescence (CL) and scanning electron microscopy (SEM; Figs. 2, 3). Out of 100 brachiopod shells, four specimens were chosen for high-resolution sampling (300 $\mu\text{m}$  steps) of the secondary (fibrous) shell calcite. Samples were taken from longitudinal sections which were first analysed by CL. The shells were mostly non-luminescent but small areas of  $\text{Mn}^{2+}$ -induced luminescence were noticed in every shell. SEM analyses were performed in transverse sections. From every sampled brachiopod shell, four or five SEM-sections were prepared perpendicular to growth direction in order to get a better overview on the preservation state of the entire shell (Figs. 2, 3). Only oxygen isotope values will be discussed because they are more prone to diagenetic alteration than  $\delta^{13}\text{C}$  values.




Fig. 1: Brachiopod from Follingbo.




Fig. 2: Well-preserved secondary shell layer.




Fig. 3: Secondary shell layer diagenetically altered.

Ber. Inst. Erdwiss. K.-F.-Univ. Graz	ISSN 1608-8166	Band 14	Graz 2009
<i>Paleozoic Seas Symposium</i>	Graz, 14-18 <sup>th</sup> September 2009		

The high-resolution isotope curves of the four brachiopod shells are remarkably similar.  $\delta^{18}\text{O}$  values of all brachiopods decrease from the umbo to the margin (Fig. 4). Three shells show a pronounced positive shift in  $\delta^{18}\text{O}$  in the anterior part and a decrease towards the margin (Fig. 4), which, however, is less clear in the fourth example.

One potential explanation for the observed pattern is that brachiopods do not secrete shell calcite in isotopic equilibrium with sea water (vital fractionation effect) and that this oxygen isotope fractionation changed systematically during the lifetime of the brachiopod. Another explanation is that the signals are diagenetically overprinted. The maximum oxygen isotope variation measured in a single shell is 1.8 ‰. This variation translates into a change of water temperatures of nearly 8°C (assuming constant salinity). Though there are areas in modern tropical shallow water where such large changes are observed it seems unrealistic that the




Fig. 4:  $\delta^{18}\text{O}$  values along a brachiopod shell.

curves indicate a systematic warming during the lifetime of the brachiopods since these organisms most probably lived for several years. A diagenetic alteration for the observed decrease in  $\delta^{18}\text{O}$  appears a more realistic explanation since the changes in  $\delta^{18}\text{O}$  coincide with a change in the ultrastructure of shell, from well-preserved fibres near the umbo (Fig. 2) to altered fibres near the margin (Fig. 3). However, two observations argue against this possibility. First, the increases of the  $\delta^{18}\text{O}$  values close to the margin of the shells are difficult to explain by diagenetic processes, and secondly, the changes in ultrastructure observed along the shells do not correlate with an increase of cathodoluminescence.

Because a diagenetic influence especially close to the margin of the shells cannot be excluded at the present state of knowledge, the brachiopod  $\delta^{18}\text{O}$  records should be interpreted critically. High-resolution trace element ratios will give further information on the preservation state and will help to answer the question if the Silurian brachiopods from Gotland preserved their primary isotopic signals.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte des Institutes für Geologie und Paläontologie der Karl-Franzens-Universität Graz](#)

Jahr/Year: 2009

Band/Volume: [14](#)

Autor(en)/Author(s): Wacker Ulrike, Joachimski M. M., Munnecke Axel

Artikel/Article: [Stable carbon and oxygen isotope values from Silurian brachiopod shells from Gotland \(Sweden\): primary or diagenetically altered signals? 82-83](#)