

Ber. Inst. Erdwiss. K.-F.-Univ. Graz	ISSN 1608-8166	Band 16	Graz 2011
IGCP 596 Opening Meeting		Graz, 19-24 th September 2011	

Biodiversity of Devonian conodonts from the West Siberia

IZOKH, N.G.

Trofimuk Institute of Petroleum Geology and Geophysics SB RAS, Acad. Koptuyug Av., 3, Novosibirsk, 630090, Russia; IzokhNG@ipgg.nsc.ru

In West Siberia the Devonian deposits compose a part of the shelf belt along the western margin of the Siberian continent and extend from W to NE: from Altai to Salair and Kuznetsk Basin deepening farther north beneath the Mesozoic-Cenozoic cover of the West Siberian Lowland. Devonian marine sediments contain abundant benthic and pelagic fauna. Conodont associations were recovered from the NW and central parts of the West-Siberia Geo syncline (WSG), Rudny and Gorny Altai, Kuznetsk Basin.

Many conodont index-species of two (deep- and shallow-water) parallel zonal scales were recovered from the several Devonian sections.

The north-west of the West-Siberia Geosyncline. The Upper Emsian – middle Famennian conodont association from the Shchuchiy Ledge (IZOKH 2011). The Emsian association includes: *Polygnathus serotinus* TELFORD, *P. bultyncki* WEDDIGE, *Neopanderodus perliniatus* ZIEGLER & LINDSTROM, *N. cf. transitans* ZIEGLER & LINDSTROM, *Belodella devonica* (STAUFFER), *B. triangularis* (STAUFFER), *B. resima* (PHILIP), *Panderodus* sp. and *Pseudooneotodus* sp.

The Eifelian association is characterized by species derived from Emsian and entering Givetian. It includes: *Polygnathus serotinus* TELFORD, *P. costatus patulus* KLAPPER, *P. costatus partitus* KLAPPER *et al.*, *P. cf. linguiformis* HINDE, *P. costatus costatus* KLAPPER, *P. aff. P. trigonicus* BISCHOFF & ZIEGLER, *P. pseudofoliatus* WITTEKINDT, *P. eiflii* BISCHOFF & ZIEGLER, *Icriodus cf. regularicrescens* BULTYNCK, *Belodella devonica* (STAUFFER).

The Upper Frasnian association is represented by *Polygnathus decorosus* STAUFFER, *P. cf. P. aequalis* KLAPPER & LANE, *P. cf. P. samueli* KLAPPER & LANE, *P. cf. P. politus* OVNATANOVA, *Palmatolepis cf. Pa. rhenana* BISCHOFF, *Pa. cf. Pa. subrecta* MILLER & YOUNGQUIST, *Nothognatella* sp., *Belodella devonica* (STAUFFER).

Upper Famennian olistostrome-turbidite complex is composed of large limestone blocks characterized by three different in age conodont associations. The first is early Emsian which includes *Pedavis aff. sherryae* LANE & ORMISTON, *Pandorinellina cf. exigua philipi* (KLAPPER) and *Pseudooneotodus beckmanni* (BISCHOFF & SANNEMANN). The other two are Famennian: Second association characterizes triangularis Zone (*Palmatolepis praetriangularis* ZIEGLER & SANDBERG, *Pa. triangularis* Sannemann, *Icriodus alternatus* BRANSON & MEHL), third – belong to *marginifera*—Early *trachitera* zones (*Palmatolepis marginifera marginifera* HELMS, *Polygnathus* sp. A).

In the central part of WSG. *Ozarkodina remscheidensis remscheidensis* (WALLISER), *Oz. r. repetitor* (CARLS & GANDL), *Pandorinellina exigua philipi* (KLAPPER), and *Pand. steinhornensis miae* (BULTYNCK) were defined from the Lower Devonian deposits. The Emsian sequences yielded *Polygnathus kitabicus* YOLKIN *et al.*, *P. excavatus* CARLS & GANDL, *Pandorinellina e. exigua* (PHILIP), *Polygnathus nothoperbonus* MAWSON, *P. inversus* KLAPPER & JOHNSON, *P. serotinus* TELFORD and *P. foliformis* SNIGIREVA. The Eifelian association being very poor, only Upper Eifelian species were identified: *Tortodus kockelianus australis* and *P. x. ensensis*. *Icriodus obliquimarginatus* BISCHOFF & ZIEGLER, *Polygnathus x. xylus* STAUFFER, *P. varcus* STAUFFER, *Ozarkodina brevis* BISCHOFF & ZIEGLER, *Klapperina disparilis* ZIEGLER & KLAPPER were found in the Givetian. The Frasnian associations include *Palmatolepis hassi* MÜLLER & MÜLLER, *Ancyrognathus triangularis* KLAPPER, *Pa. gigas* MILLER & YOUNGQUIST, *Pa. subrecta* MÜLLER & YOUNGQUIST. The Famennian associations are dominated by *Palmatolepis* (*Pa. triangularis* SANNEMANN, *Pa. rhomboidea* SANNEMANN, *Pa. quadrantinodosa inflexoidea* ZIEGLER, *Pa. postera* ZIEGLER), with rare *Polygnathus* (*Polygnathus znepolensis* Spasov) (DUBATOLOV *et al.* 1990).

Salair, Rudny and Gorny Altai. The Lochkovian conodont association was found in Salair. It includes the following taxa: *Caudicriodus woschmidti transiens* CARLS & GANDL, *Pedavis cf. Ped. breviramus* MURPHY & MATTI, *Pandorinellina exigua philipi* (KLAPPER), *Pand. optima* (MOSKALENKO), *Pelekysgnathus serratus* JENTZSCH, *Ozarkodina e. excavata* BRANSON & MEHL, *Panderodus* sp. and

Ber. Inst. Erdwiss. K.-F.-Univ. Graz	ISSN 1608-8166	Band 16	Graz 2011
<i>IGCP 596 Opening Meeting</i>		Graz, 19-24 th September 2011	

Belodella sp. Predominant are the *Panderodus* elements (up to 75%) in the conodont collection. Most diverse conodont associations were established in the Emsian of Rudny Altai, Gorny Altai, Salair and Tuva Depression. They include species belonging to *Pandorinellina*, *Caudicriodus*, *Latericriodus*, *Ozarkodina*, *Polygnathus*, *Panderodus*, *Pelekysgnathus*, *Vjaloviodus* and *Belodella*. Almost all Emsian zonal index-species have been identified. Among them are *Polygnathus kitabicus* YOLKIN *et al.*, *P. excavatus* CARLS & GANDL, *P. nothoperbonus* MAWSON, *P. inversus* KLAPPER & JOHNSON, *P. serotinus* TELFORD. The Emsian-Eifelian boundary interval is expressed by non-marine deposits with flora remains.

The Eifelian sequences of Rudny Altai, Gorny Altai and Salair are characterized by *Polygnathus*, *Icriodus* and *Tortodus*, including *Polygnathus costatus partitus* KLAPPER *et al.*, *Po. costatus costatus* KLAPPER, *Po. linguiformis klapperi* CLAUSEN *et al.*, *Icriodus regularicrescens* BULTYNCK (BAKNAREV *et al.* 2004). The Givetian interval is represented by shallow-water associations of *Icriodus* with rare *Polygnathus* in Rudny Altai, NW Kuznetsk Basin and Minusa Depression (AKSENOVA *et al.* 1994, BAKNAREV *et al.* 2004). The *Icriodus obliquimarginatus* BISCHOFF & ZIEGLER species was identified in Rudny Altai and NW Kuznetsk basin.

Most diverse Frasnian associations which typify complete zonal succession (*falsiovalis* - *linguiformis* zones) were found in Rudny Altai. The Lower and Middle Frasnian carbonates contain abundant *Polygnathus*, *Ancyrodella*, *Mesotaxis*, and rarely, *Icriodus* and *Palmatolepis*. The Late Frasnian siliceous-terrigenous sequences contain rare *Palmatolepis* and *Polygnathus* (BAKNAREV *et al.* 2004, IZOKH *et al.* 2004, OBUT *et al.* 2007).

The Frasnian conodonts, comprising common *Polygnathus*, *Ancyrognathus*, *Icriodus*, few *Pelekysgnathus* and *Ancyrodella* from SE Altai differ from the above associations.

NW Kuznetsk Basin. The Frasnian sequences contain shallow-water biofacies association, mainly *Polygnathus* and *Icriodus*, single *Palmatolepis*, *Ancyrodella*, *Ancyrognathus*, *Polylophodonta*. Among them: *Polygnathus dengleri* BISCHOFF & ZIEGLER, *P. angustidiscus* YOUNGQUIST, *Po. brevilamiformis* OVNATANOVA, *P. webbi* STAUFFER, *P. aequalis* KLAPPER & LANE, *P. robustus* KLAPPER & LANE, *P. costulatus* ARISTOV, *P. samueli* KLAPPER & LANE, *Palmatolepis hassi* MÜLLER & MÜLLER, *Ancyrodella gigas* YOUNGQUIST, *An. nodosa* ULRICH & BASSLER). F/F interval is characterized by: *Polygnathus brevilaminus* BRANSON & MEHL, *Icriodus praealternatus* SANDBERG, Ziegler & Dreesen, *I. alternatus alternatus* BRANSON & MEHL, *I. alternatus helmsi* SANDBERG & DREESEN.

Polygnathus izhmensis KUZMIN, *P. gr. P. semicostatus* BRANSON & MEHL and *I. iowaensis* YOUNGQUIST & PETERSON are found beginning from the base of the Famennian. Conodonts diversity sharply increased at the base of the Lower Famennian crepida Zone: abundant *Polygnathus* and *Palmatolepis*, rare *Icriodus* and single *Ancyrolepis* (*Palmatolepis triangularis* SANNEMANN, *Pa. quadrantinodosalobata* SANNEMANN, *Pa. crepida* SANNEMANN, *Ancyrolepis cruciformis* ZIEGLER and others).

The Uppermost Famennian complex of conodonts from the NW Kuznetsk Basin contain *Polygnathus*, *Pelekysgnathus*, *Neopolygnathus*, *Bispathodus*, *Icriodus*, and *Siphonodella*.

This study has been supported by RFBR grants № 11-05-00737, 11-05-01105, RAS Projects 21, 25. Contribution to IGCP 596 Project.

References

- AKSENOVA, L.M., RODYGIN, S.A. & KHALYMBADZHA, B.G. (1994): Konodontovaya kharakteristika pogranychykh otlozhenii severnoi okrainy Kuzbassa. - *In*: PODOBINA, B.M. & RODYGIN, S.A. (eds.): Voprosy geologii Sibiri. 2. - Izd-vo Tomskogo un-ta, Tomsk: 320-336. [in Russian]
- BAKNAREV, N.K., SENNIKOV, N.V., YOLKIN, E.A., IZOKH, N.G. *et al.* (2004): Key Devonian sections from Rudny Altai, Salair and Kuznetsk Basin regions. - Publishing House of SB RAS, Novosibirsk, 104 pp. [in Russian]
- DUBATOLOV, V.N., KRASNOV, V.I., BOGUSH, O.I., AKSENOVA, L.M. *et al.* (1990): Stratigrafiya paleozoiskikh otlozhenii yugo-vostoka Zapadno-Sibirskoi plity. - Nauka, Novosibirsk, 181 pp. [in Russian]
- IZOKH, N.G., YOLKIN, E.A., BAKHAREV, N.K. (2004): Early Frasnian conodonts from the Rudny Altai (West Siberia) // *News of Paleontology and Stratigraphy*, 6-7. - Novosibirsk: Publishing House of SB RAS: 89-101. Suppl. to *J. Geologiya i Geofizika*, 45. [in Russian]
- IZOKH, N.G. (2011): Devonian conodont association of the Shchuchiy Ledge (NW of the West Siberian Geosyncline). - GEO-Sibir-2011. Nedropol'zovanie. Gornoe delo. Novye napravleniya i tekhnologiya poiska, razvedki i razrabotki mestorozhdenii poleznykh iskopaemykh. Sbornik Materialov VII mezhdunar. Nauchnogo Kongressa "GEO-Sibir-2011", 19-29 aprelya 2011, Novosibirsk. - SGGA, Novosibirsk, 2 (1): 96-99. [in Russian]

Ber. Inst. Erdwiss. K.-F.-Univ. Graz	ISSN 1608-8166	Band 16	Graz 2011
<i>IGCP 596 Opening Meeting</i>	Graz, 19-24 th September 2011		

OBUT, O.T., IZOKH, N.G. & YOLKIN, E.A. (2007): First occurrences of radiolarians and conodonts in Frasnian siliciclastic sequences of the Rudny Altai (south of West Siberia, Russia). Subcommission on Devonian Stratigraphy and IGCP 499 Devonian Land Sea Interaction, Eureka, Nevada, 9-17 September 2007. - Minuteman Press, Rochester, NY.: 69-70.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte des Institutes für Geologie und Paläontologie der Karl-Franzens-Universität Graz](#)

Jahr/Year: 2011

Band/Volume: [16](#)

Autor(en)/Author(s): Izokh N.G.

Artikel/Article: [Biodiversity of Devonian conodonts from the West Siberia. 49-51](#)