

Ber. Inst. Erdwiss. K.-F.-Univ. Graz	ISSN 1608-8166	Band 16	Graz 2011
IGCP 596 Opening Meeting		Graz, 19-24 th September 2011	

Taxonomic diversity of the late Famennian - early Carboniferous foraminifers of the South Urals

KULAGINA, E.I.

Institute of Geology, Ufa Scientific Centre, Russian Academy of Sciences, Ufa, 450077, K. Marx St. 16/2, Russian Federation; kulagina@anrb.ru

The taxonomic diversity of foraminifers from the Devonian-Carboniferous boundary beds was studied based on material from the key sections of the western slope of the South Urals in Russia (Zigan, Sikaza, Ryauzyak, Usuli) and the Berchogur sections of Kazakhstan. These sections are mainly composed of marine carbonates, apparently formed in a shallow-water environment.

This boundary interval includes the *Quasiendothyra communis*, *Q. kobeitusana*, *Tournayellina pseudobeata*, *Earlandia minima*, and *Chenyshinella disputabilis* foraminiferal zones of the Russian Scale. The *T. pseudobeata* Zone in the South Urals corresponds to the beds with remnant *Quasiendothyra*. The first appearance of the conodont *Siphonodella sulcata* is fixed within these beds. More than 110 species of 26 genera are identified in the Devonian-Carboniferous boundary interval (KOCHETKOVA *et al.* 1985, 1987). Ten genera are represented by unilocular and bilocular foraminifers (orders Parathuramminida, Earlandiida and Incertae sedis). The others are endothyrids and tournayellids. 41 species are determined in the *Q. communis* Zone. The number of foraminiferal species reaches 90 in the overlying zone, where the maximum diversification of the subfamily Quasiendothyrinae is recorded (in the taxonomy of RAUSER-CHERNOUSOVA *et al.* 1996). About 30 *Quasiendothyra* species and subspecies are recorded in this zone. The assemblage also includes species of *Septatournayella*, *Septaglomospiranella*, *Septabrunsiina*, *Tournayellina* (order Tournayellida).

The minimum species diversity is observed in the *Earlandia minima* Zone, where only 26 species are determined (Fig.1). Most of them unilocular taxa (*Archaesphaera* – 3 species, *Vicinesphaera* - 3, *Parathurammina* – 5, *Parathuramminites* – 3, *Bisphaera* – 5, *Eotuberitina* - 2). Bilocular forms are represented by *Rauserina notata*, *Earlandia minima* and *Caligella antropovi*. Of multi-locular foraminifers, only *Glomospiranella rara* and *Tournayellina cf. pseudobeata* are found in the *E. minima* Zone.

Species diversity begins to increase in the *Ch. disputabilis* Zone and reaches 60 in the overlying *P. tchernyshinensis* Zone due to radiation of the family Chernyshinellidae (Order Tournayellida). The total number of genera reaches 30. At the end of the Tournaisian, due to the extinction of tournayellids and the decreasing diversity of Tournaisian endothyrids, the species diversity drops to 50, in spite of the appearance of numerous short lived taxa.

Viséan and Serpukhovian deposits are widespread in the South Urals. The taxonomic diversity is studied in the sections of the west and east subregions of the Urals. In the early Viséan the species diversity increased from 80 in the *E. simplex* Zone to 100 in the *U. rotundus* Zone due to the rapid evolution of Archaediscida, Endothyrida and Staffellida. The number of genera increased to 50.

In the Late Viséan, species diversity reached its maximum for the Early Carboniferous. This time is characterized by maximum diversification of Endothyrida, Archaediscida, Palaeotextulariida, Ozawainellida (family Eostaffellidae).

At the Viséan-Serpukhovian boundary, species diversity almost halved compared to the underlying beds. More than 60 species known from the Upper Viséan did not continue into the Serpukhovian. At the same time the number of species of encrusting foraminifers increased, and several new eostaffellid and archaediscid (family Howchiniidae) species appeared. In total 45 new taxa appeared in the Serpukhovian, and after reaching maximum diversity in the *E. paraprotvae* Zone, species diversity again decreased.

The study was supported by the Russian Foundation for Basic Research, grant no. 10-05-01076.

Ber. Inst. Erdwiss. K.-F.-Univ. Graz	ISSN 1608-8166	Band 16	Graz 2011
IGCP 596 Opening Meeting		Graz, 19-24 th September 2011	

References

- KOCHETKOVA, N.M., ALEKSEEV, A.S., BARSKOV, I.S., BOGOSLOVSKIY, B.I., GAGIEV, M.KH., KONONOVA, L.I., KULAGINA, E.I., PAZUKHIN, V.N., REITLINGER, E.A., ROMANOV, V.A. & JANBULATOVA, M.G. (1987): Fauna i biostratigrafiya pogranichnykh otlozhenij devona i karbona Berchogura (Mugodzhary). - Moscow, Nauka, 121 pp. [in Russian]
- KOCHETKOVA, N.M., PAZUKHIN, V.N., REITLINGER, E.A. & SINITSINA, Z.A. (1985): Opornye razrezy pogranichnykh otlozhenij devona i karbona zapadnogo sklona Yuzhnogo Urala. Issue 6. Preprint. - Magadan, North-East Research Institute, FESC USSR Academy of Sciences, 66 pp. [in Russian]
- MENNING, M., ALEKSEEV, A.S., CHUVASHOV, B.I. *et al.* (2006): Global time scale and regional stratigraphic reference scales of Central and West Europe, East Europe, Tethys, South China, and North America as used in the Devonian – Carboniferous – Permian Correlation Chart 2003 (DCP 2003). - *Palaeogeography, Palaeoclimatology, Palaeoecology*, 240: 318–372.
- RAUSER-CHERNOUSOVA, D.M., BENSH, F.R., VDOVENKO, M.V., GIBSHMAN, N.B., LEVEN, E.YA., LIPINA, O.A., REITLINGER, E.A., SOLOVIEVA, M.N. & CHEDIYA, I.O. (1996): Reference-book on the systematic of the Paleozoic foraminifera (Endothyrida, Fusulinoida) - Moscow, Nauka, 204 pp. [in Russian]




Fig. 1: The species diversity of the late Famennian - early Carboniferous foraminifera of the South Urals. 1-16 – the foraminiferal zones: 1 – *Quasiendothyra communis*, 2 – *Q. kobeitusana*, 3 – *Tournayellina pseudobeata*; 4 – *Earlandia minima*; 5 – *Chernyshinella disputabilis*; 6 – *Palaeospiroplectamina tchernyshinensis*; 7 – *Spinoendothyra costifera*; 8 – *Eotextularia diversa*; 9 – *Eoparastaffella rotunda*; 10 – *E. simplex*; 11 – *Uralodiscus rotundus*; 12 – *Paraarchaediscus koktjubensis*; 13 – *Endothyranopsis crassa* – *Archaediscus gigas* (3 subzones); 14 – *Neoarchaediscus postrugosus*; 15 – *Eostaffellina paraprotvae*; 16 – *Monotaxinoides transitorius*.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte des Institutes für Geologie und Paläontologie der Karl-Franzens-Universität Graz](#)

Jahr/Year: 2011

Band/Volume: [16](#)

Autor(en)/Author(s): Kulagina E.I.

Artikel/Article: [Taxonomic diversity of the late Famennian - early Carboniferous foraminifers of the South Urals. 63-64](#)