

Ber. Inst. Erdwiss. K.-F.-Univ. Graz	ISSN 1608-8166	Band 16	Graz 2011
IGCP 596 Opening Meeting		Graz, 19-24 th September 2011	

Preliminary study of Late Middle Devonian Bentonites in Western Black Sea (Zonguldak-Bartın) Region, NW Turkey: a possible link with climate change

TÜRKMENOGLU, A.G.¹, GÖNCÜOĞLU, M.C.¹, YILMAZ, İ.O.¹ & ÜNLÜCE, O.¹

(1) Middle East Technical University, Department of Geological Engineering, Ankara, Turkey,
 asumant@metu.edu.tr; mcgoncu@metu.edu.tr; ioyilmaz@metu.edu.tr

In addition to climatic biological changes, extensive volcanic activity associated with the closure of the Rheic Ocean took place during Late Devonian. The products of explosive eruptions in the form of volcanic ash, after long distance transportation, were settled down and the diagenetic processes converted the ashes into potassium rich claystones called as K-bentonites or metabentonites. They are originally silicic volcanic ashes which are altered to claystones rich in smectite.

In the western Black Sea region, near Bartın and Zonguldak cities, yellowish brown and gray-green colored claystones having maximum thicknesses of 50-60 cm are exposed along with thick sequences of limestones-dolomitic limestones (Middle Devonian to Early Carboniferous in age, The Yılanlı formation). These carbonates were deposited on a shallow marine carbonate platform. Microfacies analysis of limestones indicate the cyclic nature of alternation charophyta and ostracoda rich wackestones, peloidal intraclastic packstones/wackestones and breccia along the section. Claystones are alternating with these microfacies without transition or gradual passage. Cyclic alternation of very shallow marine facies are interrupted by volcanic occurrences. Our preliminary investigations indicate that these claystone horizons are K-bentonites of volcanic origin (TÜRKMENOGLU 2001, TÜRKMENOGLU *et al.* 2009). Thus, they indicate successive Devonian explosive volcanic eruptions, with yet unknown source and distances.

In this research, two sections across bentonite-bearing successions in the Yılanlı Formation were measured and sampled both for clay mineralogical and paleontological purposes. The preliminary XRD data indicate that these K-bentonites are mainly consisting of illite, trace amounts of kaolinite, some quartz, feldspar, biotite, zircon, pyrite, calcite and gypsum. Chemical analyses demonstrate the volcanic origin of these K-bentonites. The foraminiferal content of the studied sections assign the Late Middle Devonian age (TÜRKMENOGLU *et al.* 2009). Results of this investigation have implications on tephra diagenesis and event-stratigraphic applications of the Bartın K-bentonites.

References

- TÜRKMENOGLU, A.G. (2001): A Paleozoic K-bentonite occurrence in Turkey. - Mid-European Clay Conference'01, MECC201, September 9-14, 2001, Stara Lesna, Slovakia.
- TÜRKMENOGLU, A.G., GÖNCÜOĞLU, M.C. & BAYRAKTAROĞLU, Ş. (2009): Early Carboniferous K-bentonite formation around Bartın: Geological implications. - 2nd Int. Symposium on the Geology of the Black Sea Region ISGB-2, Ankara-Turkey. Abstract Book, 209.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte des Institutes für Geologie und Paläontologie der Karl-Franzens-Universität Graz](#)

Jahr/Year: 2011

Band/Volume: [16](#)

Autor(en)/Author(s): Türkmenoglu A.G., Göncüoglu M.C., Yilmaz I. Ömer, Ünlüce O.

Artikel/Article: [Preliminary study of Late Middle Devonian Bentonites in Western Black Sea \(Zonguldak-Bartın\) Region, NW Turkey: a possible link with climate change. 88](#)