Ber. Inst. Geol. Paläont. K.-F.-Univ. Graz, ISSN 1608-8166, Band 6, Graz 2002

Abstract Volume

- Brooks, P.D., Stark, J.M., McInteer, B.B., Preston, T., 1989: Diffusion method to prepare soil extracts for automated nitrogen-15 analysis. Soil Sci. Soc. Am. J. 53, 1707-1711
- Sich, I., Russow R., 1998: ¹⁵N Analysis of nitric oxide and nitrous oxide in air by gas chromatography quadrupole mass spectrometry (GC-QMS) with cryotrap enrichment. Isotopes Environ. Health Stud. 34, 279-283
- Sich, I., Russow, R., 1999: ¹⁵N analysis of nitric oxide and nitrous oxide by cryotrap enrichment using a gas chromatograph quadrupole mass spectrometer and its application to ¹⁵N-tracer investigations of NO/N₂O formation in soil. Rapid Communication in Mass Spectrometry 13, 1325–1328
- Russow, R., Stevens R.J., 1996: Rapid, sensitive and highly selective ¹⁵N analysis of ¹⁵N enriched nitrite in water samples and soil extracts by nitric oxide production and CF-QMS Measurement. Isotopes Environ. Health Stud. 32, 323–328

Sulfur isotope distribution at Bleiberg lead-zinc deposit (Austria) and its genetic implication.

E. Schroll¹ & G. Rantitsch²

¹Institute of Mineralogy and Crystallography, University Vienna, Austria ²Institut für Geowissenschaften, Montanuniversität Leoben, Austria

Bleiberg (Carinthia, Austria) is the Alpine type deposit (ATP) of the low-temperature carbonate-hosted lead-zinc mineralisations. The ores are hosted by 350m Ladinian to Carnian platform carbonates. Genetically related lead-zinc deposits are found in Mežica (Slovenia) and Raibl (Italy). These deposits are located along the Periadriatic lineament, which separates the Austroalpine from the Southalpine tectonic unit. Mining ceased at the end of the 20th century. With a total production of 2.4 Mio t Pb+Zn, the deposit of Bleiberg is the largest deposit of the ATP.

The genesis of this deposit is discussed controversially. A Carnian sedimentary-sydiagenetic origin (Brigo et al. 1977, Schroll 1996) is opposed against a (MVT) model of migrating brines during post-Norian (Jurassic) age (Zeeh et al. 199?, Kuhlemann et al. 2002).

The interpretation of the sulfur isotope distribution in the Alpine ore district is an efficient contribution to evaluate these contrary models.

The Bleiberg deposit is characterized by a complex mineralisation, distributed in six ore horizons. In spite of the uniformity of the ore mineralization, differences in mineralogy, trace element geochemistry of sphalerite, and sulfur isotopes of sulfides are observed in dependence of the stratigraphic position and the structural setting of the mineralisation (Table 1). Geochemical data, e.g. Ga/Ge-ratio of sphalerite, indicate a leaching temperature around 100°C. The burial temperature of Carnian sediments at Bleiberg is estimated with ca. 130°C (Rantitsch 2001).

Ber. Inst. Geol. Paläont. K.-F.-Univ. Graz, ISSN 1608-8166, Band 6, Graz 2002

Abstract Volume

Ore horizon	Metal Zn+Pb mio t	Zn/Pb	Gang ue	Ge (ppm)	δ^{34} S‰ sulpides	δ^{34} S‰ barite
3 rd Cardita	0.1	6/1	Ba, F	(11)	-22.1 to -15.1 (19)	15,3 to 16,3 (2)
2 nd Cardita	Subeconomic		Ba, F		-19.3 to -5.0 (5))	12.1 to 14,8 (2)
1 st Cardita	0.2	8/1	F	340	-31.9 to -13.7 (44)	
Crest	0.5	6/1	F	160	-31.9 to -10.2 (29)	
Erzkalk	1.5	1/1 to 10/1	Ba, F	180	-27.0 to -3.0 (181)	12-0 to 17,1 (27)
Maxer Baenke	0.1	8/1	F	550	-31.6 to -13.7 (48)	

Table 1: General characteristic (F=fluorite, Ba=barite, n=number of samples) of the Bleiberg ore horizons (Cerny 1989, Schroll 1996).

348 sulfur isotope data from the Bleiberg deposit, obtained during thirty years of exploration, are available for this study. The sulfur isotope distribution of the Bleiberg deposit is show in Tab. 1 and Fig.1. The δ^{34} S-values of sulfides (n=303) display a wide variation from -3.0 (ZnS) to -31.6 (PbS). Three populations of δ^{34} S-mean values around δ^{34} S -7, -17 and -25 ‰ are identified. The δ^{34} S-values of barite (n=28) vary between 12.0 and 17.1‰ (mean 14.7‰). An isotopic equilibrium between the mineral phases is not recognisable.



Figure 1. Sulfur isotope distribution in the Bleiberg deposit.

The δ^{34} S-values of barite indicate precipitation of sulphate sulphur from the coeval seawater and evaporates of the Carnian stage (16%). Main source of sulfidic sulfur is the coeval seawater produced by bacterial activity (Kucha et al 2002). Stratiform mineralizations show highly depleted sulphur (~-25%) corresponding with the difference $\Delta\delta^{34}$ S% (sphalerite) – $\Delta\delta^{34}$ S% (seawater) of ~41% and with bacterial sulphate reduction (BSR) in an open system.

Ber. Inst. Geol. Paläont. K.-F.-Univ. Graz, ISSN 1608-8166, Band 6, Graz 2002

Abstract Volume

The bacterial activity is confirmed by optical and instrumental methods (Kucha et al. 2001). The lightest sulphur (-31.6%) is reported from late stage stalactitic botryoidally sphalerite. The origin of the sulfur isotope population which is characterized by a broad peak at -17% is multimodal: BSR, of pre-existing sedimentary iron sulphide or ore sulphide, mixing with hydrothermal sulphur (<-10%) dominate in discordant ore structures. The hydrothermal sulfur probably originates from sedimentary iron sulphides of the basement or its cover. Sulphur isotope patters of other carbonate-hosted Pb-Zn deposits in the Alps and anywhere support this assumption.

The detection of bacteriogenic activities during syndiagenetic sedimentary processe in sulphides from the most important ore horizons at Bleiberg deposit excludes the MVT-model, and confirms the model of Mid-Triassic age. The proposed model for the Bleiberg deposit corresponds with the interpretation of sulfur isotopes of the (Irish type) Lower Carboniferous giant lead-zinc deposit Navan in Ireland (Fallick et al. 2001).

References

- Brigo, L., Kostelka, L., Omenetto, P., Schneider, H.-J., Schroll, E., Schulz, Štrucl, I. (1977), in: Klemm, D., D., Schneider, H.-J. (eds: Time- and stratabound ore deposits, 273-293 Springer, Berlin Heidelberg.
- Cerny, I. (1989), Archiv. f. Lagerst. forsch. Geol. B.-A. 11, 5-126. Wien.
- Fallick, A., E., Ashton, J. H., Boyce, A., J., Ellam, R., M., Russell, M., J. (2001). Econ. Geol 96, 885-890.

Kucha H., Schroll E, Stumpfl E. (2001), in: Piestrzyńsjki, A. et al.: Mineral deposits at the Beginning of the 21th Century: 149-152, Balkema Publ., Lisse etc.

Kuhlemann, J., Vennemann, T., Herlec, U., Zeeh, St., Bechstädt, T. (2001), Econ.Geol.96,1931-1941

Rantitsch, G. (2001), Schweiz. Mineral. Petrol. Mitt. 81: 181-196

Schroll, E. (1996), Soc.Econ.Geol. Special Paper 4, 182-194, Soc. Econ . Geologists, Littleton, 1997.

Zeeh, S. Bechstädt, T. (1994), in: Fontbote, L. and Boni, M.(eds.: Sediment-hosted Zn-Pb ores. Geol. Soc.Spec. Publ. v. 10, 271-296, Springer, Berlin.

New applications of hydrogen isotopes in geo- and biochemistry

Z. Sharp

Department of Earth and Planetary Sciences, University of New Mexico Albuquerque, NM, 87131, USA

In spite of the enormous hydrogen isotope fractionations that occur in nature, the applications of hydrogen isotope geochemistry have been limited, in part due to the difficulty in making the isotopic analyses. However, new continuous flow analytical techniques, developed in the last several years, have made analyses vastly easier and less time consuming than those in the past. Much smaller amounts of material can be analyzed with a minimum of sample preparation. As a result, large number of analyses can be made in a relatively short time. In this communication, I present a number of varied applications of hydrogen isotope geochemistry, archaeology and ecology in order to illustrate the types of problems that can be addressed.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: <u>Berichte des Institutes für Geologie und Paläontologie der Karl-</u> <u>Franzens-Universität Graz</u>

Jahr/Year: 2002

Band/Volume: 6

Autor(en)/Author(s): Schroll Erich, Ranitsch Gert

Artikel/Article: Sulfur isotope distribution at Bleiberg lead-zinc deposit (Austria) and its genetic implication. 29-31