

## REMARKS ON LITHOSTROTIONID PHYLOGENY IN WESTERN NORTH AMERICA AND WESTERN EUROPE

Jerzy FEDOROWSKI\* & Edward Wayne BAMBER

\* Institute of Geology, Adam Mickiewicz University, Maków Polnych 16, Poland; [jerzy@amu.edu.pl](mailto:jerzy@amu.edu.pl)

\*\* Geological Survey of Canada (Calgary), 3303-33<sup>rd</sup> Street, N.W., Calgary, Alberta, T2L 2A7, Canada; [wabamber@nrcan.gc.ca](mailto:wabamber@nrcan.gc.ca).

Faunal exchange between Tournaisian-Viséan corals of western North America and Europe was limited by paleoenvironmental factors and by the great distance between those faunas, which were separated by the land-mass of Euramerica. Morphological comparison indicates that there is no direct relationship between species groups traditionally assigned to *Siphonodendron* in those areas. The immature morphology and stratigraphic distribution of the genus *Dorlodotia* suggest that it is the common ancestor for *Siphonodendron* in Europe and for *Siphonodendron*-like species in western North America. An unnamed, Ivorian species of *Dorlodotia* from western Canada initiated the phylogenetic succession that led to the North American *Siphonodendron*-like lineage in the latest Tournaisian and subsequently to the European *Siphonodendron* lineage in the early Viséan.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte des Institutes für Geologie und Paläontologie der Karl-Franzens-Universität Graz](#)

Jahr/Year: 2003

Band/Volume: [7](#)

Autor(en)/Author(s): Fedorowski Jerzy, Bamber Edward Wayne

Artikel/Article: [Remarks on lithostrotionid phylogeny in western north america and western europe 27](#)