

Origin of nitrate in groundwaters of Kalahari, Botswana

Susanne Stadler², Karsten Osenbrück¹, Kay Knöller¹, Thomas Himmelsbach²,
Axel Suckow^{3,4}, Stephan, M. Weise¹

¹Department of Isotope Hydrology, UFZ Centre for Environmental Research Leipzig-Halle, Germany, e-mail: karsten.osenbrueck@ufz.de

²Federal Institute for Geosciences and Natural Resources (BGR), Hannover, Germany, e-mail: s.stadler@bgr.de

³Isotope Hydrology Section, International Atomic Energy Agency (IAEA), Vienna


⁴Institute for Applied Geosciences (GGA), Hannover, Germany

In groundwaters from Kalahari, Botswana, elevated nitrate concentrations exceeding the WHO-guideline value of 50 mg/L have been found in mostly uninhabited semi-arid areas. Here we present first results of an isotope hydrological investigation with main emphasis to identification of input factors and the origin of that nitrate in groundwater.

The investigation area is located on the eastern fringe of the Kalahari between Serowe and Orapa, Botswana. Mean annual rainfall in the investigation area is about 450 mm/yr. The area is characterized by a flat topography at about 1200 m above sea level. Groundwater samples were taken from the Triassic Ntane sandstone aquifer, which comprises the main aquifer of the Karoo strata in the investigated area. The aquifer is mostly

confined by fractured Early Jurassic Stormberg basalts and an extensive cover with Kalahari sands.

According to their significantly different ¹⁴C contents, the investigated groundwaters can be divided into two distinct groups: modern groundwaters (40 to 90 pmC) of



Ca-Mg-HCO₃-Cl type and Na-HCO₃ type paleowaters (< 15 pmC), respectively. The high radiocarbon contents of the first group clearly confirm the presence of active recharge in the order of 1-5 mm/yr (Verhagen 1991). Only the modern groundwaters are affected by high nitrate concentrations. According to nitrate stable isotopes (Fig. 1), natural accumulation processes in soil (Heaton et al. 1983; Aranibar et al. 2003; Walvoord et al. 2003) seem to play the important role concerning the source of nitrate. The lack of an anthropogenic nitrate pollution in the investigated groundwaters indicated in Fig. 1 is in accordance with long residence times in the unsaturated zone, as suggested by tritium below detection limit in all groundwater samples.

Within the ¹⁴C containing (> 40 pmC) modern groundwaters, nitrate concentrations positively correlate with ¹⁴C content. If not due to anthropogenic activities, the decreasing nitrate concentration with increasing (corrected) ¹⁴C age (up to 3.000 years) must be a consequence of a change in nitrate leaching and/or of a change in recharge conditions.

The impact of different recharge conditions is confirmed by a strong linear correlation of nitrate with dissolved ²⁰Ne content (Fig. 2). ²⁰Ne is found in excess in all groundwater samples, which

is due to excess air dissolved in course of the recharge process. The extend of excess air is influenced by the rate and extend of recharge and therefore may be used as an indicator to distinguish between dry and wet infiltration conditions (Heaton et al. 1983; Stute and




Fig. 2. Nitrate and ²⁰Ne contents of Kalahari groundwaters out of the Ntane sandstone aquifer. Today's low recharge rates yield low ²⁰Ne and high nitrate contents as end member of the mixing line with high ²⁰Ne/low nitrate containing groundwaters recharged under more humid conditions.

Talma 1997; Beyerle et al. 2003). High nitrate contents correspond to low ²⁰Ne excess values, which suggest a slow re-charge where ²⁰Ne concentrations are maintained at so-

Ber. Inst. Erdwissenschaften K.-F.-Univ. Graz	Bd. 8	ISSN 1608-8166	Isotope Workshop Volume	Graz 2004
---	-------	----------------	-------------------------	-----------

lubility equilibrium. Low nitrate on the other hand is connected to recharge under wet conditions, where air is captured below the water table.

References

- Aranibar, J.N., Anderson, I.C., Ringrose, S., Macko, S.A., 2003. Importance of nitrogen fixation in soil crust of southern African arid ecosystems: acetylenen reduction and stable isotope studies. *J. Arid Environ.* 54, 345-358.
- Beyerle, U., Rueedi, J., Leuenberger, M., Aeschbach-Hertig, W., Peeters, F., Kipfer, R., 2003. Evidence for periods of wetter and cooler climate in the Sahel between 6 and 40 kyr BP derived from groundwater. *Geophys. Res. Lett.* 30 (4), 1173-1177.
- Heaton, T.H.E., Talma, A.S., Vogel, J.C., 1983. Origin and history of the nitrate in confined groundwater in the western Kalahari. *J. Hydrol.* 62, 243-262.
- Stute, M., Talma, S., 1997. Glacial temperatures and moisture transport regimes reconstructed from noble gases and $\delta^{18}\text{O}$, Stamriet aquifer, Namibia. In: *Isotope Techniques in the Study of Environmental Change*, IAEA, Vienna, 307-320.
- Verhagen, B.T., 1991. Detailed geohydrology with environmental isotopes: A case study at Seroew, Botswana. In: *Isotope Techniques in Water Resources Development*, IAEA, Vienna, 345-362.
- Walvoord, M.A., Phillips, F.M., Stonestrom, D.A., Evans, R.D., Hartsough, P.C., Newman, B.D., Striegl, R.G., 2003. A Reservoir of Nitrate Beneath Desert Soils. *Science* 302, 1021-1024

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte des Institutes für Geologie und Paläontologie der Karl-Franzens-Universität Graz](#)

Jahr/Year: 2004

Band/Volume: [8](#)

Autor(en)/Author(s): Stadler Susanne, Osenbrück Karsten, Knöller Kay, Himmelsbach Thomas, Suckow Axel, Weise Stephan M.

Artikel/Article: [Origin of nitrate in groundwaters of Kalahari, Botswana. 132-134](#)