

ON DEFORMATION AND METAMORPHISM OF THE PLATTENGNEISS SHEAR ZONE, EASTERN ALPS

Kurt STÜWE, Veronika TENCZER & Martin PUTZ

Institut für Erdwissenschaften, Universität Graz, Heinrichstr. 26; A-8010 Graz, Austria

Introduction

The Plattengneiss shear zone is one of the largest shear zones of the Eastern Alps. It is about 1 km thick, flat lying and is exposed over an area of about 600 km² in the Koralpe. During the Eo-Alpine peak metamorphic event it experienced eclogite facies metamorphism with peak pressures of 14 kbar and peak temperatures of 700°C. Overall, it is one of the highlights of hard rock geology in the Alpine orogen. However, surprisingly few geologists work on the shear zone and many puzzling questions remain. This contribution summarizes our motivation to study aspects of its structural and metamorphic history and some recent approaches of our working group to tackle them.

Questions and Motivations

Our principle motivation for working on the Plattengneiss stems from considerations of the Eo-Alpine metamorphic heat budget which indicate that the Eo-Alpine thermal event may need interpretation in terms of heating and cooling mechanisms that are outside our normal understanding of Barrovian terrains (e.g. Stüwe, 1998):

- *The metamorphic grade:* The metamorphic grade of the Plattengneiss indicates a ratio of peak temperature to peak pressure of some 14°C per Kilometer (using 700°C/14 kbar and 1 kbar ~ 3.6 km) (e.g. Stüwe and Powell, 1995). Such low *T/P* ratios can only be attained and preserved if rocks are buried and exhumed in a time interval that is shorter than the time required for thermal equilibration. However, fission track ages of the Koralpe indicate that final exhumation was no earlier than about 50 Ma (Hejl, 1997), so that several tens of my were available for its metamorphic cycle. This indicates that heating and cooling may not be related to conductive equilibration and exhumation, respectively.
- *The metamorphic field gradient:* The metamorphic field gradient from the Plattengneiss in the southern Koralpe to the Gleinalm crystalline is smooth indicating that these units acted as one continuous sequence during the Eo-Alpine (Tenczer and Stüwe, 2003). However, interestingly, pressures drop much more rapidly with distance than temperatures: Some 50 km north of the major Plattengneiss exposures, rocks still experienced peak temperatures around 600°C - 650°C, but peak pressures are significantly lower around 8-10 kbar. The field gradient appears therefore to indicate a heating mechanism where rocks at higher crustal levels were more thermally perturbed than those at lower crustal levels.
- *Preservation of isotopic and metamorphic equilibria:* Despite temperatures around 700°C, the major elements and isotopes of the Plattengneiss are extremely ill-equilibrated. This has long been a problem for geochronologists and metamorphic petrologists alike. The lack of equilibration at such high temperatures is likely to indicate either (a) extremely dry conditions or (b) extreme short duration for the high temperatures with the latter being difficult to reconcile with a "normal" heat source for metamorphism.

- *Tectonic setting and timing relationships:* The Eo-Alpine metamorphic peak occurred at a time when the Austroalpine nappe pile was already thickened. However, despite the over thickened crust, there is no obvious evidence that significant topography was developed at this time. One interpretation for this is that not only the crust, but also the mantle part of the lithosphere were over thickened at this time (Stüwe and Sandiford, 1995). While this interpretation is consistent with the absence of topography, it is inconsistent with the occurrence of Eo-Alpine metamorphism which should lead to cooling rather than heating.

All these considerations indicate that there is a problem with the metamorphic heat budget. In the past we have worked on these problems by investigating details of the PT path and cooling history (Stüwe and Powell, 1995; Ehlers et al., 1994; Stüwe and Ehlers, 1998) as well as the stresses and strain rates (Tenczer and Stüwe, 1999; Biermeier and Stüwe, 2003) during the Eo-Alpine. Aspects of our recent and as yet unpublished work in this general region is summarized here.

Metamorphism in the Plattengneiss

In order to constrain some of the questions discussed above, in particular the question on the duration of metamorphism, the heat source and the dryness of the rocks, we have aimed at constraining both details of the metamorphic PT path and the evolution of fluid content of the Plattengneiss (Tenczer et al., 2004).




Fig. 1: Thermodynamic PT pseudosection in the System NCKFMASH for a bulk composition appropriate to the Plattengneiss. The bulk composition was derived from a whole rock analysis by subtracting appropriate proportions of phases that did not re-equilibrate during the Eo-Alpine metamorphic event (see Tenczer et al., 2004 for details). The Asterisk marks the Eo-Alpine and Permian metamorphic peak conditions.

As a principal result of this work, we present in Fig. 1 a PT pseudosection for kyanite-absent varieties of the Plattengneiss. Among many other interesting features, this phase diagram highlights the possibility that partial melting may have occurred during the Eo-Alpine

evolution both at the metamorphic peak (if the water content is slightly higher) and during substantial isothermal decompression.

Deformation of the Plattengneiss shear zone

While several studies have investigated the significance of the mylonitic deformation of the Plattengneiss shear zone (Krohe, 1987; Kurz et al., 2002), the large scale geometry of the shear zone and its post-Eo-Alpine deformation history remains not very well understood. However, in view of the importance of its potential contribution to the mechanical energy production in the region, its volume and geometry are of crucial importance. Fig. 2 shows some results of a recent 3D modeling study in which we determined the present geometry of the shear zone within the Koralpe.




Fig. 2: A 3 dimensional model of the Plattengneiss shear zone. (a) Field data on dip and strike of the Plattengneiss superimposed on a contoured digital elevation model for the Koralpe. The view is from SE towards the NW across the Speikkogel. (b) Model result: The image shows a Landsat image draped over the digital elevation model for the Koralpe looking west across the Speikkogel. The lines show the intersection of the modeled Plattengneiss with the topography.

References

- BIERMEIER C. and STÜWE K., 2003 Strain Rates from Snowball garnets ? *Journal of Metamorphic Geology*, 21, 253-268.
- EHLERS, K., STÜWE K., POWELL, R., SANDIFORD, M. & FRANK, W., 1994. Thermometrically inferred cooling rates from the Plattengneiss, Koralpe region, Eastern Alps. *Earth and Planetary Science Letters*, 125, 307-321.
- HEJL, E., 1997. 'Cold spots' during the Cenozoic evolution of the Eastern Alps: thermochronological interpretation of apatite fission-track data. *Tectonophysics*, 272, 159-173.
- KROHE, A., 1987. Kinematics of Cretaceous nappe tectonics in the Austroalpine basement of the Koralpe region (eastern Austria). *Tectonophysics*, 136, 171-196.
- KURZ, W., FRITZ, H., TENCZER, V. & UNZOG, W., 2002. Tectonometamorphic evolution of the Koralpe Complex (Eastern Alps): Constraints from microstructures and textures of the „Plattengneiss“ – shear zone. *Journal of Structural Geology*, 24, 1957-1970.
- STÜWE K., 1998. Heat Sources of Cretaceous Metamorphism in the Eastern Alps - a discussion. *Tectonophysics*, 287, 251-269.
- STÜWE K., and SANDIFORD M., 1995: Mantle-lithospheric deformation and crustal metamorphism with some speculations on the thermal and mechanical significance of the Tauern event, Eastern Alps. *Tectonophysics*, 242, 115-132.
- STÜWE K., and POWELL R., 1995: Geothermobarometry from modal proportions. Application to a PT path of the Koralpe complex, Eastern Alps. *Contributions to Mineralogy and Petrology*, 119, 83-93.
- STÜWE K. and EHLERS K., 1998. Distinguishing Cooling Histories using Thermometry. Interpretations of Cooling Curves with some Examples from the Glein-Koralpe Region and the Central Swiss Alps. *Mitteilungen der österreichischen Geologischen Gesellschaft*, 89, 201-212.
- TENCZER V. and STÜWE K., 1999. Some constraints on strength and rheology of the Plattengneiss shear zone (Koralpe - Austroalpine). *Mitteilungen des Naturwissenschaftlichen Vereins für Steiermark*, 129, 43-53.
- TENCZER V. and STÜWE K., 2003 The metamorphic field gradient in the eclogite type locality. *Journal of Metamorphic Geology*, 21, 377-393.
- TENCZER V., STÜWE, K., POWELL R. and DALE J., 2004. Evolution of water content in a polymetamorphic terranes. The Plattengneiss shear zone of the Eastern Alps. *Contributions to Mineralogy and Petrology*, in review.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte des Institutes für Geologie und Paläontologie der Karl-Franzens-Universität Graz](#)

Jahr/Year: 2004

Band/Volume: [9](#)

Autor(en)/Author(s): Stüwe Kurt, Tenczer Veronika, Putz Martin

Artikel/Article: [On deformation and metaorphism of the Plattengneiss shear zone, Eastern Alps 392-394](#)