

IUGS Subcomm. Silurian Stratigraphy, Field Meeting 1994; Bibl. Geol. B.-A., 30/1994, Vienna

Silurian Retiolitids from Poland

A. Kozlowska-Dawidziuk¹

Isolated retiolitids fauna from Upper Llandovery to Upper Ludlow of NE Poland deep boring and Baltic erratic boulders was investigated. The process of reduction of reticula and clathrium (anca sleeve) as well reduction of edges of interthecal septa and central position of virgula started in the Wenlock. All Ludlow retiolitids possess central virgula in reduced rhabdosomes.

The appearance of the final growth phenomenon of the colony related to reduction the number of thecae in *Gothograptus* lineage started in Upper Wenlock and continued in the Ludlow. Classification and phylogenetic relationships among subfamily Plectograptinae Boucek et Münch, 1952 are assumed.

Sokolovograptus polonicus sp.n. from *Cyrtograptus murchisoni* Biozone represents strongly reduced rhabdosom. Rhabosome of *Plectodinemagraptus gracilis* gen. et sp.n. from *Cucullograptus hemiaversus/aversus* Biozone is the most reduced and stratigraphically latest retiolitid.

The investigated retiolitids comprise genera: *Retiolites*, *Pseudoretiolites*, *Paraplectograptus*, *Pseudoplectograptus*, *Sokolovograptus*, *Eisenackograptus*, *Gothograptus*, *Neogothograptus* gen.n., *Holoretiolites*, *Spinograptus*, *Plectograptus*, *Semiplectograptus* gen.n., *Plectodinemagraptus* gen.n.

¹ Institute of Paleobiology of the Polish Academy of Sciences, 02-089 Warsaw, Poland

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte der Geologischen Bundesanstalt](#)

Jahr/Year: 1994

Band/Volume: [30](#)

Autor(en)/Author(s): Kozlowska-Dawidziuk A.

Artikel/Article: [Silurian Retiolitids from Poland 137](#)