

The Distribution of the Chitinozoans in the Cella Section (Hirnantian - Lower Lochkovian). - A Preliminary Report.

by

Helga Priewalder
Geological Survey of Austria, Vienna
with 1 figure

Introduction

The investigations of the chitinozoans from the Cella section [Ashgill - Lochkovian] were part of a project with the goal of examining the geographic and stratigraphic distribution of the palynomorphs within the different facies of the Upper Ordovician to Lower Devonian strata in the Carnic Alps.

These are: the shallow water facies with mainly calcareous deposits, the siliciclastic basin facies and the transitional facies mediating between the former two.

In none of these facies *spores* could be observed. The *acritarchs* turned out to be strongly influenced by the local environments. Their only remarkable occurrence was in the Lower Silurian of the Cella section which belongs to the calcareous facies [PRIEWALDER, 1987]. However, the *chitinozoans* proved to be the geographically and stratigraphically widest distributed group of the palynomorphs.

Concerning the chitinozoans, 79 samples from the siliciclastic and transitional facies have been examined so far by spot checks: 60% of them were found to be fossiliferous.

From the Upper Ordovician to Lower Devonian sequence in the Cella section 95 samples have been prepared. 48 [= 51%] yielded chitinozoans.

As the chitinozoans were opaque to transmitting light the investigations had to be done mainly under SEM. About 4.300 micropalaeontological objects [chitinozoans as well as chitinozoan-like and/or problematic particles] have been examined in this way.

It has to be pointed out that the names of the chitinozoans in this report are provisional because they are based only on gross determinations. Detailed studies have yet to be carried out and will result in more diverse chitinozoan associations at many horizons of the section.

In the Cella section, the chitinozoans are restricted to four formations [fig.1]:

- ◆ the Plöcken Formation [**upper Ashgill**];
- ◆ the lower part of the Kok Formation [**upper Llandovery**];
- ◆ the sequence from the uppermost Kok Formation to the top of the Cardiola Formation [**upper Ludlow**];

- ◆ the sequence from the upper part of the Alticola Limestone to the lowermost Rauckofel Limestone [**Ludlow/Pridoli boundary - lowermost Lochkovian**].

Chitinozoans of the Upper Ordovician

In the Uggwa Shale and Uggwa Limestone chitinozoans are lacking. Instead, black and glossy particles with chitinozoan-like contours, probably consisting of graphite, are frequently present. In the light-microscope they can be confused with badly preserved chitinozoans.

Stratigraphically the chitinozoans occur for the first time at the base of the Plöcken Formation [sample **126**] with a few representatives of *Conochitina* EISENACK 1931 and probably also *Tanuchitina* JANSONIUS 1964. Further numerous melanosklerits with a strong resemblance to chitinozoans can be observed, as well as chitinozoan-like graphitic particles.

In the uppermost part of the Plöcken Formation a few samples [**128**, **129**, **45**] contain taxa which are diagnostic for the Ashgill: *Armoricochitina nigerica* (BOUCHÉ 1965) and *Tanuchitina elongata* (BOUCHÉ 1965). Furthermore *Desmochitina minor* EISENACK 1931, which does not range across the Ordovician/Silurian boundary, and representatives of *Conochitina*, *Rhabdochitina* (?) EISENACK 1931, *Spinachitina* SCHALLREUTER 1963 and the first specimen of the *Ancyrochitininae* with broken processes have been extracted.

The chitinozoan assemblages of this succession suggest the Hirnantian ***Tanuchitina elongata* - Biozone** (PARIS 1990).


The Ashgillian samples yield very few chitinozoans in a rather bad state of preservation: most specimens are three-dimensionally preserved, but broken.



Chitinozoans of the upper Llandovery

Sample **46A** at the very base of the Kok Formation [= upper Llandovery], which unconformably overlies the Plöcken Formation, yields a completely different chitinozoan fauna with a great number of individuals: numerous representatives of *Lagenochitinidae* and *Ancyrochitininae*, which cannot be determined exactly; *Ancyrochitina* gr. *ancyrea* (EISENACK 1931), *A. cf. diabolus* (EISENACK 1937), *Cyathochitina caputoi* DA COSTA 1971 and *Eisenackitina dolioliformis* UMNOVA 1976 which is very characteristic of this sample.

Samples **47**, **130** and **131** contain many specimens of *Bursachitina* TAUGOURDEAU 1966 and *Conochitina* [e.g. *C.sp. cf. emmastensis* NESTOR 1982], further *E. dolioliformis*, as well as *A. cf. nigerica* and *Laufeldochitina* ? sp., which are reworked taxa of upper Ordovician age.

This part of the sequence may be assigned to the upper Aeronian - lower Telychian ***Eisenackitina dolioliformis* - Biozone** [VERNIERS et al. 1995].





In sample **49**, an *Angochitina* species appears which is similar to *A. longicolla* EISENACK 1931, the index species of the following *Angochitina longicollis* - Biozone [VERNIERS et al. 1995] of upper Telychian age.

The upper part of the Llandoveryan strata of the Kok Formation [samples **50**, **132**] is characterized by chitinozoans which closely resemble *Conochitina proboscifera* EISENACK 1937, a typical species of the upper Telychian/lower Sheinwoodian period; *Conochitina* spp. [e.g. *C. sp. cf. C. armillata* TAUGOURDEAU & DE JEKHOWSKI 1960, *C. sp. cf. C. edjelensis elongata* TAUGOURDEAU 1963], *Eisenackitina sp.* and *Lagenochitina sp.* occur less frequently.

The uppermost Llandovery sample [**133**] yields only badly preserved individuals resembling *Angochitina longicolla*, as well as *Conochitina sp.*, *Cyathochitina sp.*, *Eisenackitina sp.* and *Sphaerochitina sp.*.

The chitinozoans from this part of the section are entirely or partly flattened and frequently folded. In cases of intense folding or variable flattening of the vesicles [e.g. thinner-walled necks are more, thicker-walled body chambers less strongly deformed] their contours may be altered to an extent that the original taxon is difficult to recognize.

Chitinozoans of the Wenlock - lower Ludlow

Throughout the Wenlock, the strata of which attain a thickness of only 5 meters indicating extreme condensation [SCHÖNLAUB 1994], and also in the lower Ludlow, associations of determinable chitinozoans are missing. Only sporadic and badly preserved fossils are present: Sample **135**: one fragment of *Belonechitina sp.*; sample **54**: the internal moulds of *Conochitinidae* indet.; sample **136**: fragments of *Conochitinidae* indet. and *Lagenochitinidae* indet., sample **56**: *Bursachitina sp.*, *Lagenochitinidae* indet., *Conochitinidae* indet..

Chitinozoans of the upper Ludlow

From the uppermost bed of the Kok Formation [sample **63**] to the top of the Cardiola Formation [sample **145**] a great variety of chitinozoans occurs.

At the base of this sequence [samples **63**, **141**] abundant and diverse *Angochitina* EISENACK 1931 [e.g. *A. echinata* EISENACK 1931 and a fragment similar to *A. elongata* EISENACK 1931], *Sphaerochitina* EISENACK 1955 [e.g. *S. sp. cf. impia* LAUFELD 1974], *Conochitina* EISENACK 1931 and a few *Bursachitina sp.* and *Eisenackitina sp.*, as well as some *Ancyrochitina sp.* appear.

Above this level a fragment of *Linochitina* EISENACK 1968 [sample **142**] is found.

Some *Cingulochitina cf. convexa* (LAUFELD 1974), *Sphaerochitina* spp. and *Angochitina* spp. [like in samples **63**, **141**] and a few *Ancyrochitininae* indet. are found in sample **64**.

The middle part is dominated by numerous *Conochitina* and *Belonechitina* JANSO-
NIUS 1964. *C.sp. cf. tuba* EISENACK 1932 occurs in sample **143**. Sample **66** yields
Belonechitina sp. cf. latifrons (EISENACK 1964) and *B.sp. cf. lauensis* (LAUFELD
1974) and rare *Sphaerochitina sp.*.

Finally, in the last sample of this succession [**145**] a few *Cingulochitina sp.* and *Ancyrochitininae* indet. are present.

The chitinozoans of this sequence seem thus to be referred to the upper Gorstian -
lower Ludfordian ***Angochitina elongata* - Biozone** [VERNIERS et al. 1995].

Here an other - unusual - state of preservation can be observed: the vesicles of thin-
walled taxa from limestones had collapsed three-dimensionally similar to a deflated
rubber ball. This feature probably had been developed at an early stage of diagenesis
when the internal cavities of the chitinozoans became dehydrated before mineral
fillings occurred. These fillings are common in chitinozoans from limestones and
they are responsible for their three-dimensional preservation.

From the base of the Alticola Limestone up to the end of the Ludlow the examined
samples yield no chitinozoans.

Chitinozoans of the uppermost Ludlow to the lower Lochkovian

A rich development of chitinozoans is documented from sample **73** of the Ludlow/
Pridoli boundary beds and persists through the Pridoli up to the end of the section in
the lower Lochkovian [sample **89**]. It comprises the upper part of the Alticola Li-
mestone, the Megaerella Limestone and the lowermost part of the Rauchkofel
Limestone.

Three samples at the base of this succession [**73** = uppermost Ludfordian sample;
74, 75 = lower Pridolian samples] contain numerous *Eisenackitina barrandei* PARIS
& KRIZ 1984, *E. granulata* (CRAMER 1964), *E. intermedia* (EISENACK 1955), *Urno-
chitina gr. urna* (EISENACK 1934), and some *Sphaerochitina cf. sphaerocephala*
(EISENACK 1932), *Ancyrochitina gr. ancyrea* (EISENACK 1931), *Angochitina sp.*,
Bursachitina sp. and *Gotlandochitina sp.*.

E. barrandei is the index-fossil of the ***Eisenackitina barrandei* - (total range) - Bio-
zone** [VERNIERS et al. 1995] which is restricted to the uppermost Ludfordian. At the
global stratotype section of the Ludlow/Pridoli-boundary at Pozáry Quarry of the Pra-
gue Basin, Bohemia, *E. barrandei* ranges a few decimeters into the Pridoli, where it
coexists within a very short distance with atypical representatives of *Urnochitina gr.
urna* [typical specimens are present in the higher parts of the Pridoli]. The latter is an
index species of the Pridoli appearing in the Prague Basin within an interval of a few
centimeters below to a few centimeters above the Ludlow/Pridoli-boundary [PARIS
in KRIZ et al. 1986].

Compared to the ranges in the Cellon section an obvious difference exists: here *E.
barrandei* ranges relatively high up into the Pridoli [almost 5 meters]; moreover, in
this interval it coexists with atypical *U. gr. urna*. Detailed studies will have to settle
this discrepancy.

The next sample [149] reveals a very low fossil content consisting of only some *Eisenackitina* sp., *Angochitina* sp. and *Ancyrochitina* ? sp..

The development of typical *Urnochitina urna* starts suddenly and with a great number of specimens in sample 76. As already described in the literature the fauna here too is of almost monospecific composition with the exception of only rare representatives of *Desmochitinidae* indet..

The following sample [149A] displays a special feature: *U. urna* becomes numerically unimportant, whereas large quantities of *Bursachitina krizi* (PARIS & LAUFELD 1980) are present. The residue consists almost entirely of representatives of the latter species.

Sample 150 is dominated by *U. urna*; in addition, only very few individuals of *B. krizi* and *Desmochitinidae* indet. occur.

The next three samples [151, 152, 153] yield insignificant associations with various *Lagenochitinidae* indet., a fragment of *B. krizi* [?], some *Angochitina* similar to *A. chlupaci* (PARIS & LAUFELD 1980) and *Sphaerochitina* sp..

The only taxon in the following sample 78 is *E. granulata* with a few well preserved individuals. It is still present in sample 154, but there accompanied by rare *B. krizi*, *Linochitina klonkensis* PARIS & LAUFELD 1980 and *Ancyrochitina* sp..

The strata between the samples 78 and 154 in the lower part of the Megaerella Limestone proved to be barren of chitinozoans.

At about this level of the section the chitinozoan fauna starts to rearrange: *U. urna* occurs with more and more decreasing numbers of individuals, while *Angochitina* EISENACK 1931, *Cingulochitina* PARIS 1981, *Gotlandochitina* LAUFELD 1974, *Linochitina* EISENACK 1968, *Sphaerochitina* EISENACK 1955 and especially *Ancyrochitina* EISENACK 1955 occur more frequently.

The most abundant species in sample 81, from which large quantities of chitinozoans could be extracted, represents *Ancyrochitina* sp. A, provided with simple processes with very broad basis. Other taxa are *L. klonkensis*, *Calpichitina corinnae* JAGLIN 1986, *Sphaerochitina* cf. *sphaerocephala* (EISENACK 1932), *Gotlandochitina* ? sp. and *U. urna* with very few specimens.

Samples 82 and 83, the uppermost Pridolian samples, contain only poor associations: very few *U. urna*, *Ancyrochitina* sp. A and *S. cf. sphaerocephala*.

The Pridoli is defined by the total range of *Urnochitina urna*, which at the global stratotype section for the Silurian/Devonian-boundary at Klonk, Prague Basin, disappears exactly at the boundary, while in the Karlstejn section it ranges a few decimeters above the base of the Lochkovian [PARIS, LAUFELD & CHLUPÁČ 1981].

Due to the lack of the index-fossils, the chitinozoan biozones of the Pridoli [*Fungochitina kosovensis* -, *Margachitina elegans* - and *Anthochitina superba* - Biozones, VERNIERS et al. 1995] could not be identified at Cellon.

Sample **84** from the lowermost Lochkovian bed yields a rich fauna: comparatively numerous *U. urna*, which is the last documented occurrence in the section; in addition, many well preserved and diverse representatives of *Angochitina*, *Gotlandochitina*, *Sphaerochitina* [e.g. *S. sphaerocephala*] and a few *Ancyrochitina* with unusual processes are found.

The chitinozoan assemblage of sample **85**, in which the number of individuals is rather low, is dominated by *Eisenackitina bohémica* (EISENACK 1934), a species typical of the Lochkovian, which in the Prague Basin appears a few decimeters above the base of the Devonian, i.e. in bed 21 at the Klonk section [PARIS 1981]. Co-occurring taxa are a few *Angochitina* aff. *chlupaci*, *Cingulochitina* sp., *Desmochitinidae* indet. and *Lagenochitinidae* indet..

In sample **156** *A. chlupaci* is present with several unequivocal individuals together with a few *Angochitina* sp. and *Desmochitinidae* indet..

The remaining samples in the section [**157**, **87**, **88**, **158** and **89** with a large quantity of chitinozoans] are dominated by numerous *Ancyrochitina* [at least 5 species]. Moreover they yield numerous diverse representatives of *Angochitina*, *Sphaerochitina*, *Gotlandochitina*, *Linochitina* and *Cingulochitina* [e.g. *C. ervensis* (PARIS 1979)] .

The chitinozoans of the Pridoli/Lochkovian sequence are generally three-dimensionally preserved, especially thicker-walled taxa; thinner-walled individuals are often more or less strongly collapsed.

Conclusions.

- ◆ In contrast to the acritarchs which are mainly restricted to the upper Llandovery to lower Wenlock sequence, the chitinozoans are present in almost all series of the Upper Ordovician to Lower Devonian succession of the Cellon section.

In several samples (46A, 141, 74, 76, 149A, 150, 81, 84, 89) they occur with large numbers of individuals and usually great diversity.

- ◆ The chitinozoan assemblages of the upper Ashgill and upper Llandovery strata separated by a gap of two stages are easily to distinguish.

The boundaries between the Llandovery and Wenlock, and the Wenlock and Ludlow, respectively, cannot be established by the aid of chitinozoans as these fossils are missing throughout the Wenlock and also in the lower Ludlow.

With regard to the chitinozoans, the position of the base of the Pridoli in the Cellon section is not yet clear and needs further investigations.

However, the base of the Lochkovian is well documented by diagnostic chitinozoan associations.

Several chitinozoan biozones can be identified:

- the Hirnantian *Tanuchitina elongata* - Biozone;
- the upper Aeronian - lower Telychian *Eisenackitina dolioliformis* - Biozone;

- the upper Telychian *Angochitina longicollis* - Biozone;
- the upper Gorstian - lower Ludfordian *Angochitina elongata* - Biozone;
- the uppermost Ludfordian *Eisenackitina barrandei* - Biozone;
- the Pridolian *Urnochitina urna* - Biozone;
- the lower Lochkovian *Eisenackitina bohémica* - Biozone.

- ◆ Obviously, environmental conditions were more favourable for the chitinozoans in the upper part of the section than in the lower. Starting with the topmost layer of the Kok Formation [upper Ludlow] up to the lower Lochkovian they show greater diversities and larger numbers of individuals and also better preservation than in the lower part.

Presently, the reasons for the occurrence of at least some chitinozoans in the unfavourable high energy environment of the Plöcken Formation and their absence in the off-shore low-energy facies of the Uggwa Limestone and the Uggwa Shale are difficult to explain.

- ◆ The Hirnantian-age chitinozoans of the Cellon section show a pronounced relationship with assemblages of the Northern Gondwana cold-water realm, while in the Silurian and Lower Devonian their affinities to representatives of the warm-water environments of Baltica/ Avalonia are obvious.

Because of the palaeogeographic vicinity of the two depositional areas, the Silurian-Lower Devonian chitinozoans of the studied section are very similar to those from Bohemia which is especially true for the upper Ludlow to lower Lochkovian sequence [DUFKA, 1992; KRIZ, 1992; KRIZ et al. 1986; PARIS & KRIZ, 1984; PARIS et al., 1981].

On the other hand in the Cellon section samples from the base of the Wenlock to the lower Ludlow yield no chitinozoans whereas in Bohemia diverse faunas can be obtained from coeval strata [KRIZ, 1992; KRIZ et al., 1993]. This phenomenon might be caused by unfavorable conditions for the chitinozoans' preservation [e.g. high hydrodynamic regime in a shallow sea, at least temporary; oxidation] in the sedimentary environment of the Cellon section.

Acknowledgment

The investigations were supported by the *Austrian Science Foundation* to whom I want to express my appreciation.

I would like to thank Dr. Florentin Paris, University of Rennes, France, for his engaged and valuable discussion of my large collection from the Cellon section.

References

- ACHAB, A., BERTRAND, R. & VAN GROOTE, G.: Chitinozoan Contribution to the Ordovician and Lower Silurian Paleobiogeography.- *J. Geol.*, **100**, 621-629, 5 fig., Chicago, 1992.
- BOUCHÉ, P.M.: Chitinozoaires du Silurien s.l. du Djado (Sahara nigérien).- *Rev. Micropaléont.*, **8**, 151-164, 3 tab., 3 pl., Paris, 1965.
- CRAMER, F.H.: Microplankton from three Paleozoic formations in the province of León (NW Spain). - *Leidse Geol. Meded.*, **30**, 255-361, 56 fig., 24 pl., Leiden, 1964.
- CRAMER, F.H.: Chitinozoans of a composite section of Upper Llandovery to basal Lower Gedinnian sediments in northern León, Spain. A preliminary report. - *Bull. Soc. belge Géol.*, **75**, 69-129, 7 fig., 5 pl., Brussels, 1967a.
- DUFKA, P.: Lower Silurian Chitinozoans of the Prague Basin (Barrandian, Czechoslovakia). Preliminary Results.- *Rev. Micropaléont.*, **35**, 117-126, 1 fig., 3 tab., 3 pl., Paris, 1992.
- EISENACK, A.: Neotypen baltischer Silur-Chitinozoen und neue Arten. - *N. Jb. Geol. Paläont., Abh.*, **108**, 1-20, 4 fig., pl. 1-3, Stuttgart, 1959.
- EISENACK, A.: Neotypen baltischer Silur-Chitinozoen und neue Arten. - *N. Jb. Geol. Paläont., Abh.*, **114**, 291-316, 8 fig., 1 tab., pl. 14-17, Stuttgart, 1962.
- EISENACK, A.: Mikrofossilien aus dem Silur Gotlands, Chitinozoen. - *N. Jb. Geol. Paläont., Abh.*, **120**, 308-342, 9 fig., 7 tab., pl. 26-30, Stuttgart, 1964.
- EISENACK, A.: Über Chitinozoen des baltischen Gebietes. - *Paläontographica A*, **131**, 137-198, 13 fig., 2 tab., pl. 24-32, Stuttgart, 1968.
- EISENACK, A.: Beiträge zur Chitinozoen-Forschung.- *Paläontographica A*, **140**, 117-130, 1 fig., pl. 32-37, Stuttgart, 1972.
- ELAOUAD-DEBBAJ, Z.: Chitinozoaires Ashgilliens de l'Anti-Atlas (Maroc). - *Geobios*, **17**, 45-48, 5 fig., 3 pl., Lyon, 1984.
- GRAHN, Y.: Chitinozoan stratigraphy in the Ashgill and Llandovery. - In COCKS, L.R.M. & RICKARDS, R.B. (eds.): A Global Analysis across the Ordovician - Silurian boundary.- *Bull. Br. Mus. nat. Hist. (Geol.)*, **43**, 317-323, 27 fig., London, 1988.
- JAGLIN, J.C.: Chitinozoa from the late Ordovician glacio-marine deposits from North Africa. - *Chit. Newsletter*, **8**, 5-6, Uppsala, 1987.
- JENKINS, W.A.M. & LEGAULT, J.A.: Stratigraphic ranges of selected Chitinozoa. - *Palynology*, **3**, 235-264, 6 fig., Dallas, 1979.
- KREUTZER, L.H.: Photo-Atlas of the Variscan Carbonate Sequences in the Carnic Alps (Austria/Italy). - *Abh. Geol. B.-A.*, **47**, 129 p., 9 fig., 3 tab., 46 pl., Wien, 1992.
- KREUTZER, L.H.: Cellon Section. Facial differentiation and bathymetric environment.- In SCHÖNLAUB, H.P. & KREUTZER, L.H. (eds.): IUGS Subcomm. Silurian Stratigraphy, Field Meeting 1994. - *Ber. Geol. B.-A.*, **30**, 85-88, Wien, 1994.
- KRIZ, J.: Silurian Field Excursions. Prague Basin (Barrandian), Bohemia. - *Geol. Ser. Nation. Mus. Wales*, **13**, 111 p., 86 fig., 4 pl., Cardiff, 1992.
- KRIZ, J., DUFKA, P., JAEGER, H. & SCHÖNLAUB, H.P.: The Wenock/Ludlow Boundary in the Prague Basin (Bohemia). - *Jb. Geol. B.-A.*, **136**, 809-839, 18 fig., 1 tab., 3 pl., Wien, 1993.
- KRIZ, J., JAEGER, H., PARIS, F. & SCHÖNLAUB, H.P.: Pridoli - the Fourth Subdivision of the Silurian. - *Jb. Geol. B.-A.*, **129**, 291-360, 44 fig., 1 tab., 6 pl., Wien, 1986.
- LAUFELD, S.: Silurian Chitinozoa from Gotland. - *Fossils and Strata*, **5**, 130 p., 78 fig., Oslo, 1974.
- LAUFELD, S.: Biogeography of Ordovician, Silurian and Devonian Chitinozoans. - In GRAY, J. & BOUCOT, A.J. (eds.): Historical Biogeography, Plate Tectonics, and the Changing Environment, 75-90, 14 fig., (Oregon State University Press), o.O., 1979.
- MOLYNEUX, S.G. & PARIS, F.: Late Ordovician Palynomorphs.- In THUSU, B.T. & OWENS, B. (eds.): The Palynostratigraphy of Northeast Libya. - *J. Micropalaeont.*, **4**, 11-26, 7 pl., London, 1985.
- NESTOR, V. & K.: Correlation of the East-Baltic and Gotland Silurian by Chitinozoans. - In KALJO, D. & KLAAMAN, E. (eds.): Ecostratigraphy of the East Baltic Silurian. - *Acad. Sci. Estonian S.S.R.*, 89-96, 3 fig., Tallinn, 1982.

- NESTOR, V.: Silurian Chitinozoans. - In KALJO, D. & NESTOR, H.N.(eds.): An excursion guidebook, 80-83, fig. 15, pl. 14, 15, Tallinn, 1990.
- NESTOR, V.: Chitinozoan diversity dynamics in the east Baltic Silurian. - *Proc.Estonian Acad.Sci.Geol.*, **41**, 215-224, 5 fig., 2 tab., Tallinn, 1992.
- NESTOR, V.: Early Silurian Chitinozoans of Estonia and North Latvia. - *Academia*, **4**, 163 p., 29 fig., 32 pl., 3 tab., Tallinn, 1994.
- PARIS, F.: Les Chitinozoaires dans le Paléozoïque du sud-ouest de l'Europe. - *Mém.Soc.géol.minéral.Bretagne*, **26**, 412 p., 134 fig., 45 tab., 41 pl., Rennes, 1981.
- PARIS, F.: Chitinozoans. - In HOLLAND, C.H. & BASSETT, M.G. (eds.): A Global Standard for the Silurian System. - *Geol.Ser.Nation.Mus Wales*, **9**, 280-284, fig. 174, 175, Cardiff, 1989.
- PARIS, F.: The Ordovician chitinozoan biozones of the Northern Gondwana Domain. - *Rev. Palaeobot.Palynol.*, **66**, 181-209, 4 fig., Amsterdam, 1990.
- PARIS, F.: Application of chitinozoans in long-distance Ordovician correlations. - In: WEBBY, B.D. & LAURIE, J.R. (eds.): Global Perspectives on Ordovician Geology, 23-33, 3 fig., (Balkema), Rotterdam, 1992.
- PARIS, F.: Evolution paléogéographique de l'Europe au Paléozoïque inférieur: le test de Chitinozoaires. - *C.R.Acad.Sci.Paris*, **316**, Ser.II, 273-280, 4 fig., Paris, 1993.
- PARIS, F.: Chitinozoan Biostratigraphy and Palaeoecology. - In JANSONIUS, J. & MCGREGOR, D.C. (eds.): *Palynology: Principles and Applications*, **vol.2**, 531-552, 9 fig., 3 pl., Salt Lake City, (Publishers Press), 1996.
- PARIS, F. & GRAHN, Y.: Chitinozoa of the Silurian-Devonian boundary sections in Podolia, Ukraine. - *Palaeontology*, **39**, 629-649, 4 fig., 4 pl., London, 1996.
- PARIS, F. & KRIZ, J.: Nouvelles especes de chitinozoaires a la limite Ludlow/Pridoli en Tchécoslovaquie. - *Rev.Palaeobot.Palynol.*, **43**, 155-177, 8 fig., 3 pl., Amsterdam, 1984.
- PARIS, F., LAUFELD, S. & CHLUPÁČ, I.: Chitinozoa of the Silurian-Devonian boundary stratotypes in Bohemia. - *S.G.U., Avh., Ser.Ca.*, **51**, 1-28, 10 fig., 3 pl., Uppsala, 1981.
- PARIS, F. & ROBARDET, M.: Early Palaeozoic palaeobiogeography of the Variscian regions. - *Tectonophysics*, **177**, 193-213, 5 fig., 1990.
- PRIEWALDER, H.: Acritarchen aus dem Silur des Cella-Profiles, Karnische Alpen, Österreich. - *Abh.Geol.-B.-A.*, **40**, 121 p., 39 fig., 24 pl., Wien, 1987.
- PRIEWALDER, H.: Chitinozoans of the Cella Section (Upper Ordovician - Lower Devonian). A Preliminary Report. - In SCHÖNLAUB, H.P. & KREUTZER, L.H. (eds.): IUGS Subcomm.Silurian Stratigraphy, Field Meeting 1994. - *Ber.Geol.B.A.*, **30**, 61-69, 1 fig., Wien, 1994.
- RAUSCHER, R.: Recherches micropaléontologiques et stratigraphiques dans l'Ordovicien at le Silurien en France. Etude des Acritarches, des Chitinozoaires et des Spores. - *Mém.Sci.Géol.*, **38**, 224 p., 46 fig., 31 tab., 12 pl., Strasbourg, 1973.
- SCHÖNLAUB, H.P.: Das Paläozoikum der Karnischen Alpen. - In: Arbeitstagung der Geologischen Bundesanstalt, **1985**, 34-52, fig. 10-15, Wien, 1985.
- SCHÖNLAUB, H.P.: The Ordovician-Silurian boundary in the Carnic Alps of Austria.- In COCKS, L.R.M. & RICKARDS, R.B. (eds.): A Global Analyses of the Ordovician-Silurian boundary. - *Bull.Br.Mus.nat.Hist. (Geol)*, **43**, 107-115, 4 fig., London, 1988.
- SCHÖNLAUB, H.P.: Stratigraphy, Biogeography and Paleoclimatology in the Alpine Paleozoic and its Implications for Plate Movements. - *Jb.Geol.B.-A.*, **135/1**, 381-418, 16 fig., Wien, 1992.
- SCHÖNLAUB, H.P.: The Faunal Relationship of the Silurian of the Alps.- In SCHÖNLAUB, H.P. & KREUTZER, L.H. (eds.): IUGS Subcomm.Silurian Stratigraphy, Field Meeting 1994. - *Ber.Geol.B.A.*, **30**, 52-60, 1 fig., Wien, 1994.
- SCHÖNLAUB, H.P.: Cella Section. Lithology, Paleontology and Stratigraphy.- In SCHÖNLAUB, H.P. & KREUTZER, L.H. (eds.): IUGS Subcomm.Silurian Stratigraphy, Field Meeting 1994. - *Ber.Geol.B.A.*, **30**, 83-84, Wien, 1994.
- SCHÖNLAUB, H.P. & HEINISCH, H.: The Classic Fossiliferous Palaeozoic Units of the Eastern and Southern Alps. - In SCHÖNLAUB, H.P. & KREUTZER, L.H. (eds.): IUGS

- Subcomm.Silurian Stratigraphy, Field Meeting 1994. - Ber.Geol.B.A., 30, 6-51, 18 fig., Wien, 1994.
- SCHWEINEBERG, J.: Silurische Chitinozoen aus der Provinz Palencia (Kantabrisches Gebirge, N-Spanien). - Göttinger Arbeiten zur Geologie und Paläontologie, 33, 94 p., 24 fig., 13 pl., Göttingen, 1987.
- SUTHERLAND, S.J.E.: Ludlow Chitinozoans from the Type Area and Adjacent Regions. - Monogr.Palaeontogr.Soc., Publ.594, 104 p., 57 fig., 18 pl., London, 1994.
- TAUGOURDEAU, P.: Étude de quelques espèces critiques de Chitinozoaires de la Région d'Edjelé et compléments à la faune locale. - Rev.Micropaléont., 6, 130-144, 3 pl., 4 tab., Paris 1963.
- TAUGOURDEAU, P. & DE JEKHOWSKY, B.: Répartition et description des Chitinozoaires Siluro-dévonien de quelques sondages de la C.R.E.P.S., de la C.F.P.A. et de la S.N. REPAL au Sahara - Rév.Inst.Franc.Pétrol., 15, 1199 - 1260, 19 fig., 13 pl., Paris 1960.
- VERNIERS, J.: The Silurian of the Meuse Valley (Brabant Massif, Belgium): Biostratigraphy (Chitinozoa). - Rev.Palaeobot.Palynol., 34, 165-174, 1 fig., 2 pl., Amsterdam 1981.
- VERNIERS, J.: The Silurian Chitinozoa of the Meuse Area (Brabant Massif, Belgium). - Prof.Pap.Geol.Dienst Belgie, 1982/6, 192, 76 p., 10 fig., 9 pl., Gent, 1982.
- VERNIERS, J., NESTOR, V., PARIS, F., DUFKA, P., SUTHERLAND, S. & VAN GROOTEL, G.: A global Chitinozoa biozonation for the Silurian. - Geol.Mag., 132, 651-666, 6 fig., Cambridge, 1995.
- WALLISER, O.H.: Conodonten des Silurs. - Abh.Hess.L.-Amt Bodenforsch., 41, 106 p., 10 fig., 2 tab., 32 pl., Wiesbaden, 1964.
- WRONA, R.: Upper Silurian - Lower Devonian Chitinozoa from the Subsurface of Southeastern Poland. - Palaeont.Polonica, 41, 103-165, 111fig., 13 tab., pl. 24-37, Warsaw, 1980.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte der Geologischen Bundesanstalt](#)

Jahr/Year: 1997

Band/Volume: [40](#)

Autor(en)/Author(s): Priewalder Helga

Artikel/Article: [The Distribution of the Chitinozoans in the Cellon Section \(Hirnantian-Lower Lochkovian\): A Preliminary Report 74-85](#)