


2.2. The Neogene of the Vienna Basin

WERNER E. PILLER



Introduction

The Vienna Basin, located between the Eastern Alps, the West Carpathians and the western part of the Pannonian Basin represents one of the best studied large pull-apart basins of the world (ROYDEN, 1985; WESSELY, 1988). Similar to other European Tertiary Basins (e.g., Paris Basin, London Basin, Mainz Basin) the Vienna Basin was the goal of very early geological studies (e.g., STÜTZ, 1807; PREVOST, 1820; SUCESS, 1885; SCHAFFER, 1907). The importance for hydrocarbon exploration, however, distinctly enhanced our stratigraphic, sedimentologic and tectonic knowledge of the basin during the last 60 years. The different fields of interest studied cover all topics from palaeontology, sedimentology, stratigraphy, tectonics, to natural resources like thermal water and hydrocarbon.

Due to this overall importance the Vienna Basin was the target for several field trips in the course of earth science conferences during the last years. As a result of these activities a variety of field guides were produced (e.g., PILLER & KLEEMANN, 1991; PILLER & VAVRA, 1991; SAUER et al., 1992; PILLER, 1993; PILLER et al., 1996) and the following presentation widely duplicates earlier papers.

Geographical setting

The Vienna Basin is of rhombohedral shape, strikes roughly southwest-northeast, is 200 km long and nearly 60 km wide, and extends from Gloggnitz (Lower Austria) in the SSW to Napajedl in Czekia in the NNE. The western border is bound to the south by the morphological eastern margin of the Northern Alps (represented by several Alpine tectonic units: Greywacke Zone, Northern Calcareous Alps, Flysch Zone) and to the north by the Waschberg Zone. In the east it is bordered in the south by the hills of the Rosaliengebirge, Leithagebirge and the Hainburger Berge, and in the north by the Little Carpathian Mountains; all four hill ranges are part of the Alpine-Carpathian Central Zone. The Vienna Basin is connected with the Little Hungarian Basin via the Hainburger Pforte and with the Eisenstadt Basin via the Wiener Neustädter Pforte. The Eisenstadt Basin has a triangular shape and is bordered in the east by the Ruster Höhenzug, in the north by the Leithagebirge, in the west by the Rosalien- and Leithagebirge, and in the south by the Brennborg. Its maximum dimensions are approx. 20 by 20 km. The subsurface separation from the Vienna Basin is represented by the continuation between the Rosalien- and Leithagebirge; its tectonic and sedimentary history, however, is very similar and the Eisenstadt Basin is therefore considered as a subbasin of the Vienna Basin.

Stratigraphy, facial and tectonic development

The Vienna Basin is part of the Paratethys which formed together with the Mediterranean Sea after vanishing of the Tethys Ocean. Due to its isolated position for most of the time a regional stratigraphic stage system different from that of the Mediterranean had to be established (e. g., RÖGL & STEININGER, 1983; SENES & STEININGER, 1985; STEININGER et al., 1988; STEININGER et al., 1990; RÖGL, 1996; Fig. 14).

Due to the rhombohedral shape and the left-stepping pattern of en-echelon faults firstly ROYDEN (1985) interpreted the basin as pull-apart structure. This idea was strengthened later on (ROYDEN, 1988; WESSELY, 1988), however, based on more profound data a much more complex tectonic evolution was shown by several authors (DECKER et al., 1994; DECKER, 1996; DECKER & LANKREIJER, 1996; DECKER & PERESSON, 1996). The pull-apart mechanism started to act during the Karpatian (STEININGER et al., 1986; SEIFERT, 1992; DECKER, 1996), older sediments (Eggenburgian-Ottangian) at the base of the northern part of the Vienna Basin belong to an earlier piggy-back basin of the Molasse cycle (STEININGER et al., 1986, p. 295, PILLER et al., 1996; DECKER, 1996). Between the Karpatian and Pannonian the subsidence in the central Vienna Basin reached up to 5.5 km (WESSELY et al., 1993). Since the basin is subdivided by a morphological high structure, the Spannberg ridge, into a northern and a southern part, during the Karpatian sedimentation was restricted to the north (north of the Danube) and extended into the south only during the Badenian. Due to the complex fault system the basin was internally highly structured into horst and graben systems. Especially at the western border of the basin, relatively uplifted blocks occur; these are separated from the deep depressions located in the east along major faults (e.g., Mistelbach block along the Steinberg fault in the northern, Mödling block along the Leopoldsdorf fault in the southern basin). The interplay of highly active synsedimentary tectonics with rapid changing trans- and regression cycles (RÖGL & STEININGER, 1983) produced a complex facial pattern inside the basin depending on distance from land and on position of the particular blocks.

The basement of the basin is built by those Alpine-Carpathian nappes bordering the basin also on the surface. The Neogene sediment fill of the basin reaches a thickness of up to 6000 m. At the base mainly clastic sediments are developed representing fluvial facies; occasionally lignite deposits occur (STEININGER et al., 1989). A fully marine development over the entire basin was established only in the Early Badenian (Lower Lagenid Zone). These sediments consist not only of clastics but also carbonates were deposited. This facial development with local coral reefs and widespread coralline algal limestones is restricted to the Badenian. During the Sarmatian, a reduction in salinity already started leading to non-marine and subsequently continental conditions in the Pannonian-Pontian. Although tectonic subsidence was high the basin was rapidly filled due to the short distance to the source of clastic sediments and the basin cycle is therefore limited to the Middle Miocene.

Badenian (16.4. - 13 ma bp)

Due to the major marine transgression at the beginning of the Middle Miocene (RÖGL & STEININGER, 1983, 1984) subtropical biotas entered the Paratethys. In the Vienna Basin conditions for carbonate sedimentation and growth of coral buildups were favourable only during the Badenian stage. Within the context of the meeting the development and facial distribution for this period should be discussed in more detail.

The facial development roughly reflects a distinction between marginal and central basin facies:

Along the basin margins in dependence on the hinterland and coastal morphology the most complex facies pattern is developed. In general siliciclastics and carbonates can be differentiated, both exhibiting a rich facial diversity.

In general, the western border of the southern Vienna Basin is highly influenced by the clastic sediment influx from the Northern Alps. Around the Leithagebirge, which represented an island, a chain of islands or a shoal during the Badenian, and along the Ruster Höhenzug, autochthonous carbonate sediments dominate (irrespective of sometimes thick basal transgressive sediments).

The coastal development along the western margin shows strong fluvial influx at some locations, expressed by thick conglomerates dominated by material derived from the Northern Calcareous Alps as well as the Flysch Zone (Baden [Vöslau] Conglomerate; comp. BRIX & PLÖCHINGER, 1988). In some places, steep rocky shores with large boulders are also preserved (e.g., W' Soof), while wide coastal or marginal areas are covered by sands (Gainfarn Sands) with a rich and excellently preserved fauna. These sands interfinger with the basinal Baden Tegel.




Fig. 2.2.2.: Facial development and stratigraphy of the Vienna Basin with schematic representation of the spannberg ridge in the central part of the basin (after WESSELY, 1988, changed).

The most widespread facies unit along the Leithagebirge and the Ruster Höhenzug as well as at certain sites along the western margins of the Vienna Basin with reduced terrigenous input (e.g., around Wöllersdorf) is the Leitha Limestone. The name of this unit was already established by KEFERSTEIN (1828) and is well known also outside the Vienna Basin. The unit was redefined by PAPP & STEININGER (in:) PAPP et al. (1978) considering the broad facial range and selecting a faciostratotype (comp. Stop 4). The microfacial diversity was worked out by DULLO (1983) into detail describing 10 microfacies types.

Due to its high abundance of coralline red algae this Leithakalk is also well known as Nullipora or Lithothamnium Limestone. Historically important is the first description of a fossil coralline red algae out of this limestone: *Nullipora ramosissima* REUSS 1847. The original material of this taxon was recently rediscovered and the species was assigned to the genus *Lithothamnion* (PILLER, 1994).

In general, the limestone is characterized by the occurrence of coralline algae in various growth forms, ranging from rhodolith dominated types of various growth forms to maërl facies. Coral buildups of limited size are developed only locally. Such buildups are rare along the western margin of the Vienna Basin due to the high terrigenous input and represented only by small patch reefs. Also along the Ruster Höhenzug no significant coral settlement is developed (or preserved); organic buildups are predominantly made up of bivalve beds accompanied, in some places, by corals (comp. DULLO, 1983, p. 37). The best developed coral buildups are present at the southern tip of the Leithagebirge, where the limestones reach the greatest spatial extent and the thickest sequences (about 50 m). Here, due to the island position, no major terrigenous influx restricted coral growth. On the contrary, it can be assumed that water currents or relatively strong waves favoured their growth at the southern tip of the Leithagebirge. The corals are represented mainly by various taxa of *Porites*, accompanied by *Tarbellastraea*, *Caulastrea*, *Acanthastrea*, and *Stylocora* (PILLER & KLEEMANN, 1991).

The basinal facies is characterized by the Baden Tegel, a marl with variable sand and clay content. Intercalated into the marls are sandy layers. This latter material is transported from marginal sources. The marls and sandy interbeddings are highly fossiliferous, containing an extremely rich micro- (foraminifers, ostracods) and macrofauna as well as calcareous nannoplankton (comp. PAPP et al., 1978). The macrofauna is well documented since the 19th century (e.g., D'ORBIGNY, 1846; REUSS, 1849; KARRER, 1861; HÖRNES, 1856, 1870; HÖRNES & AUINGER, 1879) and is represented by solitary scleractinians, brachiopods, decapod crustaceans, molluscs, and fish remains (teeth and otoliths). In the sediments of the Lower Badenian the foraminiferal fauna is extremely rich, containing not only planktic and smaller benthic representatives but in the sandy interbeddings also larger forms as *Amphistegina*, *Planostegina* and *Borelis melo*. Remarkable is the high diversity and good preservation of molluscs (gastropods, bivalves, scaphopods).

The depositional depth of this fine-clastic material can be interpreted as being not deeper than 50 - 100 m (PAPP & STEININGER [in:] PAPP et al., 1978, p. 140) or 100 - 200 m (TOLLMANN, 1985, p. 500). The sandy layers are transported by gravitational transport from marginal areas. Although subsidence of the basin during the Badenian was very rapid, the relatively shallow water depth of the autochthonous sediments can be explained by a high sedimentation rate leading to a sediment accumulation of approx. 1500 m in the central basin during the Badenian (e.g., WESSELY, 1988, p. 342). In the Eisenstadt Basin thickness of the Baden Tegel is distinctly less.

Sarmatian (13 - 11.5 ma bp)

The salinity reduction which already started in the uppermost Badenian continues during the Sarmatian. Salinity decreased generally from 30-17 ‰ and reflects the isolation of the Paratethys from the world oceans. The westernmost extension of the Paratethys during the Sarmatian ended in Lower Austria (near Langenlois). The sedimentological inventory ranges from coastal gravel and sands, to calcareous sandstones ("Atzgersdorfer Stein") and marls (Tegel) exhibiting similarities to Badenian sediments. Additionally, "detrital Leitha Limestone" occurs which mainly represents reworked Badenian Leitha Limestone. The total thickness of Sarmatian sediments surpasses 1000 m in central basin positions. The reduced salinity of the Sarmatian sea caused a low diversity fauna rich in individuals. Stenohaline organisms are nearly absent, some groups (e. g., foraminifers, bryozoa, molluscs) are represented by a few genera only but occur in high densities (for molluscs comp. Stop 5). In contrast to the diverse Badenian algal flora only 2 species of coralline algae were described (KAMPTNER, 1942). These coralline algae produce small buildups in coastal areas together with the sessile foraminifer *Sinzowella caespitosa* (STEINMANN). In some locations also thin serpulid biostromes and ooliths are developed. Based on macro- and microfaunal associations an ecostratigraphic subdivision of the Sarmatian is possible into 5 zones:

Late Sarmatian:	"Verarmungszone"
Middle Sarmatian:	<i>Mactra</i> Beds (= middle <i>Nonion granosum</i> zone) Upper <i>Ervilia</i> Beds (= lower <i>Nonion granosum</i> zone) Lower <i>Ervilia</i> Beds (= <i>Elphidium hauerinum</i> zone)
Early Sarmatian:	<i>Mohrensternia</i> Beds (= <i>Rissoa</i> Beds) (= <i>Elphidium reginum</i> zone)

Pannonian (11.5 - 7.1 ma bp)

After a short regressive phase at the Sarmatian/Pannonian boundary which corresponds to a worldwide regressive tendency and local/regional tectonics the Pannonian Basin became finally isolated from the Eastern Paratethys (STEININGER & RÖGL, 1983, 1984). A following transgression was linked with a further salinity reduction to 5 ‰. In the uppermost part of the Pannonian (Zones F-H) limnic-fluvial conditions already prevailed. In central basin positions sedimentation of marls (Tegel) continued reaching a thickness > 1.500 m; in marginal positions sands and gravelly sediments were deposited.

According to the salinity reduction biotic diversity decreased further compared to the Sarmatian and particularly mollusc faunas are characterized by mass occurrences of a few taxa only which exhibit, however, a fast evolutionary development. Most important taxa are *Melanopsis*, *Congeria*, and *Limnocardium* (Stop 5), whose evolutionary lineages provide the base for the subdivision of the Pannonian into 5 zones (Zone A - E) (PAPP, 1949, 1951, 1953).

References

- ABDELGHANY, O., PILLER, W.E. & TOLEIKIS, R., 1996: Nummulitide Foraminiferen (Gattung: *Planostegina*) im Unter- und Mittel-Miozän der Paratethys und des mediterranen Raumes. - Sediment '96, 11. Sedimentologentreffen, Kurzfassungen, 1, Wien.
- BACHMAYER, F. & TOLLMANN, A., 1953: Die Crustaceen-Fauna aus dem tortonischen Leithakalk (Steinbrüche der Firma Fenk) bei Groß-Höflein im Burgenland. - [in:] KÜPPER, H., EXNER, CH. & GRUBINGER, H. (eds.): Skizzen zum Antlitz der Erde, 308-314, 1 Taf., Wien (Brüder Hollinek).
- BRIX, F. & PLÖCHINGER, B., 1988: Erläuterungen zu Blatt 76 Wiener Neustadt. - 85 pp., 7 Abb., Wien (Geol. B.-A.).
- BRIX, F. & SCHULTZ, O., 1993: Erdöl und Erdgas in Österreich. 2. Auflage. - Veröffentlichungen aus dem Naturhistorischen Museum Wien, N. F. 19, XXIV+688 S., 200 Abb., 17 Beil., Wien.
- DECKER, K., 1996: Miocene tectonics at the Alpine-Carpathian junction and the evolution of the Vienna basin. - Mitt. Ges. Geol. Bergbaustud. Österr. 41: 33-44, Wien.
- DECKER, K. & LANKREIJER, A., 1996: Das Wiener Becken - Tektonik, Subsidenz und Modellierung eines „thin-skinned“ Pull-Apart-Beckens. - Sediment '96, 11. Sedimentologentreffen, Kurzfassungen, 25, Wien.
- DECKER, K. PERESSON, H. & FAUPL, P., 1994: Die miozäne Tektonik der östlichen Kalkalpen: Kinematik, Paläospannungen und Deformationsaufteilung während der "lateralen Extrusion" der Zentralalpen. - Jahrbuch der Geologischen Bundes-Anstalt 137 (1), 5-18, Wien.
- DECKER, K. & PERESSON, H., 1996: Tertiary kinematics in the Alpine-Carpathian-Pannonian System: links between thrusting, transform faulting and crustal extension. - [in:] WESSELY, G., & LIEBL, W., (Ed.): Oil and Gas in Alpidic Thrustbelts and Basins of Central and Eastern Europe, EAPG Spec. Pub. 5, London.
- DECKER, K. & PERESSON, H., 1996: Rollover and hanging-wall collapse during Sarmatian/Pannonian syndimentary extension in the Eisenstadt Basin. - Mitt. Ges. Geol. Bergbaustud. Österr., 41: 45-52.
- DULLO, W.-Ch., 1983: Fossildiagenese im miozänen Leitha-Kalk der Paratethys von Österreich: Ein Beispiel für Faunenverschiebungen durch Diageneseunterschiede. - Facies, 8, 1 -112, 15 Taf., 22 Abb., Erlangen.
- FUCHS, W., 1965: Geologie des Ruster Berglandes (Burgenland). - Jb. Geol. B. A., 108, 155-94, Wien.
- FUCHS, W. & GRILL, R., 1984: Geologische Karte von Wien und Umgebung 1:200.000 - Wien (Geol. B.-A.).
- FUCHS, R. & STRADNER, H., 1977: Über Nannofossilien im Badenien (Mittelmiozän) der Zentralen Paratethys. - Beitr. Paläont. Österr., 2, 1-58, 5 Abb., 8 Taf., Wien.
- GEARY, D.H., 1986: The Evolutionary Radiation of Melanopsid Gastropods in the Pannonian Basin (Late Miocene, Eastern Europe). - unpubl. Ph.D. thesis Harvard Univ., Cambridge, Massachusetts, 1-221.
- HÖRNES, M., 1856: Die fossilen Mollusken des Tertiaer-Beckens von Wien. 1. Band: Univalven. - Abh. k. k. Geol. Reichsanst., 3, 1-736, 1 Karte, Atlas mit 52 Taf., Wien.
- HÖRNES, M., 1870: Die fossilen Mollusken des Tertiaer-Beckens von Wien. 2. Band: Bivalven. - Abh. k. k. Geol. Reichsanst., 4, 1-479, Atlas mit 85 Taf., Wien.
- HÖRNES, R. & AUINGER, M., 1879: Gasteropoden der Meeres-Ablagerungen der ersten und zweiten miocänen Mediterran-Stufe in der Österreichisch-Ungarischen Monarchie. - Abh. k.k. Geol. Reichsanst., 12 (1), 1 - 382, 50 Taf., Wien.
- KAMPTNER, E., 1942: Zwei Corallinaceen aus dem Sarmat des Alpen-Ostrandes und der Hainburger Berge. - Ann. Naturhistor. Mus. Wien, 52 (1941), 5-19, Wien.
- KARRER, F., 1861: Über das Auftreten der Foraminiferen in den marinen Tegeln des Wiener Beckens. - Sitzber. kaiserl. Akad. Wiss., math.-naturwiss. Cl., Abt. I, 427 - 459, 2 Taf., Wien.
- KEFERSTEIN, C., 1828: Beobachtungen und Ansichten über die geognostischen Verhältnisse der nördlichen Kalk-Alpenkette in Österreich-Bayern. - Teutschland geognostisch-geologisch dargestellt, 5/3, 425 pp., Weimar.
- D'ORBIGNY, A., 1846: Die fossilen Foraminiferen des tertiären Beckens von Wien. - XXXVII + 312 S., 21 Taf., Paris (Gide et Comp.)
- PAPP, A., 1949: Fauna und Gliederung des Sarmats im Wiener Becken. - Anz. Akad. Wiss., math.-naturwiss. Kl., 1949 (13), 256-266, 2 Tab., Wien.
- PAPP, A., 1951: Das Pannon des Wiener Beckens. - Mitt. Geol. Ges. Wien, 39-41 (1946-48), 99-193, 7 Abb., 4 Tab., Wien.
- PAPP, A., 1953: Die Molluskenfauna des Pannon im Wiener Becken. - Mitt. geol. Ges. Wien, 44 (1951), 85-222, 1 Abb., 7 Tab., 25 Taf., Wien.
- PAPP, A., 1954: Die Molluskenfauna im Sarmat des Wiener Beckens. - Mitt. Geol. Ges. Wien, 45 (1952): 1-112, Wien.

- PAPP, A., 1978: Die Entwicklung der Heterosteginen im Badenien. - [in:] PAPP, A., CICHA, I., SENES, J. & STEININGER, F. (eds.): *M₄ - Badenien (Moravien, Wielicien, Kosovien)*. - Chronostratigraphie und Neostatotypen. Miozän der Zentralen Paratethys, **6**, 284 - 288, Taf. 12 - 13, Bratislava (Slowak. Akad. Wiss.).
- PAPP, A., MARINESCU, F. & SENES, J. (1974): *M₅. Sarmatien*. - Chronostratigraphie und Neostatotypen, **4**: 1-707, Bratislava (VEDA).
- PAPP, A., CICHA, I. & CTYROKA, J., 1978: Allgemeine Charakteristik der Foraminiferenfauna im Badenien. - [in:] PAPP, A., CICHA, I., SENES, J. & STEININGER, F. (eds.): *M₄ - Badenien (Moravien, Wielicien, Kosovien)*. - Chronostratigraphie und Neostatotypen. Miozän der Zentralen Paratethys, **6**, 263 - 268, Abb. 55 - 57, Bratislava (Slowak. Akad. Wiss.).
- PAPP, A., CICHA, I., SENES, J. & STEININGER, F. (eds.), 1978: *M₄ - Badenien (Moravien, Wielicien, Kosovien)*. - Chronostratigraphie und Neostatotypen. Miozän der Zentralen Paratethys, **6**, 1 - 594, Bratislava (Slowak. Akad. Wiss.).
- PAPP, A. & KÜPPER, K., 1954: The genus *Heterostegina* in the upper Tertiary of Europe. - *Contr. Cushman Found. Foraminif. Res.*, **5**, 108-127, New York.
- PAPP, A., RÖGL, F., CICHA, I., CTYROKA, J. & PISHVANNOVA, S., 1978: Planktonische Foraminiferen im Badenien. - [in:] PAPP, A., CICHA, I., SENES, J. & STEININGER, F. (eds.): *M₄ - Badenien (Moravien, Wielicien, Kosovien)*. - Chronostratigraphie und Neostatotypen. Miozän der Zentralen Paratethys, **6**, 268 - 278, Taf. 1 - 8, Bratislava (Slowak. Akad. Wiss.).
- PAPP, A. & SCHMID, M. E., 1978: Die Entwicklung der Uvigerinen im Badenien der Zentralen Paratethys. - [in:] PAPP, A., CICHA, I., SENES, J. & STEININGER, F. (eds.): *M₄ - Badenien (Moravien, Wielicien, Kosovien)*. - Chronostratigraphie und Neostatotypen. Miozän der Zentralen Paratethys, **6**, 279 - 284, Taf. 9 - 11, Bratislava (Slowak. Akad. Wiss.).
- PAPP, A., JAMBOR, A. & STEININGER, F.F., 1985: *M₆. Pannonien*. - Chronostratigraphie und Neostatotypen, **7**: 1-636, Budapest (Akademiai Kiado).
- PILLER, W.E., 1993: Facies development and coralline algae in the Vienna and Eisenstadt Basins (Miocene). - [in:] HÖFLING, R., MOUSSAVIAN, E. & PILLER, W.E. (eds.): *Facial development of algae-bearing carbonate sequences in the Eastern Alps. Field Trip Guidebook*. - **B7**, 1-24, 6 Fig., 4 Pl., München.
- PILLER, W.E., 1994: *Nullipora ramosissima* REUSS, 1847 - a re-discovery. - *Beiträge Paläontologie*, **19**, 181-189, 2 Fig., 2 Pl., Wien.
- PILLER, W.E. & KLEEMANN, K., 1991: Middle Miocene Reefs and related facies in Eastern Austria. I) Vienna Basin. - VI. International Symposium on Fossil Cnidaria including Archaeocyatha and Porifera, Excursion-Guidebook, Excursion B4, 1-28, 12 Fig., Münster.
- PILLER, W.E. & VAVRA, N., 1991: Das Tertiär im Wiener und Eisenstädter Becken. - [in:] ROETZEL, R. & NAGEL, D. (ed.): *Exkursionen im Tertiär Österreichs, Molassezone-Waschbergzone-Korneuburger Becken - Wiener Becken - Eisenstädter Becken*. - 161-216, Abb. 41-49, Wien (Österr. Paläont. Ges.).
- PILLER, W.E., ABDELGHANY, O. & TOLEIKIS, R., 1995: Über die stratigraphische Verbreitung nummulitider Foraminiferen (Gattung: *Planostegina*) im Unter- und Mittel-Miozän der Paratethys und des mediterranen Raumes. - 2. Tagung der Österreichischen Paläontologischen Gesellschaft, Kurzfassungen der Vorträge und Poster, 25-26, Graz.
- PILLER, W.E., DECKER, K. & HAAS, M., 1996: Sedimentologie und Beckendynamik des wiener Beckens. - *Berichte Geol. B.-A. (Exkursionsführer Sediment '96, 11. Sedimentologentreffen Exkursion A 1)*, **33**, 1-41, Wien.
- PREVOST, C., 1820: *Essai sur la constitution physique et géognostique du bassin de Vienne en Autriche*. - *Journal de physique*, **1820**, 34pp, 2 Abb., Paris.
- REUSS, A.E., 1847: I. Die fossilen Polyparien des Wiener Tertiärbeckens. Ein monographischer Versuch. - *Naturwiss. Abh., Haidinger, W. (ed.)*, **2**, 1 - 109, 11 Taf., Wien.
- REUSS, A.E., 1849: Neue Foraminiferen aus den Schichten des österreichischen Tertiärbeckens. - *Denkschr. K. Akad. Wiss., math.-naturwiss. Cl.*, **1**, 365 - 390, 6 Taf., Wien.
- RÖGL, F., 1996: Stratigraphic correlation of the Paratethys Oligocene and Miocene. - *Mitt. Ges. Geol. Bergbaustud. Österr.* **41**: 65-73, Wien.
- RÖGL, F. & STEININGER, F.F., 1983: Vom Zerfall der Tethys zu Mediterran und Paratethys. Die neogene Paläogeographie und Palinspastik des zirkum-mediterranen Raumes. - *Ann. Naturhist. Mus. Wien*, **85/A**, 135 - 163, 2 Abb., 14 Taf., Wien.
- RÖGL, F. & STEININGER, F.F., 1984: Neogene Paratethys, Mediterranean and Indo-pacific Seaways. - [in:] BRENCHLEY, P.J. (ed.): *Fossils and Climate*, 171-200, Chichester (John Wiley & Sons).
- ROYDEN, L.H., 1985: The Vienna Basin: A thin-skinned pull-apart basin. - [in:] BIDDLE, K. T., CHRISTIE-BLICK, N., (eds.): *Strike slip deformation, basin formation and sedimentation. Society of Economic Paleontologists and Mineralogists, Special Publication 37*, 319-338.

- ROYDEN, L.H., 1988: Late Cenozoic tectonics of the Pannonian basin system. - [in:] ROYDEN, L. H., & HORVATH, F., (eds.): The Pannonian Basin - A study in basin evolution. American Association of Petroleum Geologists, Memoir 45, 27-48, 13 Fig., Tulsa (Oklahoma).
- SAUER, R., SEIFERT, P. & WESSELY, G., 1992: Guidbook to Excursions in the Vienna Basin and the Adjacent Alpine-Carpathian Thrustbelt in Austria. - Mitt. Geol. Ges., 85, 1-264, 200 Abb., Wien.
- SAUERZOPF, F., 1950: Fauna und Gliederung der Schichten des Pannon im südlichen Burgenland. - unpubl. Diss. Univ. Wien, 1-162, Wien.
- SCHAFFER, F.X., 1907: Geologischer Führer für Exkursionen im Inneralpinen Wiener Becken der nächsten Umgebung von Wien. - Samml. geol. Führer, 12, VIII + 127pp, 11 Abb., Berlin (Bornträger).
- SEIFERT, P., 1992: Vienna Basin. [in:] BRIX, F. & SCHULTZ, O., 1993: Erdöl und Erdgas in Österreich. - Veröff. nathist. Mus. Wien, 19, 688pp., Wien.
- SENES, J. & STEININGER, F.F., 1985: I. Mediterranean Tethys and Paratethys Neogene Stages and their current correlation. - [in:] STEININGER, F. F., SENES, J., KLEEMANN, K. & RÖGL, F. (eds.): Neogene of the Mediterranean Tethys and Paratethys. Stratigraphic correlation tables and sediment distribution maps. - vol. 1: 1 - 10, Wien (Inst. Paleontology).
- STEININGER, F.F., 1977: Integrated Assemblage-Zone Biostratigraphy at Marine-Nonmarine Boundaries: Examples from the Neogene of Central Europe. - [in:] KAUFFMANN, E. G. & HAZEL, J. E. (eds.): Concepts and Methods of Biostratigraphy, 235 - 256, 11 Text-Fig., Stroudsburg, Pennsylvania (Dowden, Hutchinson & Ross).
- STEININGER, F.F., SENES, J., KLEEMANN, K. & RÖGL, F. (eds.), 1985: Neogene of the Mediterranean Tethys and Paratethys. Stratigraphic correlation tables and sediment distribution maps. - vol. 1: XIV + 189 pp.; vol. 2: XXVI + 536 pp., Wien (Inst. Paleontology).
- STEININGER, F.F., WESSELY, G., RÖGL, F. & WAGNER, L., 1986: Tertiary sedimentary history and tectonic evolution of the Eastern Alpine Foredeep. - Giornale di Geologia, ser. 3, 48/1-2, 285-297, 10 Fig., Bologna.
- STEININGER, F.F., MÜLLER, C. & RÖGL, F., 1988: Correlation of Central Paratethys, Eastern Paratethys, and Mediterranean Neogene Stages. - [in:] ROYDEN, L.H. & HORVATH, F. (eds.): The Pannonian System. A study in basin evolution. - Amer. Assoc. Petrol. Geol. Mem. 45, 79 - 87, 3 Fig., Tulsa (Oklahoma).
- STEININGER, F.F., RÖGL, F., HOCHULI, P., & MÜLLER, C., 1989: Lignite deposition and marine cycles. The Austrian Tertiary lignite deposits - A case history. - Sitzungsbericht der Akademie der Wissenschaften Wien, mathematisch-naturwissenschaftliche Klasse 197 (5-10), 309-332.
- STEININGER, F.F., BERNOR, R.L. & FAHLBUSCH, V., 1990: European Neogene marine/continental chronologic correlations. - [in:] LINDSAY, E. H., FAHLBUSCH, V. & MEIN, P. (eds.): European Neogene Mammal Chronology. - 15 - 46, 1 Fig., New York (Plenum Press).
- STÜTZ, A., 1807: Mineralogisches Taschenbuch. Enthaltend eine Oryctographie von Unterösterreich etc. - 394pp, Wien-Triest (Geistinger).
- SUESS, E., 1885: Das Antlitz der Erde. Bd. 1. - 778 S., illustr. (Tempisky), Prag (Freytag), Leipzig.
- TOLLMANN, A., 1955: Das Neogen am Nordwestrand der Eisenstädter Bucht. - Wissenschaftliche Arbeiten Burgenland, 10, 1-75, 7 Fig., Profile A-G, 1 geol. Karte, Eisenstadt.
- TOLLMANN, A., 1985: Geologie von Österreich. Band II. Außerzentralalpiner Anteil. - 710 S., 287 Abb., Wien (Deuticke).
- WESSELY, G., 1988: Structure and Development of the Vienna Basin in Austria. - [in:] ROYDEN, L. H. & HORVATH, F. (eds.): The Pannonian System. A study in basin evolution. - Amer. Assoc. Petrol. Geol. Mem. 45, 333 - 346, 10 Fig., Tulsa (Oklahoma).
- WESSELY, G., KRÖLL, A., JIRICEK, R., & NEMEC, F. 1993: Wiener Becken und angrenzende Gebiete - Geologische Einheiten des präneogenen Beckenuntergrundes. - Geologische Themenkarte der Republik Österreich 1:200.000, Wien (Geol. B.-A.).

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte der Geologischen Bundesanstalt](#)

Jahr/Year: 1999

Band/Volume: [49](#)

Autor(en)/Author(s): Piller Werner E.

Artikel/Article: [The Neogene of the Vienna Basin 11-19](#)