

## **Environmental change during the PETM drives formation of gigantic biogenic magnetite**

Dirk SCHUMANN<sup>1, 2, \*</sup>, Timothy D. RAUB<sup>3</sup>, Robert E. KOPP<sup>4</sup>, Jean-Luc GUERQUIN-KERN<sup>5, 6</sup>, Ting-Di WU<sup>5, 6</sup>, Isabelle ROUILLER<sup>2, 7</sup>, Aleksey V. SMIRNOV<sup>8</sup>, S. Kelly SEARS<sup>2, 7</sup>, Uwe LÜCKEN<sup>9</sup>, Sonia M. TIKOO<sup>3</sup>, Reinhard HESSE<sup>1</sup>, Joseph L. KIRSCHVINK<sup>3</sup> & Hojatollah VALI<sup>1, 2, 7</sup>

<sup>1</sup>Department of Earth and Planetary Sciences, McGill University, 3450 University Street, Montréal, Québec, H3A 2A7, Canada; <sup>2</sup>Facility for Electron Microscopy Research, McGill University, 3640 University Street, Montréal, Québec, H3A 2B2, Canada; <sup>3</sup>Division of Geological and Planetary Sciences, California Institute of Technology, MC 170-25, 1200 E California Blvd., Pasadena, CA 91125, USA; <sup>4</sup>Department of Geosciences and Woodrow Wilson School of Public & International Affairs, 210 Guyot Hall, Princeton University, Princeton, NJ 08544, USA; <sup>5</sup>INSERM, U759, Institut Curie, Imagerie intégrative de la molécule à l'organisme, Orsay, 91405, France; <sup>6</sup>Institut Curie, Laboratoire de Microscopie Ionique, Orsay, 91405, France; <sup>7</sup>Department of Anatomy and Cell Biology, McGill University, 3640 University Street, Montréal, Québec, H3A 2B2, Canada; <sup>8</sup>Department of Geological and Mining Engineering and Sciences, Michigan Technological University, Houghton, MI 49931-1295, USA; <sup>9</sup>FEI Company, Nanobiology Marketing, Eindhoven, 5600KA Eindhoven, The Netherlands; hesse@eps.mcgill.ca.

About 55 Ma at the Paleocene /Eocene boundary, the planet experienced a 5-9°C jump in global temperatures within less than 10,000 years known as the Paleocene-Eocene Thermal Maximum (PETM). Debate is still ongoing on what triggered the event that lasted ~180,000 years. However, several lines of evidence suggest that large releases of greenhouse gases, in particular methane from gas hydrates, contributed to the rapidity and extent of the warming event. The event reflects a drastic perturbation of the Earth's ocean and atmospheric systems and was associated with a significant diversification of the terrestrial fauna and flora but also of marine life. Numerous deep-sea benthic foraminifera species disappeared and new forms evolved.

The sediments deposited during the PETM serve as archives that contain distinct paleontological, mineralogical, magnetic, chemical and isotopic evidence of these climatic changes. Kopp et al., (2007) and Lippert & Zachos (2007) report an extraordinary magnetofossil 'Lagerstätte' in kaolinite-rich clay sediments deposited during the PETM at subtropical paleolatitude in the Atlantic Coastal Plain of New Jersey. They used ferromagnetic resonance (FMR) spectroscopy, other rock magnetic methods, and transmission electron microscopy (TEM) of magnetic separates to characterize sediments from boreholes at Ancora (ODP Leg 174AX) and Wilson Lake, NJ, respectively. These sediments contain abundant ~40- to 300-nm cuboidal, elongate-prismatic and bullet-shaped magnetofossils, sometimes arranged in short chains, resembling crystals in living magnetotactic bacteria. Aside from abundant bacterial magnetofossils, these same sediments also contain exceptionally large and novel biogenic magnetite crystals unlike any previously reported from living organisms or from sediments (Schumann et al., 2008).

The spearhead-like, spindle-like and elongated hexaoctahedra magnetite crystals exhibit chemical composition, lattice perfection and oxygen isotopic composition consistent with a biogenic origin. The spearheads and spindles can be up to 4000 nm long (8 times larger than magnetite produced by magnetotactic bacteria). The elongated hexaoctahedra may be up to 1400 nm long and are thus "giant" magnetofossils. They are probably too big to be produced intracellularly by prokaryotes, although exceptionally large prokaryotes having cellular diameters up to 750 µm have been reported. In a few cases, we observed apparently intact, tip-outward spherical assemblages of spearhead-like particles that possibly represent the preserved original biological arrangement of these crystals in a hitherto unknown magnetite producing organism.

The discovery of these exceptionally large biogenic magnetite crystals that possibly represent the remains of new micro-organisms that appeared and disappeared with the PETM sheds some light upon the ecological response to biogeochemical changes that occurred during the warming event. Magnetotactic bacteria usually live in the oxic-anoxic transition zone of fresh, brackish, and marine environments including the suboxic zone of

sediments. The occurrence of these new forms together with conventional magnetofossils suggests that they shared a similar ecological niche. The development of a thick suboxic zone with high iron bioavailability – a product of dramatic changes in weathering and sedimentation patterns driven by severe global warming – may have resulted in diversification of magnetite-forming organisms, likely including eukaryotes.

In this study we extended the search for these new magnetofossils to other PETM locations of the Atlantic margin and to a possible modern analog environment in the Amazon delta system.

Kopp, R.E., Raub, T.D., Schumann, D., Vali, H., Smirnov, A.V., Kirschvink, J.L. (2007): Magnetofossil spike during the Paleocene-Eocene thermal maximum: Ferromagnetic resonance, rock magnetic, and electron microscopy evidence from Ancora, New Jersey, United States. *Paleoceanography*, doi:10.1029/2007pa001473.

Lippert, P.C. & Zachos, J.C. (2007): A biogenic origin for anomalous fine-grained magnetic material at the Paleocene-Eocene boundary at Wilson Lake, New Jersey. *Paleoceanography*, doi:10.1029/2007pa001471.

Schumann D et al., (2008): Gigantism in unique biogenic magnetite at the Paleocene-Eocene Thermal Maximum. *Proc. Nat. Acad. Sci*, 105:17648–17653.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte der Geologischen Bundesanstalt](#)

Jahr/Year: 2009

Band/Volume: [78](#)

Autor(en)/Author(s): Schumann Dirk, diverse

Artikel/Article: [Environmental change during the PETM drives formation of gigantic biogenic magnetite 40-41](#)