

## Lower – Middle Devonian conodont faunas from Prague Basin

BERKYOVÁ, S.<sup>1,2</sup>, FRÝDA, J.<sup>1,3</sup> & SUTTNER, T.J.<sup>4</sup>

(1) Czech Geological Survey, P.O.B. 85, 11821 Praha 11, Czech Republic; *berkyova.s@seznam.cz*

(2) Charles University in Prague, Faculty of Science, Albertov 6, 12843 Praha 2

(3) Faculty of Environmental Sciences, CULS, 16521 Praha 6, Czech Republic; *frйда@cgu.cz*

(4) Austrian Academy of Sciences (CPA) c/o University of Graz, Institute of Earth Sciences (Geology and Palaeontology), Heinrichstrasse 26, A-8010 Graz, Austria; *thomas.suttner@uni-graz.at*

The conodonts documented were sampled from Třebotov Limestone (Daleje-Třebotov Formation, Lower – Middle Devonian, upper Emsian – lowermost Eifelian) and Choteč Limestone (Choteč Formation, Middle Devonian, Eifelian) and their shallow-water equivalents (Suchomasty and Acanthopyge Limestone). The following sections were studied: Prastav Quarry, Holyně section, Na Škrábku Quarry, Barrandov roadcut section, Jelínek mill section, Červený Quarry at Suchomasty, U Němců section at Karlštejn and Na Vyhliďce section at Hostím (Fig. 1).

Třebotov Limestone (Daleje – Třebotov Formation, Lower – Middle Devonian) was defined by SVOBODA & PRANTL (1947) and redefined by CHLUPÁČ (1959). This unit is represented mainly by bioturbated, skeletal wacke- to packstones with high component content, mainly dacryoconarid tentaculites, abundant fragments of ostracods, trilobite exoskeletons, nautiloids and goniatites, gastropods, brachiopods and bivalves.

On the basis of the presence of micritic matrix, benthic fauna typical for muddy bottom environments, abundant representatives of plankton and the lack of sedimentological features that would indicate current activity, the sedimentary environment is interpreted as low-energy, relatively deep water milieu which is rich in free oxygen (judging from intense bioturbation and diverse benthic assemblages) and below storm wave base.

Choteč Limestone (Choteč Formation, Middle Devonian), a unit defined by SVOBODA & PRANTL (1948) and redefined by CHLUPÁČ (1957, 1959), reflects in its development the effect of the Basal Choteč Event, which has been regarded as an important global transgressive geo-event documented by a distinct facies and faunal change (e.g. CHLUPÁČ & KUKAL 1986, HOUSE 2002). This unit is represented by dark, organic-rich, well-washed peloidal grainstones and packstones. Characteristic feature of the grainstones is intense micritization of grains however, echinoderm ossicles as the main component could be identified. These allochthonous parts, representing gravity flow deposits (turbidite) from shallow water alternate with background sediments, which are dark lime-mudstone/wackestone with distinct lamination at some parts suggestive of lack of free oxygen at the sea bottom.

Conodonts from the Lower and Middle Devonian (*gronbergi* – *kockelianus* biozones) in the Prague Basin (Fig. 2) were previously studied by WEDDIGE & ZIEGLER (1977), KLAPPER (1977), KLAPPER, ZIEGLER & ZIKMUNDOVÁ in CHLUPÁČ et al. (1977), KLAPPER et al. (1978), ZIKMUNDOVA in CHLUPÁČ et al. (1979), ZUSKOVÁ (1991) and KALVODA in HLADIL & KALVODA (1993).

The conodont fauna present is of very low diversity and also abundance, spanning six biozones (*serotinus* – *kockelianus* biozone). The following genera and species were recovered (numbers in brackets stand for quantity of specimens): *Polygnathus serotinus* (375), *P. linguliformis bultyncki* (147), *Icridous beckmanni beckmanni* (11), *I. beckmanni sinuatus* (2), *Ozarkodina carinthiaca* (4), *P. cooperi cooperi* (8), *P. costatus patulus* (36), *P. costatus partitus* (62), *P. costatus costatus* (53), *P. sp. aff. P. trigonicus* (4), *Polygnathus linguiformis pinguis* (2), *Polygnathus trigonicus* (3), *Tortodus kockelianus australis* (2), *T. kockelianus kockelianus* (11), *Polygnathus pseudofoliatulus* (16), *Pelekysgnathus sp.* (1), *Pandorinellina sp.* (2), *Pseudooneotodus beckmanni*, *Polygnathus cf. benderi* (2), *P. cf. borealis* (1) and three new species, which are together with two last mentioned taxa and with *P. serotinus* and *P. I. bultyncki* topic of a recent manuscript.

**References:**

- CHLUPÁČ, I. (1957): Stratigraficko – paleontologický výzkum břidlic dalejských a vápenců hlubočepských ve střeďočeském devonu. - *Věstník Ústředního Ústavu Geologického*, 32: 248-258.
- CHLUPÁČ, I. (1959): Faciální vývoj a biostratigrafie břidlic dalejských a vápenců hlubočepských (Eifel) ve střeďočeském devonu. - *Sborník Ústředního Ústavu Geologického*, 25: 445-511.
- CHLUPÁČ, I., LUKEŠ, P. & ŽIKMUNDOVÁ, J. (1977): Barrandian 1977, A field trip Guidebook, Field Conference of the International Subcommittee on Devonian Stratigraphy: 1-23.
- CHLUPÁČ, I. & KUKAL, Z. (1986): Reflection of possible global Devonian events in the barrandian area, C.S.S.R., - *In: WALLISER, O.H. (Ed.): Lecture Note in Earth Sciences, Global Bio-events: 169-179.*
- CHLUPÁČ, I., LUKEŠ, P. & ŽIKMUNDOVÁ, J. (1979): The Lower/Middle Devonian boundary beds in the Barrandian area, Czechoslovakia. - *Geologica et Palaeontologica*, 13: 125-156.
- HLADIL, J. & KALVODA, J. (1993): Extinction and recovery successions of the Devonian marine shoals; the Eifelian – Givetian and Frasnian – Famennian events in Moravia and Bohemia. - *Bulletin of the Czech Geological Survey*, 68(4): 13-23.
- HOUSE, M.R. (2002): Strength, timing, setting and cause of mid-Palaeozoic extinctions. - *Palaeogeography, Palaeoclimatology, Palaeoecology*, 181: 5-25.
- KLAPPER, G. (1977): Lower-Middle Devonian boundary conodont sequence in the Barrandian area of Czechoslovakia. *Časopis pro mineralogii a geologii*, 22(4): 401-410.
- KLAPPER, G., ZIEGLER, W. & MASHKOVA, T.V. (1978): Conodonts and correlation of Lower-Middle Devonian boundary beds in the Barrandian area of Czechoslovakia. - *Geologica et Palaeontologica*, 12: 103-116.
- SVOBODA, J. & PRANTL, F. (1947): O stratigrafii a tektonice staršího paleozoika v okolí Třebotova. - *Sborník Státního Geologického Ústavu*, 14: 281-314.
- SVOBODA, J. & PRANTL, F. (1948): O stratigrafii a tektonice staršího paleozoika v okolí Chýnice. - *Sborník Státního Geologického Ústavu*: 1-39.
- WEDDIGE, K. & ZIEGLER, W. (1977): Correlation of Lower/Middle Devonian beds. - *Newsletter on Stratigraphy*, 6(2): 67-84.
- ZUSKOVÁ, J. (1991): Conodont faunas from the Lower/Middle Devonian section in Praha-Barrandov. - *Věstník Ústředního Ústavu Geologického*, 66(2): 107-112.




Fig. 1: Conodont distribution chart across the boundary of Třebotov Limestone and Choteč Lst.




Fig. 2: A, B. *Polygnathus costatus costatus* KLAPPER, 65x; C. *Polygnathus pseudofoliatus* WITTEKINDT, 57x; D. *Polygnathus costatus patulus* KLAPPER, 60x; E. *Polygnathus cooperi cooperi* KLAPPER, 40x; F, G. *Polygnathus costatus partitus*, KLAPPER, ZIEGLER & MASHKOVA, 60x; H. *Polygnathus trigonicus* BISCHOFF & ZIEGLER, 85x; I. *Polygnathus serotinus* TELFORD, 57x; J. *Tortodus kockelianus kockelianus* (BISCHOFF & ZIEGLER), 85x; K. *Polygnathus linguiformis bulyyncki* WEDDIGE, 57x.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte der Geologischen Bundesanstalt](#)

Jahr/Year: 2009

Band/Volume: [79](#)

Autor(en)/Author(s): Berkyova Stanislava, Fryda Jiri, Suttner Thomas

Artikel/Article: [Lower-Middle Devonian conodont faunas from Prague Basin 9-11](#)