

Coupling of marine and continental isotope records during the Eocene/Oligocene transition

**Melanie J. Bugler¹, Stephen T. Grimes¹, Jerry J. Hooker², Margaret E. Collinson³,
Gregory D. Price¹, Nathan D. Sheldon⁴, Paul A. Sutton¹**

¹ SoGEES, University of Plymouth, Drake Circus, Plymouth, Devon, PL4 8AA, UK.

² Dept. of Palaeontology, Natural History Museum, London, SW7 5BD, UK.

³ Dept. of Earth Sciences, Royal Holloway Univ. of London, Egham, TW20 0EX, UK.

⁴ Dept. of Geological Sciences, University of Michigan, Ann Arbor, MI 48109, USA.

During the Cenozoic, conditions existed that allowed for the formation of permanent continental-scale ice caps on Antarctica (Miller *et al.*, 1991; Zachos *et al.*, 2001). Long-term cooling commenced after the Early Eocene Climatic Optimum (52 to 50 Ma), culminating in the Oi-1 glacial maximum (~33.65 Ma) (Miller *et al.*, 1991; Zachos *et al.*, 2001; Bohaty and Zachos, 2003). This event reflects a transition from a greenhouse world to an icehouse world and is recorded in the marine realm as a stepped positive $\delta^{18}\text{O}$ shift in benthic and planktonic foraminifera (Coxall *et al.*, 2005; Katz *et al.*, 2008; Pearson *et al.*, 2008; Lear *et al.*, 2008).

Reported here is a $\delta^{18}\text{O}$ record across the continental Eocene/Oligocene transition (EOT).

The record derived from *V. lentus* displays 3 cycles followed by a positive $\delta^{18}\text{O}$ shift across Oi-1 with subsequent recovery to pre-Oi-1 values. Within these cycles, plus across the isotope shift and recovery, $\delta^{18}\text{O}$ values change in magnitude between 1.4–1.8‰. If entirely due to temperature this would equate to a 6–13°C (depending on calibration) temperature variation, which according to its high magnitude, implies a contribution from another factor, such as changes in the isotopic composition of the host water in which *V. lentus* grew. The ultimate control on this is likely to be related to changes in global ice volume. The *V. lentus* $\delta^{18}\text{O}$ Oi-1 shift and recovery are comparable to that seen in the marine record. Furthermore, in cycle 1 the most positive $\delta^{18}\text{O}$ value corresponds with the Late Eocene Event as observed in the marine record. Cycles 2 and 3 together occupy the same time interval, recording successive positive $\delta^{18}\text{O}$ shifts in the marine realm. Moreover, the negative $\delta^{18}\text{O}$ peak between cycles 1 and 2, associated with biotic and isotopic evidence for warming, matches a marine calcite compensation depth shoaling event. Although the cycle 2 positive $\delta^{18}\text{O}$ peak has no equivalent in the marine realm, fluctuations within cycle 3 appear synchronous with the named events EOT1/Step 1 and EOT2 (Coxall *et al.*, 2005; Katz *et al.*, 2008; Pearson *et al.*, 2008; Lear *et al.*, 2008). Therefore, the $\delta^{18}\text{O}$ record from *V. lentus* in a coastal floodplain environment shows substantial similarities with those from benthic foraminifera in the marine realm across the EOT and Oi-1 in timing and trajectory. Combined with multiproxy isotope derived summer temperatures, sea-level change and biotic data, some patterns of climate change with respect to the relative roles of temperature and ice-volume in this continental record are also comparable with those documented in the marine record. This suggests that when sampling resolution allows, a coupling of the marine and continental isotope records through the EOT can be observed.

References:

Bohaty, S. M. & Zachos, J. C., 2003. *Geology* 31, 1017-1020. Coxall, H. K., *et al.*, 2005. *Nature* 433, 53-57. Hooker, J.J. *et al.*, 2009. Chapter 12: In *The Late Eocene Earth—Hothouse, Icehouse, and Impacts* (eds. Koeberland, C. & Montanari, A). The Geological Society of America Special Paper, 452. Katz, M. E., *et al.*, 2008. *Nature Geoscience* 1, 329-334. Lear, C. H., *et al.*, 2008. *Geology* 36, 251-254. Miller, K. G., *et al.*, 1991. *Journal of Geophysical Research-Solid Earth and Planets* 96, 6829-6848. Pearson, P. N., *et al.*, 2008. *Geology* 36, 179-182. Zachos, J., *et al.*, 2001. *Science* 292, 686-693.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte der Geologischen Bundesanstalt](#)

Jahr/Year: 2011

Band/Volume: [85](#)

Autor(en)/Author(s): Bugler Melanie J., Grimes Stephen T., Hooker Jerry J., Collinson Margaret E., Price Gregory D., Sheldon Nathan D., Sutton Paul A.

Artikel/Article: [Coupling of marine and continental isotope records during the Eocene/Oligocene transition](#) 48