

**A new planktonic foraminifera species
(*Hantkenina gohrbandti* nov. spec.)
from the Middle Eocene of the northwestern Tethys
(Mattsee, Austria)**

Fred Rögl¹, Hans Egger²

¹ Museum of Natural History, Burgring 7, A-1010 Vienna, Austria;

² Geological Survey of Austria, Neulinggasse 38, A-1030 Vienna, Austria;

The planktonic foraminifer genus *Hantkenina* is characterized by planispiral coiling and hollow chamber extensions, called tubulospines. It evolved gradually from the genus *Clavigerinella*, which shows radial elongate, clavate or digitate chambers, but no tubulospines. This evolutionary trend and the transition from *Clavigerinella* to *Hantkenina* was demonstrated from the Austrian Holzhäusl section (Coxall et al., 2003) and from the Kilwa drill sites in Tanzania (Pearson et al., 2004). At both localities, a newly discovered species, which has been named *Hantkenina singanoae* by Coxall and Pearson (2006), was considered to be the missing link between the two genera. However, the chambers of this species are terminate in a distal hood (proto-tubulospine), and it appears unclear how, and unlikely that, straight tubulospines of the younger *Hantkenina* species could evolve from this bent feature.

Rögl & Egger (2010) reported on a newly discovered planktonic foraminifer, which is characterized by pointed chamber ends with a nub (proto-tubulospines) and in some cases by the first tubulospines appearing in a juvenile growth stage. This species forms the evolutionary link between the genera *Clavigerinella* and *Hantkenina* and is considered to be the real ancestor of the genus *Hantkenina*. For this species the name *Hantkenina gohrbandti* nov. spec. is introduced (Rögl & Egger, in press). The new species is named in honour of Klaus H. Gohrbandt (Gulf Breeze, Florida, USA; former employee of Rohöl-Aufsuchungs AG, Vienna) for his fundamental work on the Paleogene of the Helvetikum north of Salzburg.

References:

- Coxall, H.K., Huber, B.T. and Pearson, P.N., 2003. Origin and morphology of the Eocene planktonic foraminifer *Hantkenina*. *Journal of Foraminiferal Research*, 33, 237-261.
- Coxall, H.K. and Pearson, P.N., 2006. Taxonomy, biostratigraphy, and phylogeny of the Hantkeninidae (*Clavigerinella*, *Hantkenina*, and *Cribohantkenina*). *Cushman Foundation Special Publication*, 41, 213-256.
- Rögl, F. and Egger, H., 2010. The missing link in the evolutionary origin of the foraminiferal genus *Hantkenina* and the problem of the Lower/Middle Eocene boundary. *Geology*, 38, 23-26.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte der Geologischen Bundesanstalt](#)

Jahr/Year: 2011

Band/Volume: [85](#)

Autor(en)/Author(s): Rögl Fred, Egger Hans

Artikel/Article: [A new planktonic foraminifera species \(*Hantkenina gohrbandti* nov.spec.\) from the Middle Eocene of the northwestern Tethys \(Mattsee, Austria\) 140](#)