Fluid inclusion in quartz from the Aligudarz Granitoids, NW Iran

Bagheriyan, Siyamak

Islamic Azad University, Tiran Branch, Isfahan, 85318, I.R. Iran

The Aligudarz granitoid plutons intruded into meta-sediments in the Sananday-Sirjan zone, NW of Aligudarz city (west Iran). Based on petrography and geochemistry, the Aligudarz granitoid has been classified as granite, granodiorite and S-Type pegmatite with the mineral assemblage of quartz, plagioclase, Kfeldspar, biotite and muscovite. The intrusion has thermally metamorphosed the country rocks up to the albite-epidote-hornfels facies. Fluid inclusions studies in the quartz veins of the Aligudarz granitoid show four types of fluid inclusions (Fig.1): 1) low saline aqueous inclusions; 2) high saline inclusions (Fig. 2); 3) CO₂-H₂O rich inclusions; and 4) carbonic inclusions. The density data obtained from the fluids indicate entrapment temperatures of 580 to 636 °C at pressures of 3.9 to 5.1 Mbars. These conditions nearly coincide with the obtained mineral P-T estimations. There are partial melting processes during the formation of the S-Type Aligudarz granitoids.




Fig. 2. Fluid inclusion in a quartz vein of the Aligudarz granitoid (high saline inclusion)




Fig. 1. a-b: three-phase fluid inclusions (L+V+S); c: two phase fluid inclusions (L+V), rich of liquid phase; d: two-phase fluid inclusions (L+V), rich of vapour phase; e: two-phase fluid inclusions (L+V)including two phases of non-mixed vapour; f: liquid single phase fluid inclusion (L) and two phase inclusion (L+V); g: single phase of fluid inclusion (V); H: multi-phase fluid inclusion (L+V+ha+sil+opg)

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Berichte der Geologischen Bundesanstalt

Jahr/Year: 2011

Band/Volume: 87

Autor(en)/Author(s): Bagheriyan Siyamak

Artikel/Article: Fluid inclusion in quartz from the Aligudarz Granitoids, NW Iran 30