

Genesis of the Weishan REE deposit in the Shandong Province, eastern China: evidences from Rb-Sr isochron age, LA-MC-ICPMS Nd isotopic composition and fluid inclusions

Fan, Hong-Rui*, Lan, Ting-Guang*, Hu, Fang-Fang* and Yang, Kui-Feng*

*Key Laboratory of Mineral Resources, Institute of Geology and Geophysics, Chinese Academy of Sciences, Beijing 100029, PR China

Weishan REE deposit, a pegmatite-type REE deposit related to alkaline rocks, is located at Luxi Block, south-eastern North China Craton. According to muscovite Rb-Sr isochron, it formed at the age of 119.5 Ma, which belonged to the early Cretaceous large-scale mineralization in the North China Craton. LA-ICPMS Nd isotopic compositions of bastnaesite and monazite indicate that the source of the REE minerals is the enriched lithospheric mantle. Observation of fluid inclusions in quartz, fluorite and barite shows that four types of inclusions can be identified, including (1) H₂O inclusions, (2) pure CO₂ inclusions, (3) H₂O+CO₂ inclusions and (4) H₂O+CO₂+daughter mineral inclusions. The H₂O inclusions are secondary inclusions while others are primary inclusions. The daughter minerals in H₂O+CO₂+daughter mineral inclusions include thenardite, barite, celestine, calcite, apthitalite and glauberite. The homogenization temperature and capture pressure of H₂O+CO₂ and H₂O+CO₂+daughter mineral inclusions range from 205 °C to 433 °C and 120 MPa to 200 MPa, respectively. Coupled with the existence of abundant daughter minerals and sulphur stable isotopic compositions, it can be deduced that the initial ore-forming fluids were high-temperature, moderate-pressure and high-concentration orthomagmatic fluids, which were characterized by enrichment of HCO₃⁻/CO₃²⁻, SO₄²⁻ and F⁻ and multicomponent (e.g., Na⁺, K⁺, Ca²⁺, Ba²⁺, Sr²⁺ and REE³⁺). The coexistence of C, HC and HCD inclusions and the wide range of liquid/vapour ratios between these inclusions suggest that fluid unmixing may have occurred during ore-forming process. REE were most probably transported as REEF²⁺ and precipitated through fluid boiling. Fluids mixing, which

contributed little to the REE precipitation, also happened in the late stage of the ore-forming process.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte der Geologischen Bundesanstalt](#)

Jahr/Year: 2011

Band/Volume: [87](#)

Autor(en)/Author(s): Fan Hong-Rui, Lan Ting-Guang, Hu Fang-Fang, Yang Kui-Feng

Artikel/Article: [Genesis of the Weishan REE deposit in the Shandong Province, eastern China: evidences from Rb-Sr isochron age, LA-MC-ICPMS Nd isotopic composition and fluid inclusions 88](#)