

Bicarbonate-rich fluid inclusions and hydrogen diffusion in quartz gangue from the Libčice orogenic gold deposit, Bohemian Massif

Zacharias, Jiri*, Hrstka, Tomas* and Dubessy, Jean**

*Institute of Geochemistry, Mineralogy and Mineral Resources, Faculty of Science, Charles University in Prague, Albertov 6, Prague, Czech Republic

**G2R(UMR 7566), Faculté des Sciences, Université Henri Poincaré-Nancy Université, Vandoeuvre-les-Nancy Cedex, France

Unusual paleofluid composition is reported for the Libčice orogenic-type gold deposit located in a contact zone of the Central Bohemian Plutonic Complex, Czech Republic. Unexpected bicarbonate-rich fluids and their complex chemistry variations characterize primary fluid inclusions from the main gold-bearing quartz vein. A detailed microthermometry, Laser Raman Micro Spectroscopy and SEM cathodoluminescence study was used in order to decipher fluid history. The results (Zacharias, 2002; Hrstka et al., 2011) indicate the presence of H₂O and H₂O–CO₂–CH₄ (± N₂; H₂S) fluids, the latter displaying variations of the CO₂/CH₄ ratio in the gaseous phase from 6.8 to 0.06. Variation of the CH₄ content across single grains and between different levels of the mine was recorded. The presence of nahcolite, H₂ (up to 6 mole%) and ethane (0–0.2 mole%) in the fluids were also discovered by Raman probe. Potential models for the formation of different types of fluids present in the deposit are discussed, including the genesis of HCO₃⁻ rich fluids as well as H₂ and C₂H₆ presence in the primary fluid inclusions. The potential influence of organic matter-bearing sediments, as well as the impact of the intrusion of CBPC, re-equilibration and/or re-speciation of fluid inclusions during the post-entrapment history is considered to have the main impact on the complex paleofluid chemistry. Based on the thermodynamic modelling, H₂ diffusion into the fluid inclusions was shown to be the main reason for the CH₄ variation on the scale of a single grain, as well as across the whole vein. Although the exact processes of production/formation of HCO₃⁻ and H₂ at the Libčice deposit remain open to discussion, reactions in the C–O–H system are considered to be a possible formation mechanism.

This work also contributes to our understanding of the importance of post-entrapment modifications and reactions in the C–O–H system on interpretation/deciphering the processes in orogenic-type deposits.

This research benefited from financial support of the Czech Republic (GACR 205/06/0702 and MSM 0021620855) and French government (BGF 2005 and EC-HPMT-CT-2001, rf. 00381). We have also benefited from international collaboration within the scope of the UNESCO-IGCP project No. 540 "Fluid inclusions in orogenic Au deposits

REFERENCES

- Zacharias J. (2002): J. Czech Geol. Society, 47: 123-132.
Hrstka T., Dubessy J., Zacharias J. (2011): *Chem. Geol.* 281: 313-332.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte der Geologischen Bundesanstalt](#)

Jahr/Year: 2011

Band/Volume: [87](#)

Autor(en)/Author(s): Zacharias Jiri, Hrstka Tomas, Dubessy Jean

Artikel/Article: [Bicarbonate-rich fluid inclusions and hydrogen diffusion in quartz gangue from the Libcice orogenic gold deposit, Bohemian Massif 208](#)