Berichte Geol.B.-A., 94 (ISSN 1017-8880) - 18. Jahrestagung ÖPG Linz (12.-13. 10. 2012)

## Computed tomography in palaeontology – case studies on Triassic and Cretaceous cephalopods

ALEXANDER LUKENEDER<sup>1</sup>, SUSANNE LUKENEDER<sup>1</sup> & CHRISTIAN GUSENBAUER<sup>2</sup>

<sup>1</sup>Natural History Museum Vienna, Department of Geology & Palaeontology, Burgring 7, 1010 Vienna, Austria; E-mail: alexander.lukeneder@nhm-wien.ac.at <sup>2</sup>Upper Austria University of Applied Sciences, Campus Wels, Stelzhamerstrasse 23, 4600 Wels, Austria

Case studies on computed tomography on ammonites, ammonite mass-occurrences, and trace fossils, deposited during the Upper Triassic (approx. 225 my) of Turkey and during the Lower Cretaceous of Italy (approx. 130 my), are presented. X-ray computed tomography is known in palaeontology as providing data for 3D visualization and geometrical modelling techniques. Computed tomography down to a few microns (or even below) of spatial resolution are increasingly employed for geoscientific investigations, using an equally variable range of processing techniques and software packages. Additionally, internal structures are visualized without destruction of fossils, as computed tomography is a non-destructive method.

#### Experimental set-up

The scans were made at the Upper Austria University of Applied Science in Wels with a dual source industrial 3D computed tomography device (RayScan 250 E), equipped with a 225 kV microfocus and a 450 kV minifocus X-ray tube as well as a 2048x2048 pixel flat panel detector (cone beam reconstruction). The data consist of volumetric pixels (voxels), whose size limits the detail detectability. For each fossil part the optimal voxel size and tube voltage were set according to the specimen dimensions.

#### Case study 1: Triassic ammonites from Turkey (FWF Project P22109-B17)

A case study in computed tomography on the ammonite genus *Orthoceltites* is presented. The latter studies are essential for palaeontology and systematic investigations. Ammonite shells and filled phragmocones (secondary calcite) from the *Orthoceltites* beds possess the same mass-density as the matrix in which the ammonite specimens are embedded. The almost identical mass-density of the embedding matrix (about 2.8 g/cm<sup>3</sup>), the ammonite shells (secondary calcite, about 2.6-2.8 g/cm<sup>3</sup>) and the infilled matrix (about 2.8 g/cm<sup>3</sup>) avoids their visualization. It is therefore not possible to visualize the ammonites by computed tomography. In few cases ammonite shells, body chambers and secondary formed calcite fissures can be observed in computed tomographic images and movies. Future work will be done on the possibilities of computed tomography in such dense Mesozoic limestones.

### Case study 2: Cretaceous ammonites and trace fossils from Italy (FWF Project P 20018-N10)

This case is somehow different compared with the Triassic case study above. We used the same methods and equipment as within the Triassic samples. The only differences are the sediments and the material of ammonites and trace fossils. Within marly-limestones from the Lower Cretaceous numerous ammonites (e.g., *Dissimilites, Lytoceras*; Lower Barremian) and trace fossils (*Halimedides*; Lower Barremian) are preserved as limonitic steinkerns or limonitic fillings. These dense structures can be visualized by computed tomography. New morphological details as spines on ammonite shells, shape and position of suture lines, and the exact structure of trace fossils can be shown.

# **ZOBODAT - www.zobodat.at**

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Berichte der Geologischen Bundesanstalt

Jahr/Year: 2012

Band/Volume: 94

Autor(en)/Author(s): Lukeneder Alexander, Lukeneder Susanne, Gusenbauer Christian

Artikel/Article: <u>Computed tomography in palaeontology - case studies on Triassic and</u> <u>Cretaceous cephalopods. 16</u>