

## *Philonthus roeri* sp. n. from Okavango, Namibia (Coleoptera, Staphylinidae)

Horace Last

During his entomological expeditions to the Namibian part of the Okavango in January of 1988 Dr. H. Roer collected in light traps several species of *Philonthus* (Staphylinidae). Among them I found a new species which I name after the collector of the type specimen.

### *Philonthus roeri* sp. n.

Description: Head and antennae, with the exception of the basal segment, black. Pronotum clear red, elytra ferrugineous often suffused with black to varying degrees,




Fig. 1: Paratype of *Philonthus roeri* sp. n.; left, habitus, right, scutellum and elytra enlarged to show punctures and hair cover. SEM-photo by G. Oleschinski.

abdomen and legs ferrugineous. Head a little longer than broad, slightly converging to anterior corners, with two rows each of six punctures.

Elytra a little longer than broad, very finely and very densely punctured. Scutellum finely punctured.

Abdomen densely punctured and pubescent, parallel-sided.

Legs spinose.

Length 5 mm.

Holotype: ♂, South West Africa (Namibia), Nyangana/Okavango, 14.—22. 1. 1985 (H. Roer). 85 paratypes, same date. In the Zoologisches Forschungsinstitut und Museum Alexander Koenig, Bonn.

Remarks: Very similar in colouration and size to *Philonthus novellus* Last, but the punctures on the elytra are smaller and denser, the abdominal segments are mostly ferrugineous and shorter.

#### Literature

Last, H. R. (1952): New Species of Staphylinidae (Col.) from Africa. — Ent. mon. Mag. 88: 89—92.

Horace Last, Woodville, Hillside Walk, Storrington, W. Sussex, U. K.