Bird Collections – an Essential Resource for Collecting Ectoparasites, in Particular Chewing Lice

Eberhard MEY

Naturhistorisches Museum of the Thüringer Landesmuseum Heidecksburg Rudolstadt

Abstract. On study skins of birds, prepared in traditional manner (i.e. without washing), one can find traces (often in considerable numbers) of their former ectoparasitical occupants. Besides mites and ticks, it is above all the chewing lice (Insecta, Phthiraptera: Amblycera and Ischnocera) that remain in mummified form on skins. Their eggs can be found in particular feather tracts. This fact has been known at least since the beginning of the 19th century and has been well exploited by many collectors, especially those interested in chewing lice. The material gathered by this method forms a basis of our current knowledge of the occurrence and distribution of chewing lice that parasitize birds of all orders. This method, briefly outlined here, docs of course contain sources of error. The authenticity of the material is high if contamination with parasites from other hosts can be excluded. Despite this, the technique of shaking out study skins remains an effective way of studying the diversity of these host-specific insects, most of which are only 2-4 mm in size. At present, c. 1200 species of Amblycera and c. 2700 of Ischnocera are known from birds. The number of undescribed chewing lice forms is estimated to be around 4000. Chewing lice are normally highly host specific. Therefore we may expect that skins of extinct bird species will yield specimens of also extinct chewing lice forms, and occasionally other host-specific ectoparasites. For these reasons bird collections possess a high importance for scientific disciplines outside ornithology

Key words. bird collections, chewing lice, collecting technique, extinct phthirapteran taxa

1. INTRODUCTION

Bird study-skin collections have not been assembled by museums in order to later examine the specimens for the occurrence of ectoparasites. For this purpose, the examination of live, freshly collected and/or prepared birds is always more rewarding, and especially regarding the authenticity of the study material it is fundamentally the preferred method. In the relatively straightforward spectrum of ornithophilous ectoparasites (a diverse array of mite groups, ticks, bugs, louse-flies, fleas), the feather mites (Acari: Analgoidea) and in particular the chewing lice (Insecta, Phthiraptera: Amblycera and Ischnocera, which alone will be treated here), occupy a special place because of their commonness. This applies only to these two groups even if they are collected from skins in scientific collections. For at least 200 years this situation has been exploited by the small band of dedicated feather lice specialists. At present, and for the foreseeable future, this remains an indispensable and effective way of harvesting from the tremendously concentrated resource of bird skin collection material that is generally difficult to obtain. The need for research is as great as ever. Considering the present expanding interest in systematics and biodiversity, well catalogued bird collections are gaining an importance which, if not exactly new is at least greater than in the past. This importance is here examined in detail. I know from my own experience over many years (Table 1) that the hidden potential of these traditional repositories in the form of 'accidentally' con-

 Table 1: Bird study-skin collections partially examined since 1978, mostly by the author, in order to gather chewing lice material.

Basel, Naturhistorisches Museum	Halberstadt, Museum Heineanum
Berlin, Naturkundemuseum	Halle/Saale, Zoologisches Institut
Bonn, Museum Alexander Koenig	Hamburg, Zoologisches Institut und Museum
Bremen, Übersee-Museum	Karlsruhe, Landessammlungen für Naturkunde
Dresden, Staatliches Museum für Tierkunde	Leiden, Nationaal Natuurhistorisch Museum Naturalis
Funchal, Museu Municipal (Hist. Nat.), Madeira	Leipzig, Naturkundemuseum
Frankfurt am Main, Senckenberg-Museum	Munich, Zoologische Staatssammlung
Görlitz, Naturkundemuseum	Rudolstadt, Naturhistorisches Museum
Gotha, Museum der Natur	Stuttgart, Museum für Naturkunde
Greifswald, Zoologisches Institut	Wrocław, Zoologisches Institut

E. Mey



Fig. 1: Ischnoceran chewing louse, Megathellipeurus parkeri (Emerson & PRICE), shaken from the over 100-yearold study skin of a Malleefowl Leipoa ocellata in the Berlin Naturkundemuseum. During collecting it often happens that bristles, legs, antennac, and/ or other parts of the brittle mummified insects break off.

Photo: H. Reubke.

served parasitical arthropod species is often ignored by many museums, despite the recognition of their important status as bird collections.

2. CHEWING LICE – FASCINATING INSECTS

Chewing lice (Fig. 1) are insects on average 2-4 mm in length, which are highly adapted to life on the integument of their host, and which never leave these hosts (a single species) during their brief lives; in other words they are extremely host-specific. They are highly unlikely to be absent on any recent bird species, even if they arc as yet completely unknown for some families (e.g. Todidae, or Picathartidac and some other passerines). This extreme host specificity of chewing lice has led to the realization that their evolution has paralleled that of their hosts, at least since the end of the Cretaceous Period, resulting in cospeciations. This hypothesis is utilized by parasitophyletic studies in the search for evidence useful in the clarifying of taxonomic relationships between particular host groups.

Chewing lice belong to the order of the animal lice Phthiraptera and can be separated into two distinct suborders. The Amblycera (1183 species so far identified) and the Ischnocera (2683 species) are known from all avian orders (CRUICKSHANK et al. 2001). I estimate that the number of undescribed chewing lice forms is around 4000.

Against this background, chewing lice deserve special attention, particularly on the part of museum ornithologists who should be aware of the scientifically valuable material that is 'slumbering' in the bird studyskin collections in their care in the inconspicuous form of mummified chewing lice.

3. EXAMINATION AND TREATMENT OF BIRD STUDY SKINS

It is a mistaken assumption that following the death of a host all its chewing lice attempt to leave the corpse. If the opportunity to transfer to another host is lacking, the majority are forced to stay where they are. They are able to survive for some days but eventually die. The Ischnocera remain on the fcathers that are tightly gripped in their powerful mandibles, while the Amblycera are close to the skin of the host, their thick coating of bristles ensuring that they are fairly well anchored in the body plumage. Both of these situations however do not necessarily apply to all members of the chewing lice population on a host individual. Many of the Amblycera actively move to leave the host, a behaviour that perhaps explains why considerably more Ischnocera than Amblycera are recorded on study skins. Nevertheless, experience shows that despite the loss of part of the Ischnocera and/or Amblycera fauna caused by the preparation procedure, a profitable fraction usually remains on the (unwashed) skin.

The methodology of the removal of ectoparasites from bird study skins has been described by EICHLER in several publications, especially in his work of 1971. It is a simple process and as long as the skins are responsibly treated is highly recommended (Figs.



Fig. 2: Wing feathers and tail feathers are moved lightly against each other and the body feathers loosened with the fingers. This usually causes the first mummified ehewing lice to fall from the plumage. Gentle shaking or patting of the skin will increase this number. The head plumage should be examined feather by feather, and also for ehewing lice eggs (nits), using tweezers.

Photos: W. Kluh



Fig. 3: The mummified chewing lice are placed in a meticulously labelled collecting tubule using a fine paintbrush. The dark-coloured adults can be more or less easily picked out among the precipitation' of dust, powder, detritus, tow, and often single feathers. Those lice that are whitish in colour and less than 1 mm in size have to be searched for with particular care.

Photo: W. Kluh

2-4). There are two stages in the operation and they can be carried out separately or in combination:

- 1. The purposeful feather-by-feather examination of specific areas, particularly on the head including the base of the bill, using tweezers; this is sometimes done to firstly establish the presence of eggs (nits) in order to be sure that there is a chance of finding chewing lice on the skin (Fig. 5). This intensive searching method is suggested for small birds up to about *Turdus merula* size.
- 2. The careful patting of the skin after the plumage has been loosened. This is the best method for species roughly larger than *Turchus merula*. This extensive examination can be carefully repeated several times with one skin.

The study skins thus treated are held over a smooth white sheet of paper or plastic and the resulting mummified chewing lice preserved in dry collecting tubules.



Fig. 4: Utensils used in obtaining ectoparasites, principally chewing lice, from dry bird study skins. - Photo: E. MEY.

When labelling, it is of crucial importance that the correct host name is recorded (EICHLER 1970). One occasionally comes across finds from study skin collections whose authenticity has to be doubted (e.g., when shorebird chewing lice are found on a pigeon). Therefore when evaluating material obtained from study skins it is always necessary to employ a skeptical approach and not to jump to hasty conclusions.

Fig. 5: The confirmation of nits, usually in the head plumage, is a surc indication of chewing lice infestation. An unusual case is shown here, where the cggs and clutch of Hohorstiella ectootoca MEY are attached to the dorsal feather tips of a Blue-headed Quail-dove Starnoenas cyanocephala and are thus externally visible. After MEY (1984).



4. SOME RESULTS FROM BIRD STUDY SKIN EXAMINATIONS

Just how successful examination of skin material carried out using the combined 'shaking out and patting' method can be has been impressively demonstrated, quantitatively as well as qualitatively, by WARD (1957 a,b) in the case of the tinamous. It has to be said that the chewing louse fauna of the Tinamiformes, compared with all other orders of recent birds, has by far the highest diversity at ca. 200 species. This was confirmed by WARD (l.c.) inasmuch as he was able to obtain from several skins of one Crypturellus subspecies alone 12 Ischnocera species, and a maximum of 9 from only a single skin of the same subspecies. He found several thousand mummified chewing lice on a total of 1500 skins of 130 tinamou species and subspecies. Of the 20 chewing lice genera identified (using the most conservative classification), 18 infest only tinamous. According to WARD's (1957a) findings, the host distribution of the chewing lice recorded does reflect the classification of the Tinamidae by von BOETTICHER (1934) in Tinaminae, Rhynchotinae, and Eudromiinae. In addition, he was able to make fundamental taxonomic and ecological statements concerning tinamou chewing lice.

However, it is only with a great amount of effort that we will be in a position to present similar results for

well over half of all bird species, in particular for the Passeriformes. In this order the intensity and extent of infestation is much less, as well as varying in some species a great deal geographically, and many chewing lice are lost because they fall from the skins during frequent handling. Despite this a search for chewing lice can be productive, especially when nits can be identified and mummified adults extracted from where they are still clinging and/or from deep in the plumage. I examined a series of 3 study skins of the world's smallest bird, the Bumble-bee-sized Hummingbird Mellisuga helenae, and discovered on head and breast of one of them two imagines and one larva of Trochiloecetes cubanensis MEY. Three of 10 Chlorostilbon ricordii skins from Cuba held nits, but adults (of Trochiloecetes or Trochiliphagus) could not be found despite a one hour search (MEY 1982).

For years an important concern has been the examination of skins of extinct birds. In addition to the posthumous discovery of *Huiacola extinctus* there have been further such records of chewing lice species, some still unnamed (MEY 1990, Table 2).

5. DISCUSSION

The number of experienced and productive animal louse specialists, those with years of research on the taxonomy and systematics of these insects behind them, has always been tiny compared with the overwhelming number of forms they have to deal with. There are perhaps about two dozen such experts worldwide, so that in most countries there is no-one working on the faunistics and/or taxonomy of chewing lice or of any other animal louse group. Chewing and sucking lice will presumably never enjoy the popularity of butterflies, beetles, orchids, or even birds! The 2nd International Congress on Phthiraptera in Brisbane, Australia, in July 2002 will provide an overview of the present state of research and will hopefully give fresh impetus to our work. Similarly, the results of the ongoing project 'Fauna Europaea' will doubtless lead to a more intense study of those chewing louse forms occurring in Europe.

Against this background our first priority must be to concentrate on and encourage specific projects:

- Establishment of a survey of the existing collections of chewing lice, or other animal lice, e.g. in Germany or Europe.
- Examination of birds for chewing lice, immediately before their preparation, by a taxidermist and/or curator and/or other trained personnel. Collected examples that have not been properly prepared and have lain untreated for years, either dried or in alcohol, could soon be useless as study specimens.
- Examination of those bird skins whose study for mummified chewing lice has not been rendered pointless by preparation processes like the 'unfortunately' modern practice in the past few years of washing them.
- Thorough investigation of all available study skins of bird species now extinct (see table 2) and of those that have always existed, or now exist, in small populations. This would result in material towards a chewing lice Red Data List.

ACKNOWLEDGEMENT

I thank Brian Hillcoat (Neuss) for the translation of the manuscript.

Chewing lice species	Host species
Amblycera	And triaglon (PECHETERS, 1911)
Ischnocera	Ara Incolor (BECHSTEIN, 1811)
Brueelia sp., 1 Columbicola extinctus Malcomsen, 1937* Halipeurus raphanus Timmermann, 1961 Huiacola extinctus Mey, 1990 Rallicola sp. 1, 1, 1 Rallicola sp. 11, 1, 2 Saussurites defectus Tendeiro, 1969	Chloridops kona Wilson, 1888 Ectopistes migratorius (Linné, 1766) Oceanodroma macrodactyla Bryant, 1887 Heteralocha acutirostris (Gould, 1837) Porzana palmeri (Frohawk, 1892) Nesoclopeus poecilopterus (Hartlaub, 1866) Ectopistes migratorius (Linné, 1766)

Table 2: Chewing lice collected from skins of extinct bird species. From literature and unpublished records by the author.

* According CLAYTON & PRICE (1999) this species is not extinct and lives on Columba fasciata SAY in North and South America.

LITERATURE

- Boetticher, H. v. (1934): Beiträge zu einem phylogenetisch begründeten natürlichen System der Steisshühner (Tinami) auf Grund ihrer taxonomisch verwertbaren Charaktere. Jena Z. Naturwiss. **69**: 169-192.(13)
- Clayton, D. H. & Price, R. D. (1999): Taxonomy of New World Columbicola (Phthiraptera: Philopteridae) from the Columbiformes (Aves), with Descriptions of Five New Species. Ann. Entomol. Soc. Am. **92**: 675-685.
- CRUICKSHANK, R. H., JOHNSON, K. P., SMITH, V. S. ADAMS, R. J., CLAYTON, D. H. & PAGE, R. D. M. (2001): Phylogenetic analysis of Partial Sequences of Elongation Factor 1a Identifies Major Groups of Lice (Insecta: Phthiraptera). Molec. Phyl. Evol. 19: 202-215.
- EICHLER, W. (1970): Artangabe, Wirtsangabe und Wirtsspezifität bei Ektoparasiten. Beitr. Vogelkd. 16: 72-86.
- EICHLER, W. (1971): Vogelbäige-Abklopfen als Mallophagen-Sammelmethode. Angew. Parasitol. **12**: 38-52.
- MEY, E. (1982): Kubanische Mallophagen I. Reichen-

bachia 20: 213-218.

- MEY, E. (1984): Kubanische Mallophagen II. Reichenbachia 22: 151-161.
- MEY, E. (1990): Eine neue ausgestorbene Vogel-Ischnozere von Neuseeland, *Huiacola extinctus* (Insecta, Phthiraptera). Zool. Anz. 224: 49-73.
- WARD, R. A. (1957a): A study of the host distribution and some relationships of Mallophaga parasitic on birds of the order Tinamiformes Part I. Ann. entomol. Soc. Am. 50: 335-353.
- WARD, R. A. (1957b): A study of the host distribution and some relationships of Mallophaga parasitic on birds of the order Tinamiformes Part II. Ann. entomol. Soc. Am. 50: 452-459.

Dr. Eberhard MEY, Naturhistorisches Museum of the Thüringer Landesmuseum Heidecksburg Rudolstadt, Schloßbezirk 1, D-07407 Rudolstadt, Germany email: museum@heidecksburg.de

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Bonn zoological Bulletin - früher Bonner Zoologische Beiträge.

Jahr/Year: 2003

Band/Volume: 51

Autor(en)/Author(s): Mey Eberhard

Artikel/Article: <u>Bird Collections - an Essential Resource for Collecting Ectoparasites, in</u> <u>Particular Chewing Lice 131-135</u>