

# Scientific note

[urn:lsid:zoobank.org/pub:AA5B7777-C36B-4E9C-B77E-272FBCBA3D7A](https://nslid:zoobank.org/pub:AA5B7777-C36B-4E9C-B77E-272FBCBA3D7A)

## Things are not always as they seem: High-resolution X-ray CT scanning reveals the first resin-embedded miniature gecko of the genus *Ebenavia*

Juan D. Daza<sup>\*1</sup>, Jordan P. Hunziker<sup>1</sup>, Aaron M. Bauer<sup>2</sup>, Philipp Wagner<sup>2,3</sup>, Wolfgang Böhme<sup>4</sup>

<sup>1</sup> Department of Biological Sciences, Sam Houston State University, 2000 Avenue I, Life Sciences Building, Room 105, Huntsville, TX 77341, USA

<sup>2</sup> Department of Biology, Villanova University, 800 Lancaster Avenue, Villanova, PA 19085, USA

<sup>3</sup> Allwetterzoo Münster, Sentruper Str. 315, D-48161 Münster, Germany

<sup>4</sup> Zoologisches Forschungsmuseum A. Koenig, Adenauerallee 160, D-53113 Bonn, Germany

\*Corresponding author: E-mail: [juand.daza@gmail.com](mailto:juand.daza@gmail.com)

<sup>1</sup> [urn:lsid:zoobank.org/author:7F796C4B-B0D5-4669-8D6F-96CA494B62F5](https://nslid:zoobank.org/author:7F796C4B-B0D5-4669-8D6F-96CA494B62F5)

<sup>2</sup> [urn:lsid:zoobank.org/author:A54A16B8-BEE1-4996-9170-6CADB539B5D1](https://nslid:zoobank.org/author:A54A16B8-BEE1-4996-9170-6CADB539B5D1)

<sup>3</sup> [urn:lsid:zoobank.org/author:0575CE45-AB5C-4493-A8AD-DD399E18284A](https://nslid:zoobank.org/author:0575CE45-AB5C-4493-A8AD-DD399E18284A)

<sup>4</sup> [urn:lsid:zoobank.org/author:FFAC2972-9F52-404B-BA9C-489C7793FF8D](https://nslid:zoobank.org/author:FFAC2972-9F52-404B-BA9C-489C7793FF8D)

\* [urn:lsid:zoobank.org/author:78BB0EB0-8A48-4A1D-8D53-287AD1EF5FFA](https://nslid:zoobank.org/author:78BB0EB0-8A48-4A1D-8D53-287AD1EF5FFA)

**Abstract.** We identify a presumed specimen of *Sphaerodactylus* in amber from the Zoological Research Museum Alexander Koenig as being embedded in copal, rather than amber. Further, the specimen matches the morphology not of a Hispaniolan gecko, but of the extant Madagascan species *Ebenavia boettgeri*, which occurs in a known area of copal deposits.

**Key words.** *Sphaerodactylus*, *Ebenavia*, CT scan, Madagascar, Osteology.

Fossil lizards embedded in amber are frequently spectacular since they preserve, in high definition, the three dimensionality of ancient organisms. To date, fossil remains of squamates have been found in six amber deposits around the world (Daza et al., 2016). Examples include the oldest reptile in amber (*Baobdasaurus xenurus*) from the Early Cretaceous of Lebanon (Arnold et al., 2002); some scales attributable to a squamate from the Albian of France (Perrichot and Néraudeau, 2005); 14 fossil squamates from the mid-Cretaceous of Myanmar, including members with affinities to Iguania, Gekkota, Scincoidea, Anguimorpha, and Ophidia (Arnold & Poinar, 2008; Daza et al., 2016; Fontanarrosa et al., 2018; Daza et al., in press; Xing et al., 2018); a gekkotan and numerous lacertids in Baltic amber (*Succinilacerta succinea*, Böhme & Weitschat, 1998; Borsuk-Białynicka et al., 1999; *Yantarogekko balticus*, Bauer et al., 2005; see also Černaňský & Augé, 2013); and many lizards from the Miocene deposits of Mexico and Hispaniola classified in the genus *Anolis* (*A. electrum*, Lazell, 1965; Rieppel, 1980; de Queiroz et al., 1998; Polcyn et al., 2002; Castañeda et al., 2014; Sherratt et al., 2015) and *Sphaerodactylus* (*S. dommeli*, Böhme, 1984; *S. ciguapa*, Daza & Bauer, 2012). Copal specimens have received relatively less attention and were reviewed in Broschinski & Kohring

(1998). The genera *Phelsuma*, *Lygodactylus*, and *Geckolepis* have been preserved in Madagascan copal.

The study of lizards in amber has been facilitated by the use of X-rays and High-Resolution X-ray computed tomography (HRCT; Polcyn et al., 2002; Daza et al., 2013; Castañeda et al., 2014; Sherratt et al., 2015; Daza et al., 2016), allowing the rendering of the skeleton without distortion, in addition to providing incredible integumentary detail. As part of an ongoing research project, we examined all available gecko specimens in Miocene amber from Hispaniola preserved in amber using HRCT. One specimen was revealed to be neither a *Sphaerodactylus* nor embedded in Miocene amber. The specimen was scanned at the Center for Nanoscale Systems, Harvard University using a Nikon (Metris) X-Tek HMXST 225 scanner with a molybdenum target at 70kV, 135 µA, 1000 ms exposure, 3143 projections, 0.1° rotation step, and no filter. The reconstructed voxel size for the particular specimen was 14.251 µm. The original data set has been archived and is available to the public at Morphosource ([https://www.morphosource.org/Detail/ProjectDetail/Show/project\\_id/545](https://www.morphosource.org/Detail/ProjectDetail/Show/project_id/545)). Additional specimens for comparison were scanned at UTCT | The University of Texas High-Resolution CT Facility in a Xradia – Zeiss machine. The specimens were scanned with a 4X objective,



Fig. 1. Specimen ZFMK 94000, in dorsal view. Scale bar equals 5 mm.

70kV/10W using variable parameters. These specimens are part of a large data base of skull Micro-CT that includes nearly all gekkotan genera (Aaron M. Bauer digital collection). All post-processing of the scan data was

performed using Avizo Lite 9.5.0 (©FEI SAS, Thermo Fisher Scientific, 2018).

The specimen in question, from the Alexander Koenig Research Museum (ZFMK 94000, Fig. 1), had been obtained by the museum sponsoring society (Alexander-Koenig-Gesellschaft) in November 2012 under the assumption that it was a piece of Dominican amber with a *Sphaerodactylus* (Gekkota: Sphaerodactylidae) inclusion. Although the size and overall appearance is consistent with that of these miniaturized geckos (Daza et al. 2008), a more thorough analysis of this material and comparison with the Micro-CT data base of gekkotans indicated that this specimen is a Madagascan clawless gecko of the genus *Ebenavia* in the family Gekkonidae. Morphological data indicates that the specimen is a sub-fossil, and that the resin is copal, not amber.

Copal from Madagascar is botanically assigned to the fabacean species *Hymenaea verrucosa* (Penney et al., 2005). Copal can be differentiated from mature resins, such as amber, with Raman spectroscopic analyses, showing more intense bands at around  $1640\text{ cm}^{-1}$  due to more stretching vibrations of the  $\nu(\text{C}=\text{C})$  attributed to the olefinic group ( $\text{C}=\text{CH}_2$ ; Winkler et al., 2001). Thermal analyses have also been used to characterize resins; copal from Madagascar may be differentiated from amber and other copal resins in reaching a peak in differential thermogravimetric analysis at  $384^\circ\text{C}$ , while Colombian copal and amber from other localities peaks at  $400^\circ\text{C}$  or more (Ragazzi et al., 2003). The age of copal resins may be only a few hundred to up to four million years old. Some resins from Madagascar have been dated using carbon dating analyses to be as young as just a few decades (Poinar, 1999; Bosselaers et al., 2010). Other estimates





Fig. 2. Distributional map of *Ebenavia* species. Colors follow Hawlitschek et al. (2018). Additional localities for *E. maintimainty* taken from Nussbaum and Raxworthy (1998).




**Fig. 3.** Gecko in copal (*Ebenavia boettgeri*, ZFMK 94000) **A.** right lateral view, **B.** dorsal view; **C.** ventral view; **D.** left lateral view indicating the major bones. Abbreviations c#, cervical vertebrae #, l#, left toe #, r#, right toe #, t#, thoracic vertebrae #.

for the age of Madagascan copal include a range from Holocene to Recent (10,000–100 y; Schlüter & Gnielinski, 1987; Lourenço, 1996; Winkler et al., 2001). In this study, we confirmed that the specimen is embedded in copal based on the morphological similarities with modern species, and some simple tests on the resin: 1) A hot needle was pushed into the piece, causing rapid melting at the point of insertion (rather than slow melting expected in amber); the melting resin released a mild fragrance

(amber yields a sooty odor). 2) Under a UV lamp the piece did not show any color change (rather than emitting a bluish glow, as does amber).

Morphological comparisons considering members of nearly all described gekkotan genera indicate that *Ebenavia* shares most morphological characters with ZFMK 94000. The genus occurs on Madagascar and satellite islands, Pemba Island, Grand Comoro, Mohéli, Anjouan, Mayote, Nosy Bé, Nosy Komba, Île Sainte-Ma-





**Fig. 4.** HRCT of the skulls of three geckos. Sphaerodactylidae: **A.** *Sphaerodactylus semasiops* (MCZ R-55766); Gekkonidae: **B.** *Ebenavia boettgeri* (ZFMK 94000), and **C.** *Ebenavia boettgeri* (CAS 66195).

rie (Nosy Boraha), Nosy Mangabe, and Île aux Prunes (Nosy Alaïña), as well as Mauritius (Ramanamanjato et al., 2002; Hawlitschek et al., 2017, 2018; Uetz et al., 2018). Until recently *Ebenavia* included only two species (*E. maintimainty* and *E. inunguis*). *Ebenavia maintimainty* has a restricted range, being found in Toliara Province in southwestern Madagascar (Nussbaum & Raxworthy, 1998), while the more widespread *E. inunguis* was recently split into four new species (viz., *E. boettgeri*,

*E. robusta*, *E. safari*, *E. tuelinae*; Hawlitschek et al., 2018, Fig. 2). Copal deposits in Madagascar are concentrated in the northern part of the island (i.e., Cap D'Ambré and the Sava Region, Geirnaert 2002), which is compatible with the distribution of *E. safari* and *E. boettgeri* (Hawlitschek et al., 2018).

The specimen is embedded in a cone-shaped piece of orange resin (Fig. 1). The preservation is exceptional, conserving the complete anterior half of the body. It ap-




Fig. 5. Lateral view of the snout of ZFMK 94000, inset shows the separation between the rostral scale and the nostril.

pears desiccated, but the skeleton is in perfect condition, including the skull, vertebral column (all six cervical and 14 thoracic vertebrae), ribs, pectoral girdle and forelimbs (Figs 1, 3). The skull is intact, with the exception of the anterior portion of the left maxillary bone, which appears damaged, and both sclerotic rings, which are collapsed. Even fine details of the skeleton are visible (e.g., small sesamoids in the elbow; Fig. 3). Although having an intact skeleton is not necessarily an indication of its young geological age (for example, a Mesozoic gecko in amber exhibits a near pristine skeleton, Daza et al., 2016), the skeleton typically exhibits multiple fractures in the majority of Miocene *Sphaerodactylus* from Hispaniola. Using two X-rays from three ethanol preserved specimens of *Ebenavia boettgeri* (CAS 66195 [male, based on the presence of cloacal bones], CAS 66196 [gravid female with 2 eggs], 16° 54' 37.08" S, 49° 54' 40.716" E, St. Marie, and USNM 495825 25° 01' 12.0" S, 46° 58' 48.0" E [gravid female with 1 egg]) we were able to determine that the missing portion of the precloacal region (SVL) is between 18 and 25% of the SVL. Using these values,

the estimated SVL of the copal specimen is 34.8–37.8 mm. The estimated size matches several species of *Ebenavia*, although it greatly exceeds the adult size range of *E. maintimainty* (21–24 mm; Nussbaum & Raxworthy, 1998).

The specimen in copal was compared to similarly sized, formalin-fixed, ethanol preserved specimens of *E. boettgeri* from St. Marie (Fig. 4) and *E. robusta* (ZSM296/2010; Hawlitschek et al., 2018). Shared traits with *Ebenavia* include a small premaxilla with a short ascending nasal process (long in *Sphaerodactylus*); fused nasals (unfused in *Sphaerodactylus*); frontal broad with flat dorsal surface (narrow and convex in *Sphaerodactylus*); quadrate more or less straight with a slightly convex conch (curved and extremely convex conch in *Sphaerodactylus*); high number of foramina in the maxillary facial process (fewer foramina in *Sphaerodactylus*); high, discrete splenial (fused to coronoid in *Sphaerodactylus*); stapedial foramen absent (present in *Sphaerodactylus*); dentary ending at the level of the coronoid eminence (extending beyond the coronoid in *Sphaerodactylus*); and



retroarticular process narrow (broad in *Sphaerodactylus*). Morphology of the manus is very similar between the two genera, both *Ebenavia* and *Sphaerodactylus* having the same (plesiomorphic) phalangeal formula (2-3-4-5-3) and similar relative length of toes, from largest to smallest III>IV>II>V>I.

The copal gecko is clearly differentiated from *E. maintimainty*. Head length in the *E. inunguis* group is 9.2–9.5 mm (~9.5 in the copal gecko and between 5.4–5.9 in *E. maintimainty*), dorsal scales are partially keeled in the *E. inunguis* group and the copal gecko (vs. fully keeled in *E. maintimainty*), and the rostral scale is broad in the *E. inunguis* group and the copal gecko (vs. narrow in *E. maintimainty*). Using the key from Hawlitschek et al. (2018), we were able to confirm that ZFMK 94000 has the rostral scale separated from the nostril (Fig. 5), which is a character that defines *Ebenavia boettgeri*. This identification is compatible with the fact that Madagascar copal mines are only found in the distribution range of *E. safari* and *E. boettgeri*.

Even if the gecko in copal is potentially less than several hundred years old, the material provides an historical record of a living species of *Ebenavia* in the northeast of Madagascar based on the known location of copal deposits (Geirnaert, 2002). The North of Madagascar is a critical area for understanding the current distribution of *Ebenavia* in Madagascar, as ancestral area reconstructions may indicate a colonization of northern Madagascar from the Comoros Islands (Hawlitschek et al., 2017).

**Acknowledgements.** We thank Morris Flecks for taking the photograph of the specimen in copal. We also thank Emma Sherratt (The University of Adelaide), Jessie Maisano and Matt Colbert (UT|CT, The University of Texas at Austin) for assistance with the HRCT. We thank Rebecca Laver and two anonymous reviewers for comments on the manuscript. The Department of Biological Sciences sponsored Jordan P. Hunziker to present partial results of this project at The Joint Meeting of Ichthyologists and Herpetologists (JMIH) in Austin in 2017. Juan D. Daza received funding from NSF DEB 1657648 and the Biological Sciences Program at Sam Houston State University. Aaron M. Bauer received funding from NSF DEB 1555968 and 1556585, and the Gerald M. Lemole Endowed Chair Funds. PW is grateful to Jürgen Velten for an ongoing cooperation in amber research.

## REFERENCES

- Arnold EN, Azar D, Ineich I, Nel A (2002) The oldest reptile in amber: a 120-million-year-old lizard from Lebanon. *Journal of Zoology* 258: 7–10
- Arnold EN, Poinar G. (2008) A 100 million year old gecko with sophisticated adhesive toe pads, preserved in amber from Myanmar. *Zootaxa* 1847: 62–68
- Bauer AM, Böhme W, Weitschat W (2005) An early Eocene gecko from Baltic amber and its implications for the evolution of gecko adhesion. *Journal of Zoology* 265: 327–332
- Böhme W (1984) Erstfund eines fossilen Kugelfingergeckos (Sauria: Gekkonidae: Sphaerodactylinae) aus Dominikani-schem Bernstein (Oligozän von Hispaniola, Antillen). *Salamandra* 20: 212–220
- Böhme W, Weitschat W (1998) Redescription of the Eocene lacertid lizard *Nucras succinea* Boulenger, 1917 from Baltic amber and its allocation to *Succinilacerta* n. gen. *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg* 81: 203–222
- Borsuk-Białynicka M, Lubka M, Böhme W (1999) A lizard from Baltic amber (Eocene) and the ancestry of the crown group lacertids. *Acta Paleontologica Polonica* 44: 349–382
- Bosselaers J, Dierick M, Cnudde V, Masschaele B, Van Hoorebeke L, Jacobs P (2010) High-resolution X-ray computed tomography of an extant new *Donuea* (Araneae: Liocranidae) species in Madagascan copal. *Zootaxa* 2427: 25–35
- Castañeda MdR, Sherratt E, Losos JB (2014) The Mexican amber anole, *Anolis electrum*, within a phylogenetic context: implications for the origins of Caribbean anoles. *Zoological Journal of the Linnean Society* 172: 133–144
- Černáňský A, Augé ML (2013) New species of the genus *Plesi-lacerta* (Squamata: Lacertidae) from the Upper Oligocene (MP28) of Southern Germany and a revision of the type species *Plesi-lacerta hydekkeri*. *Palaeontologia* 56: 79–94
- Daza JD, Abdala V, Thomas R, Bauer AM (2008) Skull anatomy of the miniaturized gecko *Sphaerodactylus roosevelti* (Squamata: Gekkota). *Journal of Morphology* 269: 1340–1364
- Daza JD, Bauer AM (2012) A new amber-embedded sphaerodactyl gecko from Hispaniola, with comments on the morphological synapomorphies of the Sphaerodactylidae. *Breviora* 529: 1–28
- Daza JD, Bauer AM, Stanley EL, Bolet A, Dickson B, Losos JB (In press) An enigmatic miniaturized and attenuate whole lizard from the Mid-Cretaceous amber of Myanmar. *Breviora*
- Daza JD, Bauer AM, Wagner P, Böhme W (2013) A reconsideration of *Sphaerodactylus dommeli* Böhme, 1984 (Squamata: Gekkota: Sphaerodactylidae), a Miocene lizard in amber. *Journal of Zoological Systematics and Evolutionary Research* 51: 55–63
- Daza JD, Stanley EL, Wagner P, Bauer AM, Grimaldi DA (2016) Mid-Cretaceous amber fossils illuminate the past diversity of tropical lizards. *Science Advances* 2: e1501080
- de Queiroz K, Chu L-R, Losos JB (1998) A second *Anolis* lizard in Dominican amber and the systematics and ecological morphology of Dominican amber anoles. *American Museum Novitates* 3249: 1–23
- Fontanarrosa G, Daza JD, Abdala V (2018) Cretaceous fossil gecko hand reveals a strikingly modern scansorial morphology: Qualitative and biometric analysis of an amber-preserved lizard hand. *Cretaceous Research* 84: 120–133
- Geirnaert E (2002) L'ambre – miel de fortune et mémoire de vie. Les Editions du Piat, Saint-Julien-du-Pinet
- Glaw F, Vences M (2006) Field guide to the amphibians and reptiles of Madagascar. 3<sup>rd</sup> edition. Vences & Glaw Verlag GbR, Munich
- Hawlitschek O, Toussaint EFA, Gehring PS, Ratsoavina FM, Cole N, Crottini A, Nopper J, Lam AW, Vences M, Glaw F (2017) Gecko phylogeography in the Western Indian Ocean region: the oldest clade of *Ebenavia inunguis* lives on the youngest island. *Journal of Biogeography* 44: 409–420
- Hawlitschek O, Scherz MD, Ruthensteiner B, Crottini A, Glaw F. 2018. Computational molecular species delimitation and taxonomic revision of the gecko genus *Ebenavia* Boettger, 1878. *The Science of Nature* 105: 1–21
- Lazell Jr J (1965) An *Anolis* (Sauria, Iguanidae) in amber. *Journal of Paleontology* 39: 379–382

- Lourenço WR (1996) Premier cas connu d'un sub-fossile de scorpion dans le copal de Madagascar. *Comptes Rendus de l'Académie des Sciences, Paris Series Ila* 323: 889–891
- Nussbaum RA, Raxworthy CJ (1998) Revision of the genus *Ebenavia* Boettger (Reptilia: Squamata: Gekkonidae). *Herpetologica* 54: 18–34
- Penney D., Ono H, Selden PA (2005) A new synonymy for the Madagascan copal spider fauna (Araneae, Selenopidae). *Journal of Afrotropical Zoology* 2: 41–44
- Perrichot V, Néraudeau D (2005) Reptile skin remains in the Cretaceous amber of France. *Comptes Rendus Palevol* 4: 47–51
- Poinar GOJ (1999) Cenozoic fauna and flora in amber. *Estudios del Museo Ciencias Naturales de Álava* 14: 151–154
- Polcyn M, Rogers IJ, Kobayashi Y, Jacobs L (2002) Computed tomography of an *Anolis* lizard in Dominican amber: systematic, taphonomic, biogeographic, and evolutionary implications. *Palaeontologia Electronica* 5: 1–13
- Ragazzi, E, Roghi G., Giaretta A., Gianolla P (2003) Classification of amber based on thermal analysis. *Thermochimica Acta* 404: 43–54
- Ramanamanjato J-B, McIntyre PB, Nussbaum RA (2002) Reptile, amphibian, and lemur diversity of the Malahelo Forest, a biogeographical transition zone in southeastern Madagascar. *Biodiversity and Conservation* 11: 1791–1807
- Rieppel O (1980) Green anole in Dominican amber. *Nature* 286: 486–487
- Schlüter T, Gnielinski Fv (1987) The East African copal. Its geological, stratigraphic, palaeontologic significance and comparison with fossil resins of similar age. *National Museum of Tanzania Occasional Paper, Dar es Salaam, Tanzania* 8: 1–32
- Sherratt E, Castañeda MdR, Garwood RJ, Mahler DL, Sanger TJ, Herrel A, Losos JB (2015) Amber fossils demonstrate deep-time stability of Caribbean lizard communities. *Proceedings of the National Academy of Sciences of the United States* 112: 9961–9966
- Uetz P, Freed P, Database JHTR (2018) The reptile database. Online at <http://www.reptile-database.org/> (last access: July 2, 2018)
- Winkler W, Kirchner ECh, Asenbaum A, Musso M (2001) A Raman spectroscopic approach to the maturation process of fossil resins. *Journal of Raman Spectroscopy* 32: 59–63
- Xing LD, Caldwell MW, Chen R, Nydam RL, Palci A, Simões TR, McKellar RC, Lee MSY, Liu Y, Shi HL, Wang K, Bai M (2018) A Mid-Cretaceous embryonic-to-neonate snake in amber from myanmar. *Science Advances* 4: eaat5042

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Bonn zoological Bulletin - früher Bonner Zoologische Beiträge.](#)

Jahr/Year: 2018

Band/Volume: [67](#)

Autor(en)/Author(s): Daza Juan D., Hunziker Jordan P., Bauer Aaron M., Wagner Philipp, Böhme Wolfgang

Artikel/Article: [Things are not always as they seem: High-resolution X-ray CT scanning reveals the first resin-embedded miniature gecko of the genus Ebenavia 71-77](#)