

Bonn zoological Bulletin 70 (2): 351–357 2021 · Gębicki C. et al. https://doi.org/10.20363/BZB-2021.70.2.351

Research article

urn:lsid:zoobank.org:pub:C2125016-AEF6-4F7E-83D9-5718C378B405

Youngeewa simillima sp. nov. (Hemiptera: Cicadellidae: Mileewinae) from Eocene Baltic Amber

Cezary Gębicki¹, Marcin Walczak^{2,*} & Marzena Zmarzły³

¹ Faculty of Science & Technology, Jan Długosz University in Częstochowa, Al. Armii Krajowej 13/15, PL-42-200 Czestochowa, Poland

^{2,3}Zoology Research Team, Institute of Biology, Biotechnology and Environmental Protection, Faculty of Natural Sciences,

University of Silesia in Katowice, Bankowa 9, PL-40-007 Katowice, Poland

*Corresponding author: Email: marcin.walczak@us.edu.pl

¹urn:lsid:zoobank.org:author:6FC4935A-D82E-45BE-BD77-8E9EFEB9BFDD ²urn:lsid:zoobank.org:author:046F0D52-7FBA-4D7D-9F79-AEBF7F2C98B8 ³urn:lsid:zoobank.org:author:E11DD0B8-412A-41EE-A8CC-4114A2CD1F54

Abstract. The paper describes *Youngeewa simillima* sp. nov. from Eocene Baltic amber as a new species within the tribe Mileewini Evans, 1947 (Mileewinae Evans, 1947). Due to its poor condition, the specimen is only tentatively placed in the genus *Youngeewa* Gębicki & Szwedo, 2001, differing from the other known species of the genus, *Youngeewa bicolorata* Gębicki & Szwedo, 2001, mainly by the elements of wing venation.

Key words. Youngeewa, Cicadellidae, Mileewinae, new species, Baltic amber.

INTRODUCTION

A leafhopper specimen in Baltic amber is described as a new species in the present work. The described species is the third known fossil representative of the Mileewinae Evans, 1947 subfamily so far. Previously, Youngeewa bicolorata Gębicki & Szwedo, 2001 and Eomileewa eridani Gebicki & Szwedo, 2001 (Mileewinae: Mileewini) have been described from Eocene Baltic amber (Gebicki & Szwedo 2001). Specimens of these two species were better preserved than the new species described here and had well-preserved features for the subfamily. The specimen described here has details of its head obscured and is placed in the subfamily based mainly on its short clavus, an important diagnostic feature of the tribe Mileewini Evans, 1947, and it is tentatively placed in the genus Youngeewa Gębicki & Szwedo, 2001. Three other features characteristic of the extant Mileewini of the Old world have not been confirmed, i.e., vein RP (=Sc+R or R_{2+3}) extending to the anterior rather than outer margin of the wing and the marginal vein reaching the wing margin before A₁ and fold between PCu and A₁. Two other differences concerning the width and shape of the forewing apex are variable in Mileewini (see Dietrich et al. 2017; He et al. 2021). Mileewinae Evans, 1947 are an evolutionary old auchenorrhynchan group originating from the Early Cretaceous central Gondwanaland (Nielson & Knight 2000). The first representatives of the subfamily were estimated to have originated around 126 MA, based on molecular analysis and incorporating *Youngeewa* Gębicki & Szwedo, 2001 as a calibration point (Dietrich et al. 2017). Mileewini seem to be closely related and probably derived from the subfamily Cicadellinae Latreille, 1825 and are also similar in wing venation to the subfamily Typhlocybinae Kirschbaum, 1868 (Balme 2007; Dietrich et al. 2010), especially to the extinct Protodikraneurini Gębicki & Szwedo, 2006 (Gębicki & Szwedo 2006) and Alebrini McAtee, 1926 but representatives of these taxa have an acute hind basitarsomere unique to all Typhlocybinae.

MATERIAL AND METHODS

A male was examined. Only external structures were examined using a stereoscopic microscope Olympus SZX9 Nikon Eclipse. Photographs were taken with the Canon Eos camera with extension rings. All the equipment is available at the University of Silesia, Faculty of Natural Sciences, Institute of Biology, Biotechnology and Environmental Protection. All photographs and drawings were digitalised and worked on with CorelDRAW software.

Research funding was provided by the Dean of the Faculty of Natural Sciences, Institute of Biology, Biotechnology and Environmental Protection, University of Silesia in Katowice.

The holotype *Y. simillima* Gębicki & Walczak sp. nov. bears labels; quoting the labels of specimen is done in the following way: (/) is used to divide data in different rows on the label, (;) is used to divide data on different labels, ([]) is used for the author's comments.

Abbreviations for morphological terms

|--|

- CuA = Cubitus anterior vein (CuA1 / CuA2)
- CuP = Cubitus posterior vein
- M = Media vein
- MA = Media anterior vein
- MP = Media poserior vein
- PCu = Postcubitus vein
- Sc = Subcosta vein
- R = Radius vein (R / R2)

RA = Radius anterior vein (RA1) RP = Radius posterior vein

RESULTS

Youngeewa simillima Gębicki & Walczak sp. nov. (Figs 1–4) *urn:lsid:zoobank.org:act:0C8A599B-570B-4DE4-99E2-183BE988E0F3*

Type material. Holotype ♂ [written in all caps], *Young-eewa simillima* / Gębicki & Walczak / sp. nov. [red label, written in bold]; Eocen Baltic amber / No. 5008 / coll. Carsten Gröhn.



Fig. 1. Youngeewa simillima Gębicki & Walczak sp. nov. A. Head and pronotum from the dorsal side. B. Face and right leg of the first pair, right leg of the second pair, and foot of third pair of legs. C. Face and fore legs. D. Specimen from the ventral side.

Diagnosis. In general outline the species is close to *Y. bicolorata* but differs in the following characteristics: straight lateral margins of genae (no deep incision below the eye), apex of rostrum reaching beyond bases of middle coxae, veins Sc+RA₁ and transverse r-m and m-cu on fore wings are arranged in one line (as in Fig. 3C), and the distance between distal terminations of veins CuA₂ and CuP on hind wings is only 2.1 × as long as that between MP+C-uA and CuA₂ (measured on hind wing marginal vein), (in *Y. bicolorata* 2.3 ×). Hind wing cell delimited by veins R and M is somewhat broader starting from transverse vein r-m to the wing apex than the cell delimited by veins MA and MP+CuA₁. Fore wings distinctly lighter than those in *Y. bicolorata* and lacking any clear colour pattern.

Description. Total body length 3.70 mm (sex: male), detailed data showing the dimensions of selected parts of the body are included in Table 1.

Head. Coronal part of head strongly convex especially at apex, broad, "delta"-like in outline. Ocelli, along with entire surface of head, completely invisible, covered by a felt-like sheath (Fig. 1A). Coronal and frontal sutures invisible. Antennae slightly longer than head, located in distinct antennal pits in front of the eyes at inner eye margin (Figs 1B–C, 3A). Scapus short and cylindrical, pedicellus relatively narrow, strongly elongated, many times longer than first segment. Antennal flagellum about $3 \times$ narrower than pedicellus. Lateral margins of facial part of head almost straight, lacking a clear-cut incision under eyes. Frontoclypeus convex, distinctly broader in apical part, strongly tapered distally. In subapical part of frontoclypeus there is an arcuate concavity located in place of epistomal suture (Figs 1B, 3A). Anteclypeus almost square in shape, small, about $5 \times$ narrower than postclypeus at its widest point and separated from it by a deep transclypeal suture. Indistinct, 3-segmented rostrum reaching beyond bases of middle coxae (Fig. 1D).

Pronotum. Shorter than vertex with indistinct lateral carina (Figs 1A, 2A, 3B).

Wings. Fore wings relatively narrow and elongate, evidently reaching beyond the abdomen, at apex nearly as wide as corium and slightly rounded. Transverse veins of three apical cells (2–4) arranged in a single somewhat slanting line. Inner apical field longest, and largest, delimited by CuA_1 , which is slightly "S"-like; apical fields 1 and 3 narrow, rectangular in outline; outer field



Fig. 2. Youngeewa simillima Gębicki & Walczak sp. nov. A. Specimen from the dorsal side. B. Hind wing (ventral view). C. Left mid leg. D. Right mid and hind leg. E. Hind leg tarsus in large magnification.



Fig. 3. Youngeewa simillima Gębicki & Walczak sp. nov. (reconstruction). A. Head from the ventral side. B. Head and pronotum from the dorsal side. C. Fore wing. D. Hind wing.



Fig. 4. *Youngeewa simillima* Gębicki & Walczak sp. nov. (reconstruction). **A**. Right fore leg from the ventral side. **B**. Right mid leg from the ventral side. **C**. Left hind leg from the dorsal side. **D**. Right hind leg from the ventral side. **E**. Hind leg tarsus in large magnification.

Male	Size [mm]
head length in midline	0.41 mm
head width	probably 0.7 mm*
frons length in midline	0.69 mm
frons width	probably 0.73 mm*
antennae length	0.99 mm
pronotum length in midline	0.38 mm
abdomen length in midline	1.74 mm
profemur	0.58 mm
protibia	0.63 mm
protarsus	0.31 mm
mesofemur	0.53 mm
mesotibia	0.78 mm
mesotarsus	0.31 mm
metafemur	1.04 mm
metatibia	1.49 mm
metatarsus	0.82 mm
fore wing lenght	2.95 mm
fore wing width	0.88 mm
hind wing length	2.62 mm
hind wing width	1.08 mm

 Table 1. Biometric data of Youngeewa simillima Gębicki &

 Walczak sp. nov. [*data difficult to measure]

is smallest, triangular in shape. Distances between main stems of longitudinal veins R, M and Cu nearly equal. Clavus reaching halfway along the wing, with two distinct, almost parallel veins PCu and A_1 running close to each other. Corium and costal cell lacking any transverse veins (Figs 1D, 3C). Fore wings slightly darker than hind wings, with darker colouration along the fields adjacent to main longitudinal veins (Fig. 2A).

Hind wings almost as long as fore wings. Veins R and M parallel in distal part and connected by a transverse vein r-m, which is located somewhat before nodus. Costal margin (lacking a distinct, deep incision) similar to that in species representing the genus *Mileewa* and to *Y. bicolorata*. Marginal vein reaching nodus; in proximity to the costal margin it runs parallel and joins internal wing margin outside vein A_1 . Membranous wing margin is narrowest between termination of veins MA and MP+CuA₁ whereas it is broadest at internal margin opposite termination of vein PCu (Figs 2B, 3D).

Legs. Fore legs. Profemur slightly longer than protibia. Protibia with quite large setae along inner margin. Segments of protarsus small, third segments longest (Figs 1B–C, 4A). Middle legs. With sparse fine setae along inner margin of mesotibia (Figs 2C–D, 4B). Hind legs. Spines on hind metatibia long, distinctly longer than metatibia thickness. Apex of metafemur with macrochaetae in pattern 2+1+1 (Figs 1D, 2D, 4C–D). Basal segment of metatarsus longest and at apex expanded into a process with two spines, outer margin with five sharp spines and two elongate marcochaetae, which are situated near the apex on the inner side. The middle segment with three blunt apical spines and a single sharp lateral one. Hind pretarsus shortest (Figs 2E, 4E).

Male. Copulatory block relatively short, details not clearly visible in examined specimen.

Etymology. Species name originates from the Latin word "simillima" due to the fact that the species presented here is very similar to the *Youngeewa bicolorata* described earlier.

Remarks. Substantial areas of head including rostrum, parts of prothorax and some parts of legs covered with white, felt-like substance which blurs the details of the structure. Accordingly, it is impossible to determine the location of the ocelli, which constitutes an important taxonomic feature. Right fore- and hind wings are well-visible and their venation is clear. In the male the apex of abdomen is partly covered by the left hind wing, which makes analyses difficult. There are no animal syninclusions. For reasoning of the taxonomic placement of the new species, see introduction and species diagnosis.

DISCUSSION

The species of the genus *Youngeewa* are close to many extant Mileewini (Mileewinae Evans, 1947) species in having a large dome-shaped head, a convex clypeus (also in Cicadellinae Latreille, 1825), reduction of forewing venation, a distinct fore wing appendix and shortened fore wing clavus. These features are also found in the South American genus Orsalebra Young, 1952 (belonging to the Typhlocybine tribe Alebrini, see Introduction). However, in this genus the marginal vein of the forewing does not reach the apex of vein RP, which is typical of most Mileewi, but reaches only the apex of vein M; in most Alebrini the marginal vein is fused with the apical wing margin. Reduced fore wing venation in Mileewinae is typical of Typlocybinae as well as some Cicadellinae and Nirvanini Baker, 1923 (Evacanthinae Metcalf, 1939 sensu Dietrich 2004). Hind wing venation is similar in the above mentioned tribes while veins R and M running parallel along their entire length (not only in their apical part) are found in Protodikraneurini Gebicki & Szwedo, 2006 (extinct Typhlocybinae) and in the extant neotropical mileewine genus Amahuaka Melichar, 1926, as well as in South American representatives of the tribe Tungurahualini Dietrich, 2011 of the subfamily Mileewinae (Dietrich 2011; Krishnankutty & Dietrich 2011).

Acknowledgments. We would like to thank Jacek Szwedo from the University of Gdańsk for the loan of the material and

Krzysztof Kudła from the University of Silesia in Katowice for graphic corrections.

REFERENCES

- Balme GR (2007) Phylogeny and systematics of the leafhopper subfamily Typhlocybinae (Insecta: Hemiptera: Cicadellidae). PhD dissertation. Entomology, Raleigh, North Carolina
- Dietrich CH (2004) Phylogeny of the leafhopper subfamily Evacanthinae with a review of Neotropical species and notes on related groups (Hemiptera: Membracoidea: Cicadellidae). Systematic Entomology 29: 455–487. https://doi.org/10.1111/j.0307-6970.2004.00250.x
- Dietrich CH (2011) Tungurahualini, a new tribe of Neotropical leafhoppers with notes on the subfamily Mileewinae (Hemiptera, Cicadellidae). Zookeys 124: 19–39. https://doi.org/10.3897/zookeys.124.1561
- Dietrich CH, Allen JM, Lemmon AR, Lemmon EM, Takiya DM, Evangelista O, Walden KKO, Grady PGS, Johnson KP (2017) Anchored Hybrid Enrichment-Based Phylogenomics of Leafhoppers and Treehoppers (Hemiptera: Cicadomorpha: Membracoidea). Insect Systematics and Diversity 1 (1): 57– 72. https://doi.org/10.1093/isd/ixx003

- Dietrich CH, Dmitriev DA, Rakitov RA, Takiya DM, Webb MD, Zahniser JN (2010) Phylogeny of Cicadellidae (Hemiptera: Cicadomorpha: Membracoidea) based on morphological characters. Pp. 48–49 in: Proceedings of the 13th International Auchenorrhyncha Congress, 28.06–02.07.2010, Vaison-la-Romaine, France
- Gębicki C, Szwedo J (2001) The first record of fossil Mileewinae from Eocene Baltic amber (Hemiptera: Membracoidea: Cicadellidae). Annales zoologici 51 (4): 417–422
- Gębicki C, Szwedo J (2006) Protodikraneurini trib. nov. from the Eocene Baltic amber (Hemiptera: Cicadellidae: Typhlocybinae). Annales zoologici 56 (4): 763–783
- He H-L, Yan B, Yang M, Webb MD (2021) Four new species of Mileewini leafhoppers (Hemiptera: Cicadellidae: Mileewinae) from China, with a checklist to Chinese species. Zootaxa 4949(3):521–540. https://doi.org/10.11646/zootaxa.4949.3.5
- Krishnankutty SM, Dietrich CH (2011) Review of Mileewinae Leafhoppers (Hemiptera: Cicadellidae: Mileewinae) in Madagascar, with description of seven new species. Annals of the Entomological Society of America 104 (4): 636–648. https://doi.org/10.1603/AN11022
- Nielson MW, Knight WJ (2000) Distributional patterns and possible origin of leafhoppers (Homoptera, Cicadellidae). Revista Brasileira de Zoologia 17 (1): 81–156. https://doi.org/10.1590/S0101-81752000000100010

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Bonn zoological Bulletin - früher Bonner Zoologische Beiträge.

Jahr/Year: 2021

Band/Volume: 70

Autor(en)/Author(s): Gebicki Cezary, Walczak Marcin, ZmarzÅ, y Marzena

Artikel/Article: Youngeewa simillima sp. nov. (Hemiptera: Cicadellidae: Mileewinae) from Eocene Baltic Amber 351-357