

Scientific note

[urn:lsid:zoobank.org:pub:3B5488F7-326A-4276-B266-3BCA937DF656](https://zoobank.org/pub:3B5488F7-326A-4276-B266-3BCA937DF656)**First record of the invasive grapevine aphid *Aphis illinoisensis* Shimer, 1866 (Hemiptera: Aphididae) in mainland Italy****Alice Casiraghi***Instituto Valenciano de Investigaciones Agrarias. (IVIA). Unidad de Entomología, Centro de Protección Vegetal y Biotecnología. Ctra. Moncada-Náquera Km. 4,5. E-46113 Moncada, Valencia, Spain**Email: casiraghi_alia@gva.es*[urn:lsid:zoobank.org:author:905837A0-9C77-4D57-A895-B19DD9FF790B](https://zoobank.org/author:905837A0-9C77-4D57-A895-B19DD9FF790B)

Abstract. Apterous and winged viviparous females of the Nearctic *Aphis illinoisensis* Shimer, 1866 have been recorded for the first time in mainland Italy (Rome) on branches of *Vitis labrusca* L. in 2021, thanks to photos published in a Facebook garden group.

Key words. Invasive species, citizen science, grapevine pest, new record, Vitaceae.

Garden and insect groups on Facebook are typically dedicated to questions about plant health and pest problems, but sometimes they can result in being a source of information about exotic or invasive species detection and expansion (Šeat 2015, Martinović et al. 2019, Vujić et al. 2021). In our case, a photograph of aphids on fox grape *Vitis labrusca* L. taken on 17th September 2021 in a private garden in the urban area of Colli Aniene (Rome, Italy) showed the first record of the Nearctic species *Aphis illinoisensis* Shimer, 1866 (Hemiptera: Aphididae) in mainland Italy.

The colony, composed by both apterous and winged females and their nymphs, was not attended by ants (Fig. 1). Apterous females are characterized by a shiny dark reddish brown colour, wholly black hind tibiae and dark, long and divergent siphunculi. Winged females are similar, and nymphs typically present a lighter reddish-brown colouration with black siphunculi (Blackman & Eastop, 2021). The identification was confirmed by aphidologist Nicolás Pérez Hidalgo (Departamento de Artrópodos, Museo de Ciencias Naturales de Barcelona, Spain).

Unfortunately, specimens could not be collected nor more and detailed photos be taken, since the colony has been immediately fumigated after the identification and up to date no other winged female colonized the areas. Contact with the author of the photograph has been kept open in order to await new developments.

The invasive American grapevine aphid is native to North America (Blackman & Eastop, 2021), but it is also widespread in Central and South America (Ramírez Guevara & Zuluaga 1995, Ortego et al. 2004, Trejo-Loyo et al. 2004, Etienne 2005, Zamora Mejías et al. 2010), where it presents a dioecious holocycle, host-alternating

between *Viburnum prunifolium* L. (Adoxaceae) and Vitaceae plants (Blackman & Eastop 2021). In 2002 *A. illinoisensis* was first recorded in the Palearctic, more specifically in Turkey (Remaudière et al. 2003). Since then, the species has been reported throughout the Mediterranean Basin and Middle East, i.e., Crete, Greece (Tsitsipis et al. 2005), Cyprus (Kocadal & Ulusoy 2006), Spain (Pérez Hidalgo et al. 2011), Israel (Barjadze & Ben-Dov 2011), Tunisia (Kamel-Ben Halima & Mdellel 2010), Algeria (Laamari & Coeur d'Acier 2010), Montenegro (Petrović-Obradović et al. 2010), Libya (Havelka et al. 2011), Malta (Mifsud & Pérez Hidalgo 2011), Egypt (El-Gantiry et al. 2012), Saudi Arabia (Hussain et al. 2015) and Iran (Heidarnia & Derakhshan 2018). In Italy, this species was documented from the island of Sicily, in the province of Catania (Cocuzza & Barbagallo 2011). More recently, it has been recorded even in Central Europe: Slovenia (Seljak 2021) and France (Mouttet & Bálmes 2021). On the Eurasian continent *A. illinoisensis* seems to develop anholocycles on *Vitis* plants and no sexual morphs have been recorded (Blackman & Eastop 2021). Its expansion appears to be a product of the movement of winged alatae (Havelka et al. 2011).

The grapevine aphid is considered invasive because of the potential damage caused to the wine and grape production. It has been recorded as a vine pest that can decrease the developing of young shoots, leaves and fruits and produce leaf necrotic spots (McGrew & Still 1979, Cocuzza & Barbagallo, 2011), but its impact in Europe does not appear to be as important as the one produced by the grape phylloxera *Daktulosphaira vitifoliae* (Fitch, 1855) (Hemiptera: Phylloxeridae) (Kamel-Ben & Mdellel 2010, Havelka et al. 2011). For the moment, because



Fig. 1. Colony of *Aphis illinoisensis* from Rome (Italy) with winged and apterous female morphs on *Vitis labrusca*. The photo was taken by the Italian citizen Sara Bonanno in her garden in the neighbourhood of Colli Aniene and posted in a Facebook garden group about plant pests.

of its occasional records, *A. illinoisensis* can be considered a minor pest of vines in Italy.

Once again, citizen science allows specialists to gain more information about the real distribution of exotic and invasive species.

Acknowledgments. The author is sincerely thankful to Dr. Nicolás Pérez Hidalgo who confirmed aphid identification and to Sara Bonanno who decided to share their photos and geographical information. She also wants to thank the reviewers for their work and effort to improve this little scientific note.

REFERENCES

- Barjadze S, Ben-Dov Y (2011) The grapevine aphid *Aphis illinoisensis*: an invasive pest in Israel. *Phytoparasitica* 39 (1): 55–57
- Blackman RL, Eastop VF (2021) Aphids on World's plants. An online identification and information guide. Online at <http://www.aphidsonworldsplants.info> [last accessed 30 Sep. 2021]
- Cocuzza GE, Barbagallo S (2011) The appearance of the American aphid, *Aphis illinoisensis*, in grapes in Sicily. *Informatore Agrario* 67 (34): 81–83 [in Italian]
- El-Gantiry AM, El-Heneidy AH, Mousa SF, Adly D (2012) *Aphis illinoisensis* Shimer (Hemiptera: Aphididae) a recent invasive aphid species in Egypt. *Egyptian Journal of Biological Pest Control* 22 (2): 225
- Etienne J (2005) Les pucerons de Guadeloupe, des Grandes et Petites Antilles (Hemiptera, Aphididae). *Bulletin de la Société Entomologique de France* 110 (4–5): 455–462 [in French]
- Havelka J, Schukshuk AH, Ghaliow M, Laamari M, Kavalieratos NG, Tomanović Ž, Rakhshani E, Pons X, Starý P (2011) Review of invasive grapevine aphid, *Aphis illinoisensis* Shimer and native parasitoids in the Mediterranean (Hemiptera, Aphididae; Hymenoptera, Braconidae, Aphidiniinae). *Archives of Biological Science* 63 (1): 269–74
- Heidarnia M, Derakhshan A (2018) Faunistic Study of Esfarayen aphids and the new report of the species, *Aphis illinoisensis* Shimer, 1866 for Iran fauna. *Proceedings of the 23th Iranian Plant Protection Congress*. Gorgan University of Agricultural Sciences and Natural Resources
- Hussain S, Aldryhim Y, Al-Dhafer H, Halbert SE, Thomas J (2015) New aphid records for Saudi Arabia (Hemiptera: Aphidoidea). *Zoology in the Middle East* 61 (4): 368–371
- Kamel-Ben Halima M, Mdellel L (2010) First record of the grapevine aphid, *Aphis illinoisensis* Shimer, in Tunisia. *EPPO Bulletin* 40: 191–192
- Kocadal E, Ulusoy MR (2006) A survey of Aphidoidea species (Homoptera) in the Turkish Republic of Northern Cyprus (TRNC). MSc. Thesis, C. Ü. Fen bilimleri Enstitüsü 1–26
- Laamari M, Coeur d'Acier A (2010) Le puceron de la vigne *Aphis illinoisensis* arrive en algérie. *EPPO Bulletin* 40: 167–168 [in French]
- Martinović M, Gjeldum A, Koren T (2019) *Belonochilus numenius* (Say, 1832) (Heteroptera: Lygaeidae), a new invasive species in Croatia. *Natura Croatica: Periodicum Musei Historiae Naturalis Croatici* 28 (2): 481–484
- McGrew JR, Steel GW (1979) Control of grape diseases and insects in the Eastern United States. *Farmers' Bulletin* No. 1893, United States Department of Agriculture
- Mifsud D, Pérez Hidalgo N (2011) The grapevine aphid *Aphis illinoisensis*: a good example of recent invasion and rapid colonization by aphids. *Bulletin OEPP/EPPO* Bulletin 41 (2): 183–184
- Mouttet R, Balmès V (2021) Un nouveau puceron sur vigne en France: *Aphis illinoisensis* Shimer, 1866 (Hemiptera, Aphididae). *Bulletin de la Société entomologique de France* 126 (2): 206–208 [in French]
- Ortego J, Difabio ME, Mier Durante MP (2004) New records and actualized check-list of aphids (Hemiptera: Aphididae) from Argentina. *Revista de la Sociedad Entomologica Argentina* 63 (1–2): 19–30
- Pérez Hidalgo N, Laguna García F, Nieto Nafria JM (2011) First Spanish record of *Aphis illinoisensis* Schimer (Hemiptera: Aphididae), the grapevine aphid. *Boletín de la Sociedad Entomológica Aragonesa (S.E.A)* 49: 321–323 [in Spanish]
- Petrović-Obradović O, Tomanović Ž, Poljaković-Pajnik L, Hrncić S, Vučetić A, Radonjić S (2010) New invasive species of aphids (Hemiptera, Aphididae) in Serbia and Montenegro. *Archives of Biological Sciences* 62 (3): 777–782

- Ramírez Guevara G, Zuluaga CJI (1995) Identification of hymenopterous parasitoids of aphids on cotton, Citrus and horticultural crops in Valle del Cauca. *Revista Colombiana de Entomologia* 21 (2): 71–76
- Remaudière G, Sertkaya E, Özdemir I (2003) Alerte! découverte en Turquie du puceron américain *Aphis illinoisensis* nuisible à la vigne. *Revue Française d'Entomologie* 25: 170 [in French]
- Seljak G (2021) Prva najdba črne trtne uši (*Aphis illinoisensis* Shimer, 1866) v Sloveniji (Hemiptera, Aphidoidea: Aphididae). *Acta Entomologica Slovenica* 29: 1 [in Slovenian]
- Šeat J (2015) *Halyomorpha halys* (Stål, 1855) (Heteroptera: Pentatomidae) a new invasive species in Serbia. *Acta entomologica serbica* 20: 167–171
- Trejo-Loyo A, Pena-Martinez R, Villegas-Jimenez N (2004) Afidofauna (Hemiptera: Aphididae) de Cuernavaca, Morelos, Mexico. *Folia Entomologica Mexicana* 43 (2): 191–202 [in Spanish]
- Tsitsipis JA, Angelakis E, Margaritopoulos JT, Tsamandali K, Zarpas KD (2005) First record of the grapevine aphid *Aphis illinoisensis* in the island of Kriti, Greece. *Bulletin OEPP/EPPO Bulletin* 35: 541–542
- Vujić MD, Ivković S, Rekecki T, Krstić D, Stanković V, Đurić M, Tot I (2021) A first record of the alien mantis species *Hierodula tenuidentata* (Mantodea: Mantidae) in Serbia. *Acta Entomologica Serbica* 26 (1): 1–7
- Zamora Mejías D, Hanson PE, Starý P (2010) Survey of the aphid parasitoids (Hymenoptera: Braconidae: Aphidiinae) of Costa Rica with information on their aphid (Hemiptera: Aphidoidea): plant associations. *Psyche: A Journal of Entomology*, Article ID: 278643. <https://doi.org/10.1155/2010/278643>

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Bonn zoological Bulletin - früher Bonner Zoologische Beiträge.](#)

Jahr/Year: 2021

Band/Volume: [70](#)

Autor(en)/Author(s): Casiraghi Alice

Artikel/Article: [First record of the invasive grapevine aphid *Aphis illinoisensis* Shimer, 1866 \(Hemiptera: Aphididae\) in mainland Italy 373-375](#)