

8. Die Sternopleuren nur oben ein wenig weißlich schillernd. Die Vorderschenkel unten an der Basis und gegen die Spitze mit je einem abstehenden Haar. 2 Vertikale, starke Orbitale und Postvertikale (siehe die Gattungs-Übersicht!); 3,25 mm l. *Meroplus stercorarius* Rob.-Desv.

Die Sternopleuren ganz weiß schillernd. Die Vorderschenkel unten ohne abstehende Haare. 1 Vertikale, schwache Orbitale und Postvertikale (Gattungs-Übersicht!); 3—3,5 mm l.

Enicita annulipes Meig.

Description of a New Species of Parasitic *Cynipidae* from Kuching, Borneo. (Hym.)

By P. Cameron, Whitle, New Mills (England).

Psilosema rufipes sp. nov.

Black, smooth and shining; the basal 7 joints of the antennae and the legs red, wings hyaline, the nervures fuscous, the edges with a dense, distinct hair fringe; the radial cellule more than twice longer than wide, the apical abscissa of the radius curved, distinctly thicker than the rest, the nervure bounding its anterior being also thickened, there is no closed cubital cellule, only the apical transverse cubital nervure being present. There is a 6-jointed, not very clearly separated club; the basal joint is not quite so stout as the others; the basal 4 joints are as broad as long, the 5th longer than wide, the apical clearly longer than it, twice longer than wide and narrowed towards the apex. Scutellar cup small, round; the depression at the foot of the scutellum large. Middle of metanotum raised, bordered, narrowed towards the base, the sides apparently covered with short white pubescence; the whole is tinged with rufous. — Length 1 mm.

The type of *Psilosema* Kieffer has a 5-jointed antennal club, but Ashmead says of it and its ally *Schizosema* Kief. "Club of antennae 5-jointed (rarely 6-jointed)". There is no other genus in which the present species can be placed, c. f. Ashmead, Psyche, 1903, 62. This is the first species of *Eucoilinae* I have seen from the Oriental Zoological Region.