

A new *Dendroides* species from China (Coleoptera, Pyrochroidae), with a key to the Palaearctic species

Xin-Mei Yang¹, Daniel K. Young², Bao-Lin Yang³, Zhao Pan¹

1 Key Laboratory of Zoological Systematics and Application of Hebei Province, School of Life Sciences, Institute of Life Science and Green Development, Hebei University, Baoding 071002, Hebei, China

2 Department of Entomology, University of Wisconsin, Madison, WI 53706, USA

3 Huanglong National Scenic Reserve, Songpan 623300, Sichuan, China

<https://zoobank.org/47D4FC20-681C-4363-BBA9-962A44681C98D>

Corresponding author: Zhao Pan (panzhao86@yeah.net)

Academic editor: Dmitry Telnov ♦ Received 16 January 2025 ♦ Accepted 6 March 2025 ♦ Published 17 April 2025

Abstract

Dendroides Blair, 1914 is distributed in North America and East Asia, with eight described species. Currently, a new species, *Dendroides huanglong* Pan, sp. nov., is described and illustrated from Sichuan Province, China, representing the third *Dendroides* species for the Palaearctic Region. A key to the Palaearctic species of *Dendroides* is provided and the phylogenetic position of *Dendroides* is briefly discussed.

Key Words

China, *Dendroides*, fire-coloured beetle, key, new species

Introduction

Dendroides Latreille, 1810 is a small pyrochroid genus within the subfamily Pyrochroinae, with eight species: six from North America and two from East Asia.

The generic scope of *Dendroides* was first clarified by Blair (1914), with a key to the six species, including a new species, *D. lesnei* Blair, 1914, described from Honshu, Japan. Fourteen years later, another Nearctic species was described by Van Dyke (1928). Since then, no new species have been added to the fauna of the Nearctic Region. Half a century later, in the revision of North American Pyrochroidae, Young (1975) provided diagnoses for adults, larvae and pupae of all reliably confirmed species and keys to them, respectively.

In 1936, Kôno described *Dendroides nakabusana* from Honshu in Japan, based on its compound eyes being slightly larger than those of *D. lesnei*. However, it was synonymised by Young (2005a), based on examination of types of both species. The other Palaearctic species is *D. ussuriensis* L.N. Medvedev, 1977.

This species was described from the Primorye territory of Russia, Heilongjiang of China [“ст. Гаолинцы” (= Gaolingzi Zhan, 高岭子站) and “ст. Шитоухецы” (= Shitouhezi Zhan, 石头河子站). Both are stations on the original Chinese Eastern Railway and are in Shangzhi, Harbin, Heilongjiang] and North Korea (Medvedev 1977).

During an investigation of the insect diversity in Huanglong National Nature Reserve (Sichuan, China), individuals of a new species, *Dendroides huanglong* Pan sp. nov., were collected using a light trap. The new species is described and illustrated herein and a key to *Dendroides* species from the Palaearctic Region is provided.

Materials and methods

The type material is deposited in the Museum of Hebei University, Baoding, China (MHB). The specimens were studied using a Nikon SMZ1500 and the images were taken using a Canon EOS 5D Mark III (Canon Inc.,

Tokyo, Japan) with a Laowa FF 100 mm F2.8 CA-Dreamer Macro 2 \times or Laowa FF 25 mm F2.8 Ultra Macro 2.5–5 \times (Anhui Changgeng Optics Technology Co., Ltd, Hefei, China). Figures of the antennae were hand drawn using a Nikon SMZ1500 with a camera lucida. Label data are presented verbatim. Line breaks on labels are denoted by a single slash (/).

Most of the terms in the description are from previous literature (e.g. Young (1975)). The ocular index (OI) = 100 \times minimum dorsal distance between compound eyes / maximal dorsal width across compound eyes (Campbell and Marshall 1964).

Results

Dendroides huanglong Pan, sp. nov.

<https://zoobank.org/465475DB-954B-47A4-94FB-23730FE68BC7>

Type locality. CHINA • Sichuan Province, Aba Tibetan and Qiang Autonomous Prefecture, Songpan County, Huanglong National Natural Reserve.

Type specimens. *Holotype*: ♂, labelled “2024. VII.17 / 四川黄龙自然保护区 [Sichuan, Huanglong Natural Reserve] / 丹云峡保护站 (灯诱) [Danyunxia Protection Station (light trap)] / 河北大学博物馆 [Museum of Hebei University]”, “32°44'59"N / 103°57'38"E / elev. 2542 m, 王成涛 [Cheng-Tao Wang] leg./ 河北大学博物馆 [Museum of Hebei University]”, “HOLOTYPE / *Dendroides huanglong* sp. n. / Det. Pan” (MHBU). *Paratypes*: 1♂, 1♀, with same collection event labels as the holotype and additional labels: “P4H1 or P4H2”, “PARATYPE / *Dendroides huanglong* sp. n. / Det. Pan” (MHBU).

Diagnosis. Like *D. ussuriensis*, this new species can be easily distinguished from other *Dendroides* species by its colouration: body reddish testaceous, but maxillary palpomeres, antennae and most parts of legs black (Fig. 1). The body size of *D. huanglong* sp. nov. is larger than *D. ussuriensis*; the body length is 14.5–17.2 mm in the former and 12.0–13.0 mm in the latter. The compound eyes of *D. huanglong* sp. nov. are larger than those of *D. ussuriensis*: they meet above the head in the male of the former (Fig. 2) but not in the latter and OI is 8.1 in the female of the former (Fig. 3), but more than 25.0 in the latter. Additionally, the rami of the antennal flagellomeres are more prominent and elongate in *D. huanglong* sp. nov. than those of *D. ussuriensis* (Figs 4, 5), especially in the female, wherein flagellomere I has a distinct ramus, while flagellomere I in *D. ussuriensis* lacks a ramus; rami of flagellomeres III–VII are each longer than the preceding flagellomere in *D. huanglong* sp. nov., but shorter than the preceding flagellomere in *D. ussuriensis*.

Description. Male: Body (Fig. 1) reddish testaceous, except apex of mandibles, maxillary palpomeres, antennae, eyes and legs (except base of coxae) black. Body covered with medium length yellow-brown setae; setae of antennae, tibiae and tarsi little shorter; setation dense,

but sparse on pronotum. Body length: 14.5–15.4 mm; humeral width: 3.1–3.4 mm.

Head (Fig. 2) widest across compound eyes, with dense irregular punctures, diameter of punctures more than spacing of punctures. Compound eyes very large, contiguous dorsally. Clypeus and labrum flattened; labrum short, anterior margin slightly emarginate in middle. Frons with one sub-rounded depression between antennal fossae; occiput abruptly, transversely concave; genae reduced, short. Antennae (Fig. 4) long, extending back to near middle of elytra; scape slightly and gradually widened apically; pedicel short, approximately 1/4 length of scape; flagellomere I approximately twice as long as pedicel and slightly shorter than flagellomere II; flagellomeres III–V subequal in length and slightly longer than II, flagellomeres VI–VIII subequal in length and approximately as long as scape; flagellomere IX very long, approximately as long as preceding four flagellomeres; rami of flagellomeres extremely elongate and subcylindrical, at least 4.0 \times length of respective flagellomere; rami of flagellomeres V–VIII with gradually decreasing lengths, ramus of flagellomere VIII approximately as long as flagellomere IX.


Pronotum (Fig. 6) subcampanulate, maximum width slightly narrower than head, length approximately 0.89 \times width; posterior 2/3 of lateral margins subparallel; disc shining, sparsely covered with small punctures, diameter of punctures distinctly less than spacing of punctures; disc with conspicuous longitudinal mesal groove, distinct depression at centre of base and transverse groove along posterior margin. Scutellar shield rounded posteriorly, depressed at centre of base, sparsely punctate. Legs slender; prothoracic tarsomeres I and V subequal in length, II–IV gradually shorter; mesothoracic tarsomere I longest, V second longest, II–IV gradually shorter; metathoracic tarsomere I longest, IV second longest, II–III gradually shorter. Pretarsal claws simple.

Posterior margin of abdominal sternites III–VI subparallel, posterior margin of sternite VII almost straight, VIII with posterior margin acutely emarginate mesally. Parameres elongate, approximately twice as long as phallobase (Figs 7, 8), more than basal 9/10 of parameres fused in dorsal view (Fig. 7). Penis elongate, somewhat dorsoventrally flattened, abruptly narrowed apically and produced into a bluntly recurved hook (Figs 9–11).

Female: Similar to male, except body larger, length 17.2 mm, humeral width 4.2 mm; head slightly narrower than pronotum; compound eyes relatively small, not dorsally contiguous (OI = 8.1); genae slightly prominent (Fig. 3); antennal rami distinctly shorter than those of male, each longer than that of preceding flagellomere, but less than twice as long; ramus of flagellomere I sub-triangular (Fig. 5); pronotum slightly wider than that of male, aspect ratio ca. 0.84; posterior margin of abdominal sternites VII and VIII roundly convex.

Etymology. The specific name is derived from the name of the type locality.

Distribution. China: Sichuan.



Figures 1–6. *Dendroides huanglong* Pan, sp. nov., from Sichuan, China. 1. Habitus; 2, 3. Head; 4, 5. Antennae; 6. Pronotum. 1, 2, 4, 6. Male, holotype; 3, 5. Female, paratype. Scale bars: 5 mm (1); 1 mm (2–6).

Key to the species of *Dendroides* from Palaearctic Region

- 1 Body unicoloured, piceous. Japan..... *D. lesnei* Blair, 1914
- Body reddish testaceous, but antennae and most of legs black (Fig. 1)..... 2
- 2 Body smaller, length 12.0–13.0 mm; eyes relatively smaller, not dorsally contiguous in male and with wider range in female (OI = 25.8–26.8) (fig. 1 in Medvedev (1977); fig. 7 in Young (2016)); in female, antennal flagellomere I with ramus indistinct, rami of flagellomeres II–VIII each shorter than that of preceding flagellomere (fig. 6 in Medvedev (1977)). Russia (Far East), China (Heilongjiang, Jilin), North Korea..... *D. ussuriensis* L.N. Medvedev, 1977
- Body larger, length 14.5–17.2 mm; eyes dorsally contiguous in male (Fig. 2) and short separated in female (OI = 8.1) (Fig. 3); females with antennal flagellomere I produced into a distinct ramus, rami of flagellomeres III–VII each longer than that of preceding flagellomere (Fig. 5). China (Sichuan)..... *D. huanglong* Pan, sp. nov.



Figures 7–11. Male genitalia of *Dendroides huanglong* Pan, sp. nov., from Sichuan, China. 7, 8. Tegmen: 7. Dorsal view; 8. Lateral view; 9–11. Penis: 9. Lateral view; 10. Dorsal view; 11. Apex, lateral view. Scale bars: 1 mm (7–10); 0.5 mm (11).

Discussion

In the preliminary molecular phylogenetic results for selected taxa of Pyrochroidae, based on mtDNA *COI* barcode sequences (Pan et al. 2021; Gao et al. 2024), *Dendroides* formed a sister group with the clade of

the genera *Pseudopyrochroa* Pic, 1906 and *Schizotus* Newman, 1838. These three genera share the following morphological characters of the adult: the temples are reduced and not prominent; the antennal pedicel is short and apically widened; the male genitalia have the parameres fused for most or all of their length (usually

more than 9/10 their length). Based on the above, we propose that the genera *Dendroidopsis* Young, 2004, *Neopyrochroa* Blair, 1914 and *Sinodendroides* Young, 2005 [for descriptions and figures, see Young (1975, 2004, 2005b)] also belong to the clade *Dendroides* + *Pseudopyrochroa* + *Schizotus*.

However, as mentioned in Pan et al. (2021), this relationship is not entirely congruent with that suggested by comparative morphological results of larvae, pupae and adults (for details, see Pan et al. (2021)); it needs to be expanded, that the penis with recurved apical hooks occurs not only in *Pseudopyrochroa*, but also *Dendroides*, at least in *D. huanglong* sp. nov. and *D. canadensis* Latreille, 1810, as well as *Neopyrochroa flabellata* (Fabricius, 1787).

If only adults are considered, those of *Dendroides* differ from *Pseudopyrochroa* and *Schizotus* as follows: frons lacking cranial pits; compound eyes very large, especially in male, dorsally contiguous or nearly so; rami of antennal flagellomeres very long, in usual 4.0 × longer than the length of the respective flagellomere; pronotum slightly wider than long; aedeagal parameres very elongate, more than twice as long as phallobase.

Acknowledgements

We wish to thank Mr Cheng-Tao Wang who collected the specimens in the field and his supervisor, Prof. Yu-Xia Yang (both Hebei University, Baoding, China) who presented the specimens to us. Thanks are extended to Prof. Fu-Ming Shi (Hebei University, Baoding, China) and the staff of Huanglong National Nature Reserve, Sichuan for organizing the expedition. We are also grateful to Prof. Darren A. Pollock (Eastern New Mexico University, Portales, USA), Prof. Hiroyuki Yoshitomi (Ehime University, Matsuyama, Japan) and Dr Dmitry Telnov (Natural History Museum, London, UK) for constructive comments.

This study was supported by the Special Program for Basic Resource Investigation of the Ministry of Science and Technology, China (No. 2022FY202100) and Baseline Survey of Huanglong Natural Reserve (No. N5132112023000495).

References

Blair KG (1914) A revision of the family Pyrochroidae (Coleoptera). Annals and Magazine of Natural History 13(75): 310–326. <https://doi.org/10.1080/00222931408693484>

Campbell JM, Marshall JD (1964) The ocular index and its application to the taxonomy of the Alleculidae (Coleoptera). The Coleopterists Bulletin 18(2): 42. <https://doi.org/10.5962/p.372397>

Gao Q, Young DK, Zhan ZH, Cheng HY, Pan Z (2024) Revision of *Eupyrochroa* Blair, 1914 (Coleoptera: Pyrochroidae) based on morphological comparison and molecular phylogenetics. Zootaxa 5406(2): 238–252. <https://doi.org/10.11646/zootaxa.5406.2.2>

Medvedev LN (1977) Ognetsvetki (Coleoptera, Pyrochroidae) fauna SSSR. Entomofauna Dal'nego Vostoka, Trudy Biologo-pochvennogo instituta, Novaya seriya 46(149): 47–51. [in Russian]

Pan Z, Duan JC, Gao Q, Young DK (2021) The adult, larva, and pupa of a new *Pseudopyrochroa* (Coleoptera: Pyrochroidae: Pyrochroinae) from China, with molecular phylogenetic inferences. Insects 12: 1089. <https://doi.org/10.3390/insects12121089>

Van Dyke EC (1928) New species of heteromorous Coleoptera. Bulletin of Brooklyn Entomological Society 23: 251–262.

Young DK (1975) A revision of the family Pyrochroidae (Coleoptera: Heteromera) for North America based on the larvae, pupae, and adults. Contributions of the American Entomological Institute 11(3): 1–39. <https://doi.org/10.25335/zj0y-nf23>

Young DK (2004) *Dendroidopsis* and *Frontodendroidopsis*: two new Asian genera of fire-colored beetles (Coleoptera: Pyrochroidae: Pyrochroinae), with descriptions of two new species of *Dendroidopsis*, and lectotype designations for *Dendroides assamensis* Blair and *Pseudopyrochroa binhana* Pic. Oriental Insects 38: 197–217. <https://doi.org/10.1080/00305316.2004.10417387>

Young DK (2005a) *Dendroides nakabusana* Kôno: proposed new junior synonym of *Dendroides lesnei* Blair (Coleoptera: Pyrochroidae: Pyrochroinae), with lectotype designation for *Dendroides lesnei*. Oriental Insects 39: 89–92. <https://doi.org/10.1080/00305316.2005.10 417420>

Young DK (2005b) *Sinodendroides chinensis*: a new genus and species of fire-colored beetle (Coleoptera: Pyrochroidae: Pyrochroinae) from China. Oriental Insects 39: 93–100. <https://doi.org/10.1080/00305316.2005.10417421>

Young DK (2016) *Pyroghatsiana*: A new genus of fire-colored beetles (Coleoptera: Pyrochroidae: Pyrochroinae) from the Southern Ghats, India. Insecta Mundi 0468: 1–8.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Deutsche Entomologische Zeitschrift \(Berliner Entomologische Zeitschrift und Deutsche Entomologische Zeitschrift in Vereinigung\)](#)

Jahr/Year: 2025

Band/Volume: [NF_72](#)

Autor(en)/Author(s): Yang Xin-Mei, Young Daniel K., Yang Bao-Lin, Pan Zhao

Artikel/Article: [A new Dendroides species from China \(Coleoptera, Pyrochroidae\), with a key to the Palaearctic species 15-19](#)