



Research article

urn:lsid:zoobank.org:pub:266D4101-D1C7-4150-8EAD-B87EAE05E694

Description of a new species of *Paracrobeles* Heyns, 1968 (Nematoda, Rhabditida, Cephalobidae) from Kelso Dunes, Mojave National Preserve, California, USA

Sven BOSTRÖM¹ & Oleksandr HOLOVACHOV²

^{1,2}Department of Zoology, Swedish Museum of Natural History, Box 50007,
SE-104 05 Stockholm, Sweden.

¹ E-mail: sven.bostrom@nrm.se (corresponding author)

² E-mail: oleksandr.holovachov@nrm.se

¹ urn:lsid:zoobank.org:author:528300CC-D0F0-4097-9631-6C5F75922799

² urn:lsid:zoobank.org:author:89D30ED8-CFD2-42EF-B962-30A13F97D203

Abstract. A new species of *Paracrobeles*, *P. kelsodunensis* sp. nov. is described from the Kelso Dunes area, Mojave National Preserve, southern California. *Paracrobeles kelsodunensis* sp. nov. is particularly characterised by a body length of 469–626 µm in females and 463–569 µm in males; lateral field with four incisures, extending almost to tail terminus; three pairs of asymmetrical lips, separated by U-shaped primary axils with two long guarding processes, each lip usually with four tines along its margin; three long labial probolae, deeply bifurcated, with slender prongs without tines; metastegostom with a strong anteriorly directed dorsal tooth; pharyngeal corpus anteriorly spindle-shaped, posteriorly elongate bulbous with dilated lumen; spermatheca 24–87 µm long; postvulval uterine sac 60–133 µm long; vulva in a sunken area; spicules 33–38 µm long; and male tail with a 5–8 µm long mucro. The generic diagnosis is emended on the basis of recently described species and a key to the species of *Paracrobeles* is provided.

Keywords. Morphology, new species, *Paracrobeles*, SEM, taxonomy.

Boström S. & Holovachov O. 2015. Description of a new species of *Paracrobeles* Heyns, 1968 (Nematoda, Rhabditida, Cephalobidae) from Kelso Dunes, Mojave National Preserve, California, USA. *European Journal of Taxonomy* 117: 1–11. <http://dx.doi.org/10.5852/ejt.2015.117>

Introduction

The genus *Paracrobeles* was erected by Heyns (1968), with *Paracrobeles laterellus* Heyns, 1968 as type species, for a cephalobid from South Africa having: pharyngeal metacarpus developed into an elongate median bulb with a large chamber; labial probolae long, bifurcate, sharply pointed, prongs without fringe; and cephalic probolae prominent, sharply pointed without fringe. *P. laterellus* has also been reported from Namibia by Rashid *et al.* (1990). Since then three new species have been described, viz. *P. psammophilus* Navarro & Lluch, 1999 from Spain, *P. mojavicus* Taylor, Baldwin & Mundo-Ocampo, 2004 from California, USA and *P. deserticola* Abolafia, Divsalar, Panahi & Shokoohi, 2014 from Iran. *P. psammophilus* was also recorded from Italy by Orselli & Vinciguerra (2002), who presented SEM

pictures of the species. A population of *Paracrobeles* from Kelso Dunes, Mojave National Preserve, southern California, is described here from studies by light and scanning electron microscope. It is close to *P. mojavius*, but differs in some characters and is thus considered to represent a new species. The generic diagnosis is emended on the basis of recently described species and a key to the species of *Paracrobeles* is provided.

Materials and methods

During a tour in the Kelso Dunes area, Mojave National Preserve, southern California, the junior author collected several samples of sand in the rhizosphere of different species of desert plants along the southern slope of the dunes. Nematodes were recovered from sand samples using a modified Baermann funnel extraction technique. For LM, specimens were relaxed by gentle heat, fixed in cold 4% formaldehyde solution, transferred to pure glycerine by a slow evaporation method and mounted on permanent slides in glycerine with paraffin wax as support for the coverslip. For SEM, specimens were post-fixed in 1% osmium tetroxide (OsO₄) and transferred to pure acetone through an acetone/distilled water series. Specimens were critical point dried in liquid CO₂, mounted on stubs, gold-plated under vacuum to a thickness of 200 Å in an Agar High Resolution Sputter Coater Model 20, and examined in a Hitachi S-4300 SEM at an accelerating voltage of 5 kV.

Morphometric characters applied herein and their abbreviations are as defined for Cephalobidae in De Ley *et al.* (1999); terminology of the labial and cephalic region and stoma follows that of Holovachov *et al.* (2009). Type specimens are deposited in the invertebrate collections of the Department of Zoology, Swedish Museum of Natural History, Stockholm, Sweden (SMNH).

Results

Phylum Nematoda Diesing, 1861
Class Chromadorea Inglis, 1983
Order Rhabditida Chitwood, 1933
Family Cephalobidae Filipjev, 1934

Genus *Paracrobeles* Heyns, 1968

Type species

Paracrobeles laterellus Heyns, 1968

Diagnosis (emended after Holovachov *et al.* 2009)

Cuticle annulated, without distinctly annulated internal layer; annuli with longitudinal striation (tessellated). Lateral field with two wings (three or four incisures), ending near tail terminus in females and in males. Lip region weakly offset, consisting of six lips arranged in three pairs: one dorsal and two subventral. Pairs of lips separated by primary axils with one or two acute triangular guarding processes; secondary axils shallow. Cephalic probolae with three or four long and slender tines. Labial probolae deeply bifurcated without tines along the slender prongs. Six outer labial and four cephalic papilliform sensilla arranged in a cephaloboid manner. Amphidial aperture rounded, located on lateral lips. Stoma divided into cheilo-, gymno- and stegostom: cheilostom barrel-shaped, with strongly sclerotized bacilliform cheilorhabdia; gymnostom narrow tubular, as wide as stegostom, with weakly sclerotized plate-like gymnorrhabdia; stegostom consists of a funnel-shaped prostegostom and variably shaped mesostego-, metastego- and telostegostom parts. Metastegostom tooth absent or present. Pharynx cephaloboid: pharyngeal procorpus cylindrical; metacarpus elongate bulbous; lumen of metacarpus often expanded to a large triradiate chamber with seemingly sclerotized lining; isthmus narrower than metacarpus; basal pharyngeal bulb oval, with strongly developed valves. Nerve ring encircling metacarpus, metacarpus-isthmus junction or

BOSTRÖM S. & HOLOVACHOV O., New species of *Paracrobeles* from Kelso Dunes

anterior part of isthmus. Excretory pore opens at level of nerve ring. Deirids present. Female reproductive system cephaloboid; posterior part of ovary straight and relatively short; spermatheca present; postvulval uterine sac present; vulva flat with contour of body or in a depression; vagina straight or directed anteriorly. Male reproductive system cephaloboid; spicules cephaloboid, with corpus and manubrium of approximately equal width; gubernaculum plate-like; cornua crurum absent. Male genital papillae: two ventrosublateral pairs located anterior to cloaca; one ventrosublateral pair located just posterior to cloacal opening; two pairs located at middle of tail length; and three pairs (lateral, subventral and subdorsal) near tail terminus; there is a midventral papilla on anterior cloacal lip. Rectum short (about as long as anal body diameter). Phasmid openings located at about one-third to half of tail length in both sexes. Female tail conoid, straight or slightly arcuate ventrad, tail terminus pointed or finely rounded; male tail conoid, slightly arcuate ventrad, tail terminus finely rounded.

Valid species

P. deserticola Abolafia, Divsalar, Panahi & Shokoohi, 2014

P. laterellus Heyns, 1968

P. mojavicus Taylor, Baldwin & Mundo-Ocampo, 2004

P. psammophilus Navarro & Lluch, 1999

Paracrobeles kelsodunensis sp. nov.

urn:lsid:zoobank.org:act:3D3142CA-44FA-422F-9B6A-8A988B3B75EC

Figs 1–2, 3A–C, I; Table 1

Diagnosis

Paracrobeles kelsodunensis sp. nov. is characterised by a body length of 469–626 μm in females and 463–569 μm in males; cuticle coarsely annulated, annuli 3.0–4.8 μm wide at midbody; lateral field with two wings, each separated by a broad groove, extending almost to tail terminus; three pairs of asymmetrical lips, pairs of lips separated by U-shaped primary axils, with two long guarding processes emerging from the first annule, each lip with four (usually) or three (rarely) tines along its margin: two long acute, with or without one shorter in the middle, and one long acute extending along the secondary axil; three long labial probolae, deeply bifurcated, with slender prongs without tines; metastegostom with a strong, anteriorly directed dorsal tooth; pharynx corpus anteriorly spindle-shaped, posteriorly elongate bulbous with dilated lumen; isthmus narrow, demarcated by a break in muscular tissue: anteriorly with heavy musculature, posteriorly further narrowing with reduced musculature; pharyngeal corpus 2.5–3.3 times isthmus length; nerve ring and excretory pore at level of metacarpus to metacarpus-isthmus junction; spermatheca 24–87 μm long; postvulval uterine sac 60–133 μm long; vulva in a depression; spicules 33–38 μm long; and male tail with a 5–8 μm long mucro.

Etymology

The new species name refers to the place where it was found.

Material examined

USA: holotype ♀, paratypes 15 ♀♀ and 9 ♂♂, SMNH Type-8715–Type 8720, 28 Mar. 2010, California, Mojave National Preserve, Kelso Dunes, soil around roots of desert plants (34°53.698' N, 115°42.155' W; 34°53.754' N, 115°42.248' W and 34°54.226' N, 115°42.200' W), *legit* O. Holovachov & P. De Ley.

Description

Adult

Body variably arcuate when killed by heat. Cuticle coarsely annulated, annuli 3.0–4.8 μm wide at midbody. Irregular longitudinal striae give the cuticle a tiled appearance. Lateral field with two wings,

areolated, each separated by a broad groove, appearing as four incisures under LM, occupying about 20% of body diameter, extending almost to tail terminus in both sexes. Lip region weakly offset, carrying 6 + 4 papillae and two rounded amphid apertures. Three pairs of asymmetrical lips, one dorsal and two ventrolateral. Pairs of lips separated by U-shaped primary axils, with two long guarding processes emerging from the first annule. Each lip with four (commonly) or three (rarely – seen in one specimen so far) tines along its margin: two long acute, with or without one shorter in the middle, and one long acute extending along the secondary axil. Three labial probolae, 13.0–15.5 µm long, deeply bifurcated, with slender prongs without tines. Stoma about one lip region diameter long. Stomatal parts not clearly discernible. Cheilorhabdia oval in latero-median view; metastegostom with a strong, anteriorly directed dorsal tooth. Pharyngeal corpus anteriorly spindle-shaped, posteriorly elongate bulbous with dilated lumen; isthmus narrow, demarcated by a break in muscular tissue: anteriorly with heavy musculature, posteriorly further narrowing with reduced musculature; bulb oval, with valves. Nerve ring and excretory pore vary in position, from the level of metacarpus to metacarpus-isthmus junction, at 60–68% of neck length and at 57–68% of neck length, respectively. Deirids at level of isthmus, at 69–82% of neck length. Excretory canal cuticularised distally.

Female

Reproductive system monodelphic, prodelphic, in dextral position in relation to intestine. Ovary reflexed posteriorly at oviduct, ovary straight posterior to vulva. Spermatheca well developed. Postvulval uterine sac large, 1.4–3.1 times vulval body diameter (VBD) long. Vagina straight and perpendicular to body axis, about one-third to two-fifths of VBD. Vulva in a sunken area. Intra-uterine eggs 43–66 x 30–36 µm. Tail conoid, generally slightly curved ventrad, curved dorsad in some specimens, with 16–21 ventral annuli, non-annulated in terminal 6–12 µm, terminus minutely rounded. Rectum sigmoid, about half of ABD long. Phasmid openings located at about one-third to two-fifths of tail length.

Male

Similar to female in most respects, except for the sexual characters. Reproductive system monorchic, dextral in position; testis reflexed ventrad anteriorly. Spicules slender, paired and symmetrical, strongly curved ventrad; with oval manubrium and subcylindrical, gradually narrowing shaft. Gubernaculum plate-like. Genital papillae distributed as follows: two pairs ventrosublateral precloacal (at 5–11 µm and at 45–57 µm anterior to cloaca), one pair ventrosublateral adcloacal, a single midventral on anterior cloacal lip; two pairs (one ventrosublateral and one lateral) at midtail; three pairs (one lateral, one subventral and one dorsosublateral) near tail terminus. Phasmid openings located at about two-fifths of tail length, one to two annuli posterior to the lateral midtail papillae. Tail slightly curved ventrad, conoid, with a 5–8 µm long mucro and minutely rounded terminus.

Remarks

The population of *Paracrobeles kelsodunensis* sp. nov. from Kelso Dunes described here agrees in many respects with the description of *P. mojaviensis* collected from sandy soil in a lava field, Mojave Desert, California. The new species differs from *P. mojaviensis* by having a prominent, anteriorly directed, dorsal metastegostom tooth (vs no metastegostom tooth); a somewhat more anterior position of the excretory pore (at level of metacarpus or metacarpus-isthmus junction vs at level of isthmus); male tail with a 5–8 µm long mucro (vs male tail without mucro). Type specimens of *P. mojaviensis* were examined (see Fig. 3D–I) and among them, one male (Fig. 3I) was found to have a prominent metastegostom tooth. The presence of a metastegostom tooth has not been described for any other species than *P. kelsodunensis* sp. nov., which might indicate that the population described by Taylor *et al.* (2004) is a mixture of species. One possible explanation could be a polymorphism based on food sources, and a comparison between molecular characters would probably be needed to resolve the status of the *Paracrobeles* populations from the Mojave Desert.




Fig. 1. *Paracrobeles kelsodunensis* sp. nov. **A.** Pharyngeal region. **B.** Female gonad. **C.** Female tail. **D.** Male tail. Scale bar = 20 µm.




Fig. 2. *Paracrobeles kelsodunensis* sp. nov. SEM micrographs. **A.** Anterior end, left lateral view. **B.** Anterior end, semi-en face view. **C–D.** Anterior end, subventral view. **E.** Vulval region. **F.** Male tail, ventral view. **G.** Female tail, right sublateral view (arrow points at phasmid). **H.** Male tail, left lateral view (arrows point at papillae). **I.** Male tail, left lateral view (arrow points at phasmid). Scale bars: A–D = 5 μ m, E–I = 10 μ m.




Fig. 3. A–C. *Paracrobeles kelsodunensis* sp. nov. LM micrographs. A. Male anterior end, ventral side to the right. B–C. Female anterior end, ventral side to the right. D–H. *Paracrobeles mojavicus* Taylor, Baldwin & Mundo-Ocampo, 2004. D–F. Female anterior end, ventral side to the right. G–H. Male anterior end, ventral side to the right. I. *Paracrobeles* cf. *kelsodunensis* sp. nov., male anterior end, ventral side to the right. Scale bar: A–I = 10 μ m.

Table 1. Measurements (in μm) of *Paracrobeles kelsodunensis* sp. nov. from Kelso Dunes, Mojave Desert, California (presented as mean \pm s.d. and (range) or only range). * Number of annuli from anterior end to nerve ring, excretory pore and deirid, respectively; – indicates that data is not applicable.

	Holotype ♀	16 ♀♀ (incl. holotype)	9 ♂♂
Body length	626	538 \pm 51 (469–626)	514 \pm 33 (463–569)
Body diameter (BD)	42	43.0 \pm 3.2 (37–49)	37.5 \pm 2.3 (34–41)
Pharynx length	185	169 \pm 11.8 (150–188)	161.5 \pm 9.5 (147–173)
Tail length	66	54.5 \pm 6.4 (42–66)	54.0 \pm 3.9 (47–58)
Anal or cloacal diameter (ABD)	29	27.6 \pm 1.7 (25–31)	29.9 \pm 0.9 (28–31)
Vulva or Testis	391	334 \pm 29.1 (297–391)	265 \pm 31 (223–315)
V-A/T	2.8	2.8 \pm 0.3 (2.5–3.4)	–
a	14.9	12.5 \pm 1.0 (10.9–14.9)	13.7 \pm 1.2 (12.1–16.7)
b	3.4	3.2 \pm 0.2 (2.9–3.7)	3.2 \pm 0.2 (2.9–3.6)
c	9.5	10.0 \pm 0.6 (9.1–11.5)	9.6 \pm 1.0 (8.0–11.8)
c'	2.3	2.0 \pm 0.2 (1.6–2.3)	1.8 \pm 0.2 (1.6–2.1)
V or T (%)	62.5	62.0 \pm 1.2 (60–65)	51.1 \pm 5.1 (44–59)
Lip region diameter	18	17.7 \pm 0.4 (17–18)	17.1 \pm 0.6 (16–18)
Labial probolae	15.5	14.7 \pm 0.7 (13.0–15.5)	14.1 \pm 0.6 (13.0–14.5)
Stoma length	14.5	14.9 \pm 0.5 (14.5–15.5)	15.1 \pm 0.5 (14.5–15.5)
Corpus length	114	106.1 \pm 5.3 (100–119)	99.7 \pm 3.9 (94–105)
Isthmus length	39	35.9 \pm 3.5 (30–42)	33.4 \pm 2.3 (29–37)
Bulb length	30	29.6 \pm 1.2 (28–33)	29.9 \pm 1.0 (28–31)
Bulb diameter	22	22.3 \pm 1.0 (20–23)	21.0 \pm 0.9 (20–22)
Corpus/isthmus ratio	2.9	3.0 \pm 0.3 (2.5–3.3)	3.0 \pm 0.2 (2.7–3.3)
Nerve ring from anterior end	133	123.9 \pm 7.7 (112–142)	119.0 \pm 8.0 (109–137)
Excretory pore from anterior end	141	119.4 \pm 10.4 (102–142)	114.8 \pm 10.8 (101–133)
Deirid from anterior end	161	152.8 \pm 11.1 (135–163)	134.0 \pm 13.4 (110–155)
R _{NR} *	24	26.1 \pm 1.4 (23–29)	26.4 \pm 1.2 (24–28)
R _{EP} *	26	24.9 \pm 1.3 (22–27)	25.3 \pm 1.2 (24–27)
R _{DEI} *	31	31.8 \pm 2.2 (29–36)	30.8 \pm 2.2 (27–34)
Annuli width at midbody	4.0–4.8	3.0–4.8	3.4–4.0
Annuli width anteriorly	4.0–4.8	3.4–4.8	3.0–4.0
Vagina or Testis flexure length	14.5	13.6 \pm 0.8 (13.0–15.5)	47.7 \pm 13.0 (31–72)
Spermatheca or Spicule length	60	49.7 \pm 16.8 (24–87)	35.1 \pm 1.5 (33–38)
PUS or Gubernaculum length	129	102.1 \pm 20.4 (60–133)	19.9 \pm 1.3 (18–22)
PUS/VBD	3.1	2.6 \pm 0.5 (1.4–3.1)	–
Rectum	18	17.9 \pm 1.3 (14.5–20.5)	–
Rectum/ABD	0.6	0.6 \pm 0.1 (0.5–0.7)	–
Phasmid	20.5	19.1 \pm 1.9 (16–29)	21.5 \pm 2.5 (18–26)
Phasmid (% of tail)	31	35.5 \pm 3.4 (31–40)	40.1 \pm 2.8 (37–45)

Key to species of *Paracrobeles* (emended after Abolafia *et al.* 2014; when using the key, please also consult Table 2 for additional diagnostic information)

1. Spicules less than 40 μm long 2
 - Spicules more than 40 μm long, primary axils with single guarding process 4
2. Vulva at 58–60% of body length *P. laterellus* Heyns, 1968
 - Vulva at 60–71% of body length, primary axils with two guarding processes 3
3. Metastegostom tooth absent *P. mojaviensis* Taylor, Baldwin & Mundo-Ocampo, 2004
 - Metastegostom tooth present, anteriorly directed *P. kelsodunensis* sp. nov.

BOSTRÖM S. & HOLOVACHOV O., New species of *Paracrobeles* from Kelso Dunes

Table 2. Morphometrics of *Paracrobeles* species. Measurements in µm (except L, in mm) (emended from Abolafia *et al.*, 2014). * Measured from the illustrations or calculated from other measurements; ** times the corresponding body diameter; – indicates that data are not applicable; ¹Abolafia *et al.* (2014); ²Heyns (1968); ³Rashid *et al.* (1990); ⁴Taylor *et al.* (2004); ⁵Navarro & Lluch (1999); ⁶Orselli & Vinciguerra (2002); ⁷Present paper.

Species	n	L	a	b	c	c'	V (%)	Stoma length	Labial probolae	Excretory pore –ant. end.	Tail	Spermatheca or Spicules	Postvulval sac or Gubernaculum	Country
<i>deserticola</i> ¹	8 ♀♀	0.48–0.60	14–18	3.0–4.0	9–10	2.0–3.0	50–63	11–14	14–16	96–111	53–67	33–50	60–96 2.9–4.5**	Iran
	8 ♂♂	0.46–0.65	13–19	3.0–4.0	9–13	1.8–2	–	12–14	15–16	101–120	48–54	46–56	16–24	
<i>laterellus</i> ²	4 ♀♀	0.53–0.74	17–20	3.3–4.1	11–15	2.3*	58–60	9*	10*	106*	50–60	–	1.0–1.5**	Namibia
	6 ♂♂	0.57–0.66	17–25	3.5–3.9	11–12	2.0*	–	–	–	–	50–55	32–35/43–46	17–22	
<i>laterellus</i> ³	1 ♀	0.57	18	3.7	11	2.2	59	12	9*	107	50	26	46	Namibia
<i>mojavicus</i> ⁴	17 ♀♀	0.47–0.71	13–18	2.6–3.5	9–13	1.8–2.5	62–71	13–18	11–17	115–163	41–66	38–77	54–110	California, USA
	19 ♂♂	0.45–0.75	14–19	2.7–3.7	10–12	1.7–2.3	–	13–18	11–17	103–160	43–67	28–37	17–22	
<i>psammophilus</i> ⁵	10 ♀♀	0.36–0.55	10–17	2.9–3.7	7–11	1.9–2.7	59–63	10–14	10–12	69–117	46–53	48*	57–101	Spain
	10 ♂♂	0.42–0.53	10–18	3.1–3.6	9–12	1.6–2.2	–	10–14	–	75–107	42–53	42–57	20–39	
<i>psammophilus</i> ⁶	34 ♀♀	0.49–0.58	13–16	2.3–2.9	8–10	2.2–2.8	59–62	10–18	11–18	114–139	58–68	30–37	65–106	Italy
	42 ♂♂	0.52–0.61	12–20	2.4–3.2	8–9	1.7–2.8	–	12–19	13–17	–	56–70	68–81	28–31	
<i>kelsoadunensis</i> ⁷	16 ♀♀	0.47–0.63	11–15	2.9–3.7	9–12	1.6–2.3	60–65	14–16	13–16	102–142	42–66	24–87	60–133 1.4–3.1**	California, USA
sp. nov.	9 ♂♂	0.46–0.57	12–17	2.9–3.6	8–12	1.6–2.1	–	14–16	13–15	101–133	47–58	33–38	18–22	

4. Lips with three tines (including the ones in secondary axils); postvulval sac not swollen
..... *P. deserticola* Abolafia, Divsalar, Panahi & Shokoohi, 2014
- Lips with four tines (including the ones in secondary axils); postvulval sac swollen
..... *P. psammophilus* Navarro & Lluch, 1999

Discussion

The Mojave Desert and especially the Kelso Dunes seem to be an area with a high diversity of species of the family Cephalobidae Filipjev, 1934. Representatives of this family are mostly terrestrial and bacteria-consuming nematodes with a worldwide distribution. They seem to be especially diverse and abundant in deserts, and sand dunes appear to be a suitable habitat for cephalobids. About ten new species have been described from the Mojave Desert, which might indicate a hot spot for cephalobid diversity (De Ley 2014). Hitherto one new genus and six new species, including the one described here, and one already known species have been described from the Kelso Dunes (Boström & Holovachov 2012, 2013a, 2013b, 2014). Some of the new species described from the Mojave Desert are closely related, which might further indicate that speciation processes are going on in this area. Species of the genus *Paracrobeles* are rare inhabitants of terrestrial ecosystems and have a rather restricted distribution. So far they have been found in warm, dry sandy soils in southern Africa (South Africa and Namibia), the Mediterranean (Spain and Italy), California and Iran (Abolafia *et al.* 2014; Heyns 1968; Navarro & Lluch 1999; Orselli & Vinciguerra 2002; Rashid *et al.* 1990; Taylor *et al.* 2004). The description of the new species adds morphological data important for species identification and broadens the diagnosis of *Paracrobeles*.

Acknowledgements

The second author was supported in part by an award from the NSF Partnerships for Enhancing Expertise in Taxonomy (PEET) program grant “PEET: Training the Next Generation of Nematode Taxonomists: Applying the Tools of Modern Monography Across Free-living and Parasitic Tylenchina” (DEB-0731516). Sampling was performed by O. Holovachov and P. De Ley under permit # MOJA-2010-SCI-0003 from the National Park Service, United States Department of Interior.

References

- Abolafia J., Divsalar N., Panahi H. & Shokoohi E. 2014. Description of *Paracrobeles deserticola* sp. n. and *Nothacrobeles hebetocaudatus* sp. n. (Nematoda: Rhabditida: Cephalobidae) from Iran and the phylogenetic relationships of these two species. *Zootaxa* 3827 (1): 1–19. <http://dx.doi.org/10.11646/zootaxa.3827.1.1>
- Boström S. & Holovachov O. 2012. Description of *Chilodellus eremus* gen. n., sp. n. and *Stegelleta arenaria* sp. n. (Rhabditida: Cephalobidae) from Kelso Dunes, Mojave National Preserve, California, USA. *Journal of Nematode Morphology and Systematics* 15 (1): 21–31.
- Boström S. & Holovachov O. 2013a. Description of two new species of *Nothacrobeles* Allen & Noffsinger, 1971 (Rhabditida: Cephalobidae) from Kelso Dunes, Mojave National Preserve, California, USA. *Journal of Nematode Morphology and Systematics* 16 (1): 25–34.
- Boström S. & Holovachov O. 2013b. Description of one new species of *Heterocephalobellus* Rashid, Geraert & Sharma, 1985 (Rhabditida, Cephalobidae) from Kelso Dunes, Mojave National Preserve, California, USA and Monte desert, Usno, Argentina. *Journal of Nematode Morphology and Systematics* 16 (2): 161–166.
- Boström S. & Holovachov O. 2014. Descriptions of species of *Stegelleta* Thorne, 1938 (Nematoda, Rhabditida, Cephalobidae) from California, New Zealand and Senegal, and a revision of the genus. *European Journal of Taxonomy* 87: 1–19. <http://dx.doi.org/10.5852/ejt.2014.87>

BOSTRÖM S. & HOLOVACHOV O., New species of *Paracrobeles* from Kelso Dunes

De Ley I.T., De Ley P., Baldwin J.G., Mundo-Ocampo M. & Nadler S.A. 1999. Three new species of *Nothacrobeles* (Nematoda: Cephalobidae) from the Mojave Desert, California. *Journal of Nematology* 31 (4): 482–497.

De Ley P. 2014. Past and present highlights of nematode research in the Mojave Desert. *Mojave National Preserve Science Newsletter* May 2014: 5–10.

Heyns J. 1968. *Paracrobeles laterellus* n. gen., n. sp. from South Africa (Nematoda: Cephalobidae). *Nematologica* 14: 511–514. <http://dx.doi.org/10.1163/187529268X00192>

Holovachov O., De Ley I.T., Mundo-Ocampo M. & De Ley P. 2009. *Identification of Cephaloboidea (Nematoda)*. EUMAINE, Gent and Nematology, UC Riverside. Available from <http://www.nrm.se/download/18.9ff3752132fdaecb6800015606/CEPHALOBOIDEA.pdf> [accessed 14 Jan 2015]

Navarro P. & Lluch J. 1999. *Paracrobeles psammophilus* sp. nov. (Nematoda: Cephalobidae) from El Saler, Valencia (Spain). *Journal of Zoology* 249 (4): 481–485. <http://dx.doi.org/10.1111/j.1469-7998.1999.tb01220.x>

Orselli L. & Vinciguerra M.T. 2002. Nematodes from Italian sand dunes. 6. Two new and three rare species of Cephalobidae (Nematoda). *Nematologia mediterranea* 30: 211–220.

Rashid F., Heyns J. & Coomans A. 1990. *Paracrobeles* and *Acrobeles* species from South West Africa/Namibia with description of a new *Acrobeles* species (Nematoda: Cephalobidae). *Phytophylactica* 22 (1): 41–49.

Taylor T.M., Baldwin J.G. & Mundo-Ocampo M. 2004. *Paracrobeles mojavius* sp. n. (Nematoda: Cephalobidae) from the Mojave Desert, California. *Journal of Nematode Morphology and Systematics* 6 (2): 151–160.

Manuscript received: 9 December 2014

Manuscript accepted: 15 January 2015

Published on: 23 March 2015

Topic editor: Rudy Jocqué

Desk editor: Charlotte Thionois

Printed versions of all papers are also deposited in the libraries of the institutes that are members of the *EJT* consortium: Muséum national d'Histoire naturelle, Paris, France; Botanic Garden Meise, Belgium; Royal Museum for Central Africa, Tervuren, Belgium; Natural History Museum, London, United Kingdom; Royal Belgian Institute of Natural Sciences, Brussels, Belgium; Natural History Museum of Denmark, Copenhagen, Denmark.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [European Journal of Taxonomy](#)

Jahr/Year: 2015

Band/Volume: [0117](#)

Autor(en)/Author(s): Boström Sven, Holovachov Oleksandr

Artikel/Article: [Description of a new species of Paracrobeles Heyns, 1968 \(Nematoda, Rhabditida, Cephalobidae\) from Kelso Dunes, Mojave National Preserve, California, USA 1-11](#)