



This work is licensed under a Creative Commons Attribution 3.0 License.

Research article

<urn:lsid:zoobank.org:pub:541EA4D0-3883-4190-8E10-FF7CAC1BB035>

Predatory midges of the tribes Palpomyiini and Sphaeromiini (Diptera: Ceratopogonidae) from the Middle East, with keys and descriptions of new species

Alicja ALWIN-KOWNACKA^{1,*}, Ryszard SZADZIEWSKI² & Jacek SZWEDO³

Department of Invertebrate Zoology and Parasitology, University of Gdańsk,
59, Wita Stwosza Street, PL80-308, Poland.

* Corresponding author: alicja.alwin@biol.ug.edu.pl

² Email: ryszard.szadziewski@biol.ug.edu.pl

³ Email: jacek.szwedo@biol.ug.edu.pl

¹ <urn:lsid:zoobank.org:author:0995CC31-5CC8-46F3-BB72-A4D76919750A>

² <urn:lsid:zoobank.org:author:083FF55D-C4C0-4C7D-AE23-562619624664>

³ <urn:lsid:zoobank.org:author:36BAE161-ECCD-4D89-B383-C7CC84FF0A89>

Abstract. The Middle East biting midges of the tribes Palpomyiini (20 species in three genera) and Sphaeromiini s. lat. (six species in five genera) are reviewed. Three new species are described and illustrated: *Bezzia libanensis* Alwin & Szadziewski sp. nov., *B. sharjahi* Alwin & Szadziewski sp. nov. and *Palpomyia freidbergi* Alwin & Szadziewski sp. nov. *Bezzia aegyptia* Kieffer, 1925 is recognized as a new junior synonym of *B. albicornis* (Meigen, 1818) (syn. nov.) and *B. omanensis* Boorman & van Harten, 2002, is recognized as a junior synonym of *B. (Sivabezzia) pachypyga* Remm, 1974 (syn. nov.). Keys to the genera and species of the tribes Palpomyiini and Sphaeromiini of the Middle East are also provided.

Keywords. *Bezzia*, *Palpomyia*, new synonymy, Middle East, distribution.

Alwin-Kownacka A., Szadziewski R. & Szwedo J. 2017. Predatory midges of the tribes Palpomyiini and Sphaeromiini (Diptera: Ceratopogonidae) from the Middle East, with keys and descriptions of new species. *European Journal of Taxonomy* 318: 1–30. <https://doi.org/10.5852/ejt.2017.318>

Introduction

In our first article in a series on Ceratopogonidae of the Middle East, all faunistic studies on the family were reviewed, and keys for identification of subfamilies and tribes were provided (Alwin *et al.* 2016). In that article, we also included diagnoses and keys to the seven genera and 22 species of the tribe Ceratopogonini in this region (Alwin *et al.* 2016). Herein, we provide faunistic and taxonomic summaries of species of two other predaceous tribes (Palpomyiini and Sphaeromiini) that inhabit the Middle East.

The Middle East fauna of the tribe Palpomyiini is rich and includes 18 recorded species in the genera *Bezzia* Kieffer, 1899, *Palpomyia* Meigen, 1818 and *Phaenobezzia* Haeselbarth, 1965. These three genera

European Journal of Taxonomy 318: 1–30 (2017)

include 629 species worldwide (Borkent 2016). Conversely, the tribe Sphaeromiini includes only six species and five genera in the Middle East: *Homonelea* Kieffer, 1917, *Johannsenomyia* Malloch, 1915, *Macropeza* Meigen, 1818, *Nilobezzia* Kieffer, 1921 and *Sphaeromias* Curtis, 1829. These five genera contain 172 extant species worldwide (Borkent 2016).

Material and methods

Specimens were collected in Israel, Lebanon, the United Arab Emirates and Yemen, then subsequently dissected, mounted on microscope slides in Canada balsam and studied as described by Alwin *et al.* (2016).

Holotypes, paratypes and other specimens are deposited in the Department of Invertebrate Zoology & Parasitology, University of Gdańsk, Poland (CEIUG) and the Department of Zoology, Tel Aviv University, Israel (TAU).

Results

Class Insecta Linnaeus, 1758
Order Diptera Linnaeus, 1758
Family Ceratopogonidae Newman, 1834
Subfamily Ceratopogoninae Newman, 1834

Tribe **Palpomyiini** Enderlein, 1936

Key to genera of Middle East Palpomyiini

1. Wing with two radial cells ***Palpomyia*** Meigen, 1818
- Wing with one radial cell 2
2. Female fifth tarsomeres armed with stout ventral setae with sharp, bent tips; gonostylus of male genitalia greatly reduced in size or absent ***Phaenobezzia*** Haeselbarth, 1965
- Female fifth tarsomeres usually unarmed (some species in the subgenus *Homobezzia* have 2 partial ventrolateral rows of setae); gonostylus of male genitalia well developed ***Bezzia*** Kieffer, 1899

Genus ***Bezzia*** Kieffer, 1899

Type species

Ceratopogon ornatus Meigen, 1830; by original designation.

Diagnosis

Body nearly bare, with robust thorax. Wing with 1 radial cell; membrane without macrotrichia. Legs slender; fore femur with 0–12 ventral spines, mid and hind femora sometimes with ventral spines; fourth tarsomeres short, more or less cordiform; fifth tarsomeres of female without ventral batonnets; female claws simple, equal-sized, usually with small inner tooth. Female abdomen with paired internal tergal apodemes and usually 2 seminal capsules. Male genitalia with short tergite IX; parameres fused, usually rod-like but sometimes variable in shape; gonostylus well developed (de Meillon & Wirth 1991).

Key to males of subgenera of Middle East *Bezzia*

1. Aedeagus shield-shaped, with slender, long submedian process (Fig. 3C) 2
- Aedeagus Y-shaped, without slender, long submedian process (Fig. 1J) 3

ALWIN-KOWNACKA A. et al., Palpomyiini and Sphaeromiini biting midges from the Middle East

2. Gonocoxite elongate, slender; gonostylus short, divided into 3–4 tooth-like lobes
..... ***Pygobezzia*** Remm, 1974
- Gonocoxite short, stout; gonostylus long, slender, not divided into lobes
..... ***Sivabezzia*** Remm, 1974
3. Aedeagus bare; flagellomere 10 longer than 11 (Fig. 1C)
..... ***Bezzia*** Kieffer, 1899
- Aedeagus covered with short, spine-like setae; flagellomere 10 shorter than 11 (Fig. 2B)
..... ***Homobezzia*** Macfie, 1932

Subgenus ***Bezzia*** Kieffer, 1899

Diagnosis

Fore femur with or without ventral spines. Male flagellum with flagellomere 10 elongate, longer than distal segments 11–13. Male flagellar plume well developed. Male aedeagus without spicules on ventral membrane; gonostylus elongate, simple (Krzywiński 1995).

Key to males of *Bezzia* (*Bezzia*) of the Middle East

1. Fore femur with 1 or more ventral spines 2
- Fore femur without ventral spines 3
2. Hind femur and tibia uniformly brown; gonostylus with blunt apex
..... ***B. flavicornis*** (Staeger, 1839)
- Hind femur and tibia yellow, with brown subapical and subbasal bands; gonostylus with sharply pointed apex ***B. naseri*** Boorman & van Harten, 2002
3. Sternite IX of male genitalia very long; gonostylus with sharply pointed apex
..... ***B. libanensis*** Alwin & Szadziewski sp. nov.
- Sternite IX of male genitalia short; gonostylus with blunt apex
..... ***B. mellori*** Boorman & van Harten, 2002

Bezzia* (*Bezzia*) *flavicornis (Staeger, 1839)

Ceratopogon flavicornis Staeger, 1839: 599 (♂, ♀, Denmark).

Ceratopogon flavipalpis Winnertz, 1852: 80 (♀, Germany).

Bezzia flavipluma Kieffer, 1919: 126 (♂, Hungary).

Bezzia spinifera Goetghebuer, 1920: 106 (♂, Belgium).

Bezzia spinosula Clastrier, 1962: 100 (♂, France).

Bezzia flavipalpis – Kieffer 1901: 153 (♀, key).

Bezzia flavicornis – Edwards 1926: 423 (♀, England; = *C. flavipalpis*, *B. spinifera*?). — Goetghebuer 1934: 79 (♀, key, distribution). — Remm 1988: 30 (distribution; = *C. flavipalpis*, *B. flavipluma*, *B. spinifera*, *B. spinosula*). — Krzywiński 1995: 138 (♀, ♂, Poland, Bulgaria, Switzerland, France, Denmark).

Bezzia spinifera – Zilahi-Sebess 1940: 106 (♂, Hungary).

Diagnosis

A large orange-brown species. Scutum and scutellum darker than remainder of thorax. Wing pale; wing length 2.20 mm; costal ratio 0.80. Palpus yellowish; palpal ratio 4.50. Fore femur with 2–4 ventral spines; fore and mid femora brownish; fore and mid tibiae brown with distinct yellow subapical band; hind femur and tibia dark brown; tarsi yellow, fifth tarsomeres slightly darker; tarsal ratio of fore leg 2.10,

European Journal of Taxonomy 318: 1–30 (2017)

of mid leg 2.20, of hind leg 2.00. Abdomen pale brown. Female with 2 large ovoid seminal capsules with short necks, one larger than the other. Male gonocoxite short, straight and thick; gonostylus short, simple, with blunt apex. Parameres fused, slightly expanded in middle, with V-shaped excavation on apical portion. Aedeagus Y-shaped, with long submedian process and blunt apex (Clastrier 1962; Krzywiński 1995).

Material examined

ISRAEL: 1 ♀, Baniass, 24 Apr. 1982, F. Kaplan leg. (TAU).

Distribution

Europe (Austria, Belarus, Belgium, Bulgaria, Czech Republic, Denmark, Estonia, France, Germany, Great Britain, Hungary, Latvia, Lithuania, Netherlands, Poland, Switzerland, Ukraine), central Russia, Azerbaijan and Kyrgyzstan (Remm 1988; Szadziewski *et al.* 2013). We provide the first record of this species from Israel.

***Bezzia (Bezzia) libanensis* Alwin-Kownacka & Szadziewski sp. nov.**

[urn:lsid:zoobank.org:act:B1B97EC8-DFA5-4537-BB13-FAC1B73487C6](http://urn.lsid:zoobank.org:act:B1B97EC8-DFA5-4537-BB13-FAC1B73487C6)

Fig. 1

Diagnosis

A small dark brown species without fore femoral spines. The male genitalia are unique in having a greatly elongate sternite IX, with narrow V-shaped caudomedian cleft, and a Y-shaped aedeagus without ventral spicules and an abruptly tapered, sharply pointed apex. Female unknown.

Etymology

The specific name refers to the Latin name of Lebanon (Libanus), where the holotype was collected.

Material examined

Holotype

LEBANON: ♂, Fanar, 33°52.680' N, 35°33.547' E, 138 m a.s.l, at light, 6 May 2012, P. Dominiak leg. (CEIUG).

Description

Male

HEAD. Uniformly brown. Antenna (Fig. 1C) pale brown; flagellomeres 10–13 predominately dark brown; flagellomere 10 elongate, considerably longer than 11–13; total flagellum length 0.90 mm; antennal ratio 0.70. Palpus slender, brownish; segment 3 slender with few sensilla capitata; palpal ratio 4.00.

THORAX (Fig. 1B). Dark brown. Wing (Fig. 1A) veins pale; 1 poorly marked elongate radial cell; wing length 1.30 mm; costal ratio 0.60. Halter brown. Legs (Figs. 1D–F) with brown coxae; all femora without ventral spines; fore and mid femora yellowish, proximal ½ of venter brown and with subapical, apical dark brown bands; hind femur brown, with apical, subapical yellow bands; tibiae brown, with sub-basal, subapical yellow bands; tarsi yellow, with pale brown apices, tarsomeres 4–5 slightly darker than 1–3; tarsal ratio of fore leg 2.20, of mid leg 2.40, of hind leg 2.20.

ABDOMEN (Fig. 1G). Brown.

GENITALIA (Fig. 1G–J). Sternite IX heavily sclerotized, long, apex extending to mid-length of gonocoxites, with narrow, V-shaped caudomedian cleft. Tergite IX short, slender, with elongate cerci. Gonocoxite

ALWIN-KOWNACKA A. et al., Palpomyiini and Sphaeromiini biting midges from the Middle East




Fig. 1. *Bezzia libanensis* Alwin & Szadziewski sp. nov., ♂. **A.** Wing. **B.** Thorax. **C.** Antennal scape and flagellum. **D.** Fore leg. **E.** Mid leg. **F.** Hind leg. **G.** Male genitalia. **H.** Male genitalia, ventral view. **I.** Parameres. **J.** Aedeagus.

short, stout; gonostylus moderately long, nearly straight, slightly curved distally, apex tapered, with pointed tip. Parameres (Fig. 1I) fused, rod-like; basal arms short, broad, bifurcate with unequal-length prongs; distal portion slightly vasiform, with heavily sclerotized margins, apical $\frac{1}{3}$ narrower, more lightly sclerotized, with rounded tip. Aedeagus (Fig. 1J) with bare ventral membrane; basal arms broad, heavily sclerotized; basal arch curved, extending $\frac{1}{3}$ of total aedeagus length; distal portion short, margin heavily sclerotized, with sharply pointed apex.

Female

Unknown.

Distribution

Lebanon; known only from the type locality.

Remarks

The male antennal flagellum of this new species has an elongate flagellomere 10 and a bare aedeagus, characters typical of members of the subgenus *Bezzia*. This new species differs from all other Palearctic and Afrotropical members of the genus by its greatly elongate sternite IX, which extends to the mid-length of the gonocoxites and covers their bases.

Bezzia (Bezzia) mellori Boorman & van Harten, 2002

Bezzia mellori Boorman & van Harten, 2002: 456 (♂, figure, description, Oman).

Diagnosis

An orange-brown species. Scutum with darker band on lateral sides, humeral areas paler. Legs pale; femorotibial joints darker; femora unarmed; fore femur with darker dorsal central area near apex; mid femur with narrow, darker dorsal area on apical portion; hind femur and tibiae with vague indications of similar markings. Male genitalia with rather long gonocoxite and short, blunt gonostylus; parameres fused, slender with broad base and expanded slender, triangular, with blunt appendix and high basal arch (Boorman & van Harten 2002).

Distribution

Oman.

Bezzia (Bezzia) naseri Boorman & van Harten, 2002

Bezzia naseri Boorman & van Harten, 2002: 457 (♂, figure, description, Yemen).

Diagnosis

Scutum dark brown, scutellum paler. Legs yellowish, with brown bands; fore femur with faint dark band at apical $\frac{2}{3}$; fore and mid tibiae with dark bands at/near proximal $\frac{1}{3}$ and apex; mid femur with a single dark apical band; hind femur and tibia with wide dark sub-basal and subapical bands; fore femur armed with 2 spines. Male genitalia short, with long, evenly curved gonostylus, with pointed apical portion; parameres fused, with low basal arch, distal portion moderately stout, apex with 2 bifid pointed divergent processes; aedeagus slender, with broad base, low basal arch and blunt apex (Boorman & van Harten 2002).

Distribution

Yemen.

ALWIN-KOWNACKA A. et al., Palpomyiini and Sphaeromiini biting midges from the Middle East

Subgenus *Homobezzia* Macfie, 1932

Type species

Homobezzia nyasae Macfie, 1932; by original designation.

Diagnosis

Fore femur with or without ventral spines (all Middle East species with fore femoral spines). Male flagellomere 10 shorter than 11; flagellar plume poorly developed. Male aedeagus covered with fine setae on ventral surface; gonostylus elongate (Krzywiński 1995).

Key to males of *Bezzia (Homobezzia)* of the Middle East

1. Legs pale yellow, with dark bands; submedian process of male parameres straight
..... *B. kuhetiensis* Remm, 1967
- Legs pale brown, without dark bands; submedian process of male parameres slightly curved ventrally
..... *B. sharjahi* Alwin & Szadziewski sp. nov.

Bezzia (Homobezzia) kuhetiensis Remm, 1967

Bezzia kuhetiensis Remm, 1967: 33 (♂, ♀, Azerbaijan).

Bezzia kuhetiensis – Turgut & Kilic 2015: 14 (♀, record, Turkey).

Diagnosis

Thorax brown, scutellum paler. Head brown. Palpus yellow. Flagellomeres 1–8 yellow, 9–13 brown on distal portions. Femora pale yellow or white, with dark brown band on apex; fore femur with small smoky area near apex, with 2 ventral spines; tibiae white, with dark brown bands on apical and basal portions; fore tibia with another dark brown band on middle portion; tarsomeres white, with darker apices. Wing pale. Abdomen pale brown. Female with 2 round seminal capsules with long, thick necks. Male genitalia with long tergite IX; gonocoxite stout, moderately short, slender, curved and gonostylus with sharply pointed apex; parameres fused, base W-shaped, with long, straight, slender distal process; aedeagus triangular, slender, with low basal arch and barely visible rounded apex (Remm 1967).

Distribution

Azerbaijan, Ukraine (Remm 1988). In the Middle East recorded from Turkey (Turgut & Kilic 2015).

Bezzia (Homobezzia) sharjahi Alwin-Kownacka & Szadziewski sp. nov.

[urn:lsid:zoobank.org:act:07610F8D-C955-4FD8-851A-395BA06AFB87](http://urn.nbn.se/resolve?urn=urn:nbn:se:liu-thesis-10700)

Fig. 2

Diagnosis

This new species differs from all other Middle East congeners in having the following combination of characters: thorax orange-brown with distinctly paler scutellum; legs uniformly brownish; fore femur with 6–9 ventral spines; male gonostylus long, with pointed apex; parameres long, slender, slightly curved dorsally; female with 2 subspherical seminal capsules with very long necks.

Etymology

The specific name refers to the Sharjah Desert, where most of the type series was collected.

Material examined

Holotype

UNITED ARAB EMIRATES: ♂, adult, Sharjah Desert Park, 18 Jan. 2005, A. van Harten leg. (CEIUG).

Paratypes

UNITED ARAB EMIRATES: 3 ♀♀, 3 ♂♂, same collection data as holotype except 18–25 Jan. 2005, light trap; 3 ♀♀, 1 ♂, same collection data except 9 Mar. 2005; 1 ♀, same collection data except 30 Apr. 2005; 1 ♀, same collection data except 30 Jun. 2005; 1 ♀, 1 ♂, same collection data except 21 Jul. 2005; 1 ♀, Wadi Wurayah, light trap, 15 Jan.–22 Feb. 2009, A. van Harten leg. (CEIUG).

YEMEN: 2 ♀♀, Al Mukalla, light trap, 1. Feb. 2003, A. van Harten leg. (CEIUG).

Description

Male

HEAD. Uniformly pale brown. Antenna (Fig. 2B) pale brown; flagellar plume poorly developed; flagellomeres 2(3)–13 with paler basal halves; flagellomeres 10–13 short; total flagellum length 0.50–0.70 mm; antennal ratio 0.50–0.70. Palpus yellow, slender; third palpal segment short, with distinct capitate sensilla; palpal ratio 2.00.

THORAX. Orange-brown; scutellum paler. Wing pale, with barely visible veins; 1 poorly marked long radial cell; wing length 0.90–1.30 mm; costal ratio 0.70–0.80. Legs uniformly pale brown, tarsi slightly paler, coxae dark brown; claws small and equal on all legs; fore femur armed with 4–8 short ventral spines; tarsal ratio of fore leg 1.60–1.90, of mid leg 1.80–2.30, of hind leg 2.10–2.20.

ABDOMEN. Orange brown.

GENITALIA (Fig. 2D–F). Tergite IX short, reaching apex of gonocoxite; sternite IX short. Gonocoxite short, stout; gonostylus long, narrow, distal half curved, with sharp apex. Parameres (Fig. 2E) fused, tongue-like, slightly curved dorsally, with blunt apex. Aedeagus (Fig. 2F) triangular, with low basal arch and barely visible rounded apex.

Female

HEAD. Brown. Antennal flagellum (Fig. 2C) with short flagellomeres, brown, proximal portions yellow; total flagellum length 0.50–0.70 mm; antennal ratio 0.80–1.10. Palpus slender, yellow; third palpal segment short, rather stout, without sensory pit; palpal ratio 2.00–2.50. Mandible with 6–8 massive distal teeth and numerous smaller proximal teeth.

THORAX. Orange brown; scutellum paler. Wing (Fig. 2A) venation similar to that of male; wing length 1.70–1.80 mm; costal ratio 0.80. Femoral coloration as in male; fore femur armed with 6–9 ventral spines; fore and mid tibiae with slightly dark bases, occasionally with slightly darker central areas, and dark apices; hind tibia with slightly darker base; tarsal ratio of fore leg 1.80–2.10, of mid leg 1.90–2.50, of hind leg 2.00–2.50.

ABDOMEN. Two subspherical seminal capsules (Fig. 2G) with long, thick necks; measuring 0.04–0.05 × 0.03–0.04 mm and 0.05–0.06 × 0.04–0.05 mm.

Distribution

United Arab Emirates and Yemen.

ALWIN-KOWNACKA A. et al., Palpomyiini and Sphaeromiini biting midges from the Middle East




Fig. 2. *Bezzia sharjahi* Szadziewski & Alwin sp. nov. **A.** Wing, ♀. **B.** Antennal flagellum, ♂. **C.** Antennal flagellum, ♀. **D.** Genitalia, ♂. **E.** Parameres. **F.** Aedeagus. **G.** Seminal capsules, ♀.

Remarks

This new species is a typical member of the subgenus *Homobezzia*. It most closely resembles *B. kuhetiensis*, which has similar male genitalia and seminal capsules. However, *B. kuhetiensis* differs from *B. sharjahi* sp. nov. by its whitish legs with distinct, dark bands, the fore femur has only 2 ventral spines, and the parameres are straight.

Subgenus ***Pygobezzia*** Remm, 1974

Type species

Bezzia strobli Kieffer, 1919 (= *B. albicornis* Meigen, 1818); by original designation.

Diagnosis

Fore femur with 2–4 ventral spines. Male flagellomere 10 elongate, longer than 11–13; and the flagellar plume is well developed. Aedeagus with slender and very long median process, which is strongly bent dorsally on proximal portion. Gonocoxite slender, elongate; gonostylus reduced, very short, divided into 3–4 tooth-like lobes.

Key to males of *Bezzia* (*Pygobezzia*) of the Middle East

1. Parameres swollen on distal portion, with lateral ribs; scutellum with six large setae
..... ***B. albicornis*** (Meigen, 1818)
- Parameres swollen on proximal portion, without lateral ribs; scutellum with four large setae
..... ***B. fuliginata*** Clastrier, 1962

Bezzia* (*Pygobezzia*) *albicornis (Meigen, 1818)

Ceratopogon albicornis Meigen, 1818: 74 (♂, Germany).

Ceratopogon calceatus Haliday in Walker, 1856: 239 (Great Britain).

Ceratopogon pallidetarsatus Strobl, 1900: 171 (♀, Spain).

Bezzia strobli Kieffer, 1919: 122 (♂, ♀, Hungary).

Bezzia brevinervis Kieffer, 1919: 122 (♂, ♀, Hungary).

Bezzia trilobata Kieffer, 1922: 237 (♂, ♀, Germany).

Bezzia aegyptia Kieffer, 1925: 264 (♀, Egypt – Suez). **Syn. nov.**

Homobezzia atrata Macfie, 1944: 126 (♂, ♀, figures, description, Egypt).

Bezzia sajana Remm, 1972: 89 (♂, Russia – Siberia).

Bezzia quadridens Remm, 1972: 89 (♂, Russia).

Ceratopogon pallidetarsatus – Becker 1903: 73 (♀, Egypt).

Bezzia albicornis – Goetghebuer 1922: 58 (combination). — Szadziewski 1984: 184 (= *C. pallidetarsatus*). — de Meillon & Wirth 1987: 65 (♂, ♀, South Africa; = *C. pallidetarsatus*, *Bezzia strobli*, *Bezzia brevinervis*, *Homobezzia atrata*). — Boorman & van Harten 2002: 456 (♂, ♀, Yemen). — Szadziewski *et al.* 2011: 649 (♂, ♀, United Arab Emirates).

Bezzia atrata – Clastrier 1962: 112 (♂, ♀, descriptions, figures, Algeria).

Bezzia calceata – Tokunaga 1966: 285 (♂, ♀, Afghanistan).

non Bezzia sajana – Remm 1974: 441 (= *B. quadridens*).

Diagnosis

Entirely blackish brown species, except tarsomeres 1–3 pale. Scutellum with 6 large setae. Fore femur with 2–4 ventral spines. Male genitalia large; gonocoxite very long and slender, gonostylus short, divided into 3–4 tooth-like lobes; aedeagus with slender, very long median process that is nearly straight

ALWIN-KOWNACKA A. et al., Palpomyiini and Sphaeromiini biting midges from the Middle East

on distal portion; parameres fused, expanded on distal portion, with distinct lateral ribs, apex with 2 small processes. Female with 2 ovoid, distinctly unequal seminal capsules.

Material examined

ISRAEL: 1 ♀, Sadom, 26 May 1976, A. Freidberg leg. (TAU); 2 ♀♀, Berekhat Ya'ar, 6 Jun. 2003, A. Freidberg leg. (TAU); 1 ♀, Besor Nature Reserve, Tel Sharuhen, 31°17' N, 34°29' E, 11 May 2005, A. Freidberg leg. (TAU); 1 ♂, Herzliya, swamp, 32°10.3' N, 34°49.4' E, 11 May 2008, A. Freidberg leg. (TAU).

LEBANON: 1 ♂, Tyre, 33°16.232' N, 35°12.706' E, net, 12 May 2012, P. Dominiak leg. (CEIUG).

Distribution

Afghanistan, Algeria, Egypt, Europe (Austria, Belgium, Croatia, Czech Republic, France, Germany, Hungary, Lithuania, Poland, Slovakia, Spain, Sweden, Switzerland, Ukraine, former Yugoslavia; Szadziewski et al. 2013), United Arab Emirates, Yemen, South Africa. We provide the first records from Israel and Lebanon.

Remarks

In the Middle East and North Africa, the subgenus *Pygobezzia* is represented by two very similar species, *B. albicornis* and *B. fuliginata*. The original description of *B. aegyptia* by Kieffer (1925), based on a single female, is very poor; however, it matches the description of *B. albicornis*. Unfortunately, the female holotype of *B. aegyptia* was probably not saved by Kieffer. Accordingly, we regard *B. aegyptia* as a new junior synonym of *B. albicornis*, because the latter species is more common in the region and was reported earlier from Egypt by Becker (1903) as *Ceratoogon pallidetarsata* and by Macfie (1944) as *B. atrata*.

Bezzia (Pygobezzia) fuliginata Clastrier, 1962

Bezzia fuliginata Clastrier, 1962: 115 (♂, Serbia).

Bezzia fulginata – Remm 1967: 32 (Azerbaijan, Georgia); 1973: 351 (Hungary). — Krzywiński 1995: 182 (♂, Afghanistan, Azerbaijan, Belgium, Georgia, Poland, Serbia).

Diagnosis

Very similar to *B. albicornis*, but differs from that species in having a scutellum with 4 large setae; parameres swollen on proximal portion, without lateral ribs, with 2 distinct apical lobes; apex of median process of aedeagus strongly bent and gonocoxite with 3 ventral bristles.

Material examined

ISRAEL: 1 ♂, Enot Zukim, 7 Dec. 1992, A. Freidberg leg. (TAU); 1 ♀, Park Rosh ha'Ayin, 16 Apr. 1993, A. Freidberg and F. Kaplan leg. (TAU); 1 ♂, En Mor, 30 Jun. 1994, A. Freidberg leg. (TAU); 1 ♀, Enot Samar, 21 Jun. 1998, A. Freidberg leg. (TAU).

LEBANON: 1 ♂, Tyre, 33°15.535' N, 35°12.726' E, beach, net, 6 May 2012, P. Dominiak leg. (CEIUG).

Distribution

Afghanistan, Azerbaijan, Europe (France, Hungary, Poland, Serbia, Spain, Ukraine – Crimea), Georgia, Kazakhstan, Turkmenistan, Tajikistan and Uzbekistan (Remm 1988; Krzywiński 1995; Szadziewski et al. 2013). We provide the first records from the Middle East, from Israel and Lebanon.

Subgenus *Sivabezzia* Remm, 1974

Type species

Bezzia campanai Clastrier, 1962; by original designation.

Diagnosis

Fore femur with 2–4 ventral spines. Male flagellomere 10 elongate, longer than 11–13. Gonocoxite massive; gonostylus well developed, simple; aedeagus composed of small lamella and long median process, which is bent dorsally and directed posteriorly. Female with 2 large seminal capsules, more or less similar in size, both with long necks that are narrowed proximally.

Remarks

Females of *B. melanoflava* Clastrier, 1958 and *B. pachypyga* Remm, 1974 are very similar, with only slight differences in the shapes of their seminal capsules. In *B. melanoflava* they are broadest near the middle portion, whereas in *B. pachypyga* the widest part is near the proximal portion.

Key to species of *Bezzia* (*Sivabezzia*) of the Middle East

1. Male genitalia with broad parameres (Fig. 3B); submedian process of aedeagus longer than parameres (Fig. 3C); ventral surface of gonocoxite with distinct apical process (Fig. 3A); seminal capsule broadest on proximal portion (Fig. 3D) *B. pachypyga* Remm, 1974
- Male genitalia with parameres slender (Fig. 3F); submedian process of aedeagus shorter than parameres (Fig. 3G–H); gonocoxite without subapical process (Fig. 3E); seminal capsule broadest on middle portion (Fig. 3I) *B. melanoflava* Clastrier, 1958

Bezzia (*Sivabezzia*) *pachypyga* Remm, 1974

Fig. 3A–D

Bezzia pachypyga Remm, 1974: 441 (♂, ♀, figures, Tajikistan, Turkmenistan).

Bezzia omanensis Boorman & van Harten, 2002: 456 (♂, figures, Oman). **Syn. nov.**

Bezzia omanensis – Szadziewski *et al.* 2011: 649 (♂, record, United Arab Emirates).

Diagnosis

Male

Dark orange-brown species with brownish head; scutum and scutellum same color as rest of thorax. Antenna pale brown, except apical portion of flagellomeres 10 and 11–13, which are darker; flagellomere 10 elongate, longer than 11–13. Palpus pale brown, slender; third segment slender, slightly longer than fourth segment. Legs slender, orange-brown; mid and hind femora darker towards apical portion, without ventral spines; fore femur uniformly brownish, with 3–4 ventral spines; tibiae with a vaguely defined wide, pale central band and narrow, pale basal band; tarsomeres 1–4 yellow, 5 brownish. Male genitalia short (Fig. 3A); sternite IX short, with V-shaped caudomedian excavation; gonocoxite almost 2 × longer than broad, with distinct stout apical process on ventral surface; gonostylus simple, stout, slightly curved, with distinct apical tooth; parameres broad, shield-shaped, distal portion expanded, with massive W-shaped base and concave apex (Fig. 3B); aedeagus short, with low basal arch and armed with slender, dorsal apicomedian process, longer than parameres (Fig. 3C).

Female

Similar to male with usual sexual differences. Two large seminal capsules, widest in proximal part, both with long necks (Fig. 3D).

ALWIN-KOWNACKA A. et al., Palpomyiini and Sphaeromiini biting midges from the Middle East




Fig. 3. *Bezzia (Sivabezzia)* spp. **A–D.** *Bezzia pachypyga*, ♂ (from Remm 1974). **A.** Genitalia, ♂. **B.** Parameres. **C.** Aedeagus. **D.** Seminal capsules, ♀. – **E–I.** *Bezzia melanoflava* (from Clastrier 1958). **E.** Genitalia, ♂. **F.** Parameres. **G.** Aedeagus, ventral view. **H.** Aedeagus, lateral view. **I.** Seminal capsules, ♀.

Material examined

UNITED ARAB EMIRATES: 4 ♂♂, Sharjah Desert Park, light trap, 18–25 Jan. 2005, A. van Harten leg. (CEIUG); 4 ♀♀, 1 ♂, same collection data except 16 Jun.–21 Jul. 2005 (CEIUG); 1 ♂, 1 ♀, same collection data except 20 Oct. 2005 (CEIUG); 1 ♂, same collection data except 10 Dec. 2005 (CEIUG); 1 ♂, 7 ♀♀, same collection data except 1–30 Nov. 2008 (CEIUG); 1 ♀, same collection data except 27 Nov.–11 Dec. 2008 (CEIUG); 2 ♂♂, 5 ♀♀, same collection data except 11 Dec. 2008–6 Jan. 2009 (CEIUG); 1 ♀, same collection data except 12 Jan.–2 Feb. 2009 (CEIUG); 1 ♂, 1 ♀, Fujairah, light trap, 16–24 Feb. 2005, A. van Harten leg. (CEIUG); 1 ♀, Hatta, 30 Jan.–26 Feb. 2006, A. van Harten leg. (CEIUG); 1 ♀, Al Ajban, 2–9 Apr. 2006, A. van Harten leg. (CEIUG); 1 ♂, Waldi Bih dam, 1–15 Mar. 2007, A. van Harten leg. (CEIUG).

Distribution

Tajikistan, Turkmenistan. In the Middle East known from Oman and the United Arab Emirates.

Remarks

The coloration of the legs of Middle East specimens of *Bezzia pachypyga* is highly variable and different from that in the original description by Remm (1974). However, diagnostic characters of the highly complicated male genitalia match the original descriptions of *B. pachypyga* and its junior synonym *B. omanensis* Boorman & van Harten, 2002.

***Bezzia (Sivabezzia) melanoflava* Clastrier, 1958**
Fig. 3E–I

Bezzia melanoflava Clastrier, 1958: 235 (♂, ♀, figures, description, Senegal).

Bezzia melanoflava – Boorman & van Harten 2002: 456 (♂, ♀, figure, description, Yemen).

Diagnosis

This species differs from *B. pachypyga* in having yellowish legs with indistinct dark brown bands and femora with 2–3 spines. Male gonocoxite simple, without any process (Fig. 3E); parameres slender, rod-like; aedeagus short, with low basal arch, and dorsal, submedian processes shorter than parameres (Figs. 3F–H) (Clastrier 1958; Boorman & van Harten 2002). Female with 2 conical seminal capsules, widest on middle portion, with long necks (Fig. 3I).

Distribution

Senegal, Yemen.

Genus ***Palpomyia*** Meigen, 1818

Type species

Ceratopogon flavipes Meigen, 1804; by original designation.

Diagnosis

Wing with 2 radial cells, costa extending at least $\frac{2}{3}$ wing length, but not to apex. Fore femur slender to swollen, usually with stout ventral spines; mid and hind femora slender, often with ventral spines; fifth tarsomeres with or without 2 ventro-lateral rows of stout bristle-like setae; female claws equal, small to enlarged, with or without small, basal inner tooth. Female abdomen with paired internal tergal apodemes and 2 seminal capsules. Male genitalia with prominent cerci; gonocoxite simple or with lobe; aedeagus

ALWIN-KOWNACKA A. *et al.*, Palpomyiini and Sphaeromiini biting midges from the Middle East

conical or triangular, with well developed basal arms; parameres fused or divided, usually with well-developed basal apodemes (de Meillon & Wirth 1991).

Remarks

There are no universally recognized subgenera of *Palpomyia*; however, four species groups have been proposed for this genus: *distincta* group, *flavipes* group, *lineata* group and *tibialis* group (Grogan & Wirth 1975, 1979; Spinelli *et al.* 2009).

Key to males of Middle East *Palpomyia*

1. Gonostylus short, triangular 2
- Gonostylus long, more or less subcylindrical 3

2. Ventral surface of gonostylus covered with numerous coarse ventral setae *P. tibialis* (Meigen, 1818)
- Ventral surface of gonostylus bare *P. ebejeri* Boorman & van Harten, 2002

3. Parameres completely separated *P. schmidti* Goetghebuer, 1934
- Parameres fused 4

4. Parameres divided into 2 lobes on distal portion *P. serripes* (Meigen, 1818)
- Parameres completely fused on distal portion 5

5. Femora with ventral spines 6
- Only fore femur with ventral spines 7

6. Hind tibia with long dorsal bristles; fore and hind femora with 11–12 spines *P. buettikeri* Boorman & van Harten, 2002
- Hind tibia without long dorsal bristles; fore and hind femora with 3 spines *P. mahyoubi* Boorman & van Harten, 2002

7. Hind tibia yellow; fore femur with 3 spines *P. nakali* Boorman & van Harten, 2002
- Hind tibia dark brown; fore femur with 4–8 spines 8

8. Body with flattened setae with thin whip-like ends. Femora as well as fore and mid tibiae with dark apices. Male gonocoxite short; gonostylus slender; parameres with blunt expanded apex *P. flavipes* (Meigen, 1804)
- Body with simple bristle-like setae. Only hind femur with dark apex. Male gonocoxite elongate; gonostylus swollen on basal portion; parameres with pointed apex *P. freidbergi* Alwin & Szadziewski sp. nov.

Palpomyia distincta group

Palpomyia buettikeri Boorman & van Harten, 2002

Palpomyia buettikeri Boorman & van Harten, 2002: 458 (♂, figure, description, Saudi Arabia).

Diagnosis

Scutum dark brown. Costa reaching more than ¾ of wing length. Legs uniformly brown; fore femur with 11–12 dark ventral spines; mid femur with 4 longer, more slender spines; hind femur with about 12 very long, slender spines; hind tibia with long dorsal bristles; fifth tarsomeres without ventral setae.

European Journal of Taxonomy 318: 1–30 (2017)

Male gonocoxite short, stout, with subapical ventral lobe; gonostylus slender, curved, apex pointed; parameres fused, narrowed basally, distal half cylindrical, expanded; aedeagus triangular, with broad apex (Boorman & van Harten 2002).

Distribution

Saudi Arabia.

***Palpomyia ebejeri* Boorman & van Harten, 2002**

Palpomyia ebejeri Boorman & van Harten, 2002: 458 (♂, figure, description, Oman).

Palpomyia ebejeri – Szadziewski *et al.* 2011: 651 (♂, ♀, description, United Arab Emirates).

Diagnosis

Pale brown species. Legs brown, tarsi paler; fore femur and bases of mid and hind femora pale; male fore femur with 2–3 short spines; mid and hind femora with 1 ventral spine near apex; female fore femur with 8–10, mid femur with 5, hind femur with 7 spines. Male gonostylus short, triangular, without ventral setae; gonocoxite short, massive, base ventrally expanded; parameres apparently completely separated, apex pointed, strongly bent outwardly; aedeagus with high basal arch and expanded apex. Female with 2 small seminal capsules with short necks.

Material examined

UNITED ARAB EMIRATES: 1 ♂, Wadi Midaq, light trap, 27 Nov. 2005, A. van Harten leg. (CEIUG); 1 ♀, Hatta, light trap, 30 Jan.–26 Feb. 2006, A. van Harten leg. (CEIUG).

Distribution

Oman, United Arab Emirates.

***Palpomyia mahyoubi* Boorman & van Harten, 2002**

Palpomyia mahyoubi Boorman & van Harten, 2002: 458 (♂, figure, description, Yemen).

Diagnosis

Abdomen, scutum, scutellum, femora and tibiae dark brown, without markings. Femora slightly paler toward bases; tarsomeres 1–4 pale, tarsomeres 5 uniformly dark brown; femora with 3 ventral spines. Male gonocoxite slender, elongate, with inner expansion; gonostylus nearly straight (the illustration of the male genitalia in Boorman & van Harten (2002) is inaccurate); parameres fused, slender, apex not visible in original illustration; aedeagus with very high basal arch, apex plate-like (Boorman & van Harten 2002). Female unknown.

Distribution

Yemen.

***Palpomyia serripes* (Meigen, 1818)**

Ceratopogon serripes Meigen, 1818: 82 (♀, Germany).

Ceratopogon flavitarsis Meigen, 1838: 20 (♀, locality unknown).

Ceratopogon transfuga Staeger, 1839: 598 (♀, Denmark).

Ceratopogon tarsatus Zetterstedt, 1855: 4874 (♀, Sweden).

Palpomyia ruficeps Kieffer, 1918: 59 (♀, Tunisia).

ALWIN-KOWNACKA A. *et al.*, Palpomyiini and Sphaeromiini biting midges from the Middle East

Palpomyia serripes – Kieffer 1906: 63 (combination). — Goetghebuer 1920: 20 (♂, ♀, Belgium); 1934: 72 (♂, ♀; = *P. ruficeps*). — Edwards 1926: 422 (♂, ♀; = *C. transfuga*, *P. tarsatus*). — Szadziewski 1986: 87 (♀, redescription, distribution, Sweden). — Krzywiński 1995: 55 (♂, ♀, Poland, Sweden). — Turgut & Kilic 2015: 14 (♀, Turkey).

Diagnosis

Body blackish-brown. Palpus and flagellum brown. Legs blackish-brown; fore femur and tarsomeres 1–2 paler, darker distally; tarsomeres 3–5 brown. All femora armed with spines; fore femur with 3–6, mid femur with 0–3 and hind femur with 0–3 spines. Female with 2 round seminal capsules with distinct necks. Parameres in male genitalia broad and separated on distal half.

Distribution

Armenia, Europe (Austria, Belgium, Czech Republic, Croatia, Denmark, Estonia, Finland, France, Germany, Great Britain, Hungary, Ireland, Italy, Lithuania, Netherlands, Norway, Poland, Slovakia, Spain, Sweden, Switzerland), Tunisia, Georgia and Japan. In the Middle East known from Turkey (Szadziewski *et al.* 2013; Turgut & Kilic 2015).

Palpomyia flavipes group

Palpomyia flavipes (Meigen, 1804)

Ceratopogon flavipes Meigen, 1804: 28 (♂, locality not given).

Ceratopogon hortulanus Meigen, 1818: 81 (♀, Europe).

Ceratopogon flavipes – Walker 1856: 228 (♀, Great Britain; = *C. hortulanus*).

Palpomyia flavipes – Kieffer 1901: 157 (♂, combination). — Krzywiński 1995: 75 (♂, ♀, distribution).

Diagnosis

Body with flattened setae with thin, whip-like tips; thorax uniformly dark. Legs yellow; femora as well as fore and mid tibiae with apical brown bands; hind tibia brown; fore femur with 5–8 spines. Male genitalia with relatively short gonocoxite; gonostylus almost straight, as long as gonocoxite; parameres fused, slender, expanded apically; aedeagus triangular with low basal arch and evenly rounded apex. Females with 2 large oval seminal capsules with short necks.

Material examined

ISRAEL: 1 ♀, Dan, 21 Jul. 1983, I. Nussbaum leg. (TAU); 1 ♀, same collection data except 28 Sep. 1983 (TAU); 1 ♀, Park Hayarden, 7 May 1987, I. Yarom leg. (TAU); 2 ♀♀, same collection data except 7 May 1987, F. Kaplan leg. (TAU); 1 ♀, same collection data except 14 Apr. 1999, A. Freidberg leg. (TAU); 1 ♂, Lifta, 10 May 1987, I. Nussbaum leg. (TAU).

Distribution

Armenia, Europe (Andorra, Austria, Belarus, Belgium, Czech Republic, Denmark, Estonia, Finland, France, Germany, Great Britain, Hungary, Ireland, Italy, Latvia, Lithuania, Norway, Poland, N and NW Russia, Slovakia, Ukraine), Georgia, Kazakhstan and Turkey (Remm 1988; Szadziewski *et al.* 2013; Turgut & Kilic 2015). The above records are the first from Israel.

Palpomyia freidbergi Alwin-Kownacka & Szadziewski sp. nov.

[urn:lsid.zoobank.org:act:740DBF40-E804-477B-A437-E73110DF1287](http://lsid.zoobank.org:act:740DBF40-E804-477B-A437-E73110DF1287)

Fig. 4

Diagnosis

This new species is distinguished by the following combination of characters: males with 4–5 fore femoral ventral spines; very short sternite IX; parameres fused, curved ventrally, apex pointed; and aedeagus with very high basal arch and plate-like apex. Females with 6–9 fore femoral spines; claws without basal inner teeth; and 2 large seminal capsules.

Etymology

The species is named for Amnon Freidberg, Department of Zoology, Tel Aviv University, Israel, who kindly sent us biting midges from Israel.

Material examined

Holotype

ISRAEL: ♂, Enot Zukim, 11 Oct. 1994, A. Freidberg and F. Kaplan leg. (TAU).

Paratypes

ISRAEL: 4 ♀♀, same collection data as holotype except 22 Apr. 1998, A. Freidberg leg. (TAU); 1 ♀, En Feshkha, 11 Aug. 1986, A. Freidberg leg. (TAU); 9 ♀♀, Roch Zukim, 10 Apr. 1994, A. Freidberg and F. Kaplan leg. (TAU).

Description

Male

HEAD. Uniformly brown. Antenna pale brown, with sparse plume; flagellomeres 9–13 slightly darker than 1–8; flagellomeres 10–13 elongate; total flagellum length of 1.00 mm; antennal ratio 1.40. Palpus (Fig. 4C) yellowish, slender; third segment cylindrical, with several distinct capitate sensilla; palpal ratio 3.00.

THORAX (Fig. 4B). Brown with scutum, scutellum, katepisternum and mediotergite slightly darker; scutum, scutellum, anteroanepisternum with fine setae, katepisternum and mediotergite bare; 1 row of 3 bristles posterior to sutura transversalis; 1 bristle on scutum just anterior of scutellum; scutellum with 4 marginal bristles. Wing (Fig. 4A) with distinct veins; first radial cell about 2 × shorter than second; wing length 1.34 mm; costal ratio 0.70. Legs (Fig. 4B) slender; fore coxa yellow, mid and hind coxae brownish; fore and mid femora yellow, hind femur yellow with dark brown apex; fore femur with 4–5 stout, black ventral spines; fore and mid tibiae yellow, slightly smoky on proximal portions; hind tibia uniformly dark brown; tarsi yellow, tarsomeres 4–5 slightly darker than 1–3; tarsal ratio of fore leg 1.90, of mid leg 3.30, of hind leg 2.40.

ABDOMEN (Fig. 4B). Pale brown.

GENITALIA (Fig. 4E–G). Sternite IX (Fig. 4G) narrow. Gonocoxite slender, elongate; gonostylus swollen on basal portion, distal half slender, greatly curved, with blunt apex. Parameres (Fig. 4F) fused, base W-shaped, expanded on mid portion, apex pointed, bent ventrally. Aedeagus (Fig. 4G) triangular; basal arch very high; membrane with numerous fine microtrichia; apex plate-like.

Female

Similar to male with the usual sexual differences. Antenna uniformly pale brown; total flagellum length of 1.40–1.60 mm; antennal ratio 1.40–1.60. Palpal ratio 4.50. Mandible with 6–7 large teeth.

ALWIN-KOWNACKA A. et al., Palpomyiini and Sphaeromiini biting midges from the Middle East




Fig. 4. *Palpomyia freidbergi* Szadziewski & Alwin sp. nov. **A.** Wing, ♀. **B.** Thorax, ♂. **C.** Palpus, ♀. **D.** Fore femur, ♀. **E.** Genitalia, ♂. **F.** Parameres. **G.** Aedeagus and sternite IX. **H.** Seminal capsules.

European Journal of Taxonomy 318: 1–30 (2017)

THORAX. Coloration and arrangement of bristles as in male. Wing as in male but larger with longer costa (Fig. 4A); wing length 2.10–2.20 mm; costal ratio 0.80. Coloration of legs as in male; fore femur armed with 6–9 stout, black ventral spines (Fig. 4D); tarsal ratio of fore leg 2.10–2.40, of mid leg 3.80–4.50, of hind leg 2.30–2.50.

ABDOMEN. Pale brown, with 2 pairs of internal tergal apodemes. Two large, subequal, ovoid seminal capsules (Fig. 4H) with short necks, measuring 0.08–0.10 × 0.12–0.14 mm and 0.06–0.08 × 0.10–0.12 mm.

Distribution

Israel; known only from the type locality.

Remarks

Males of this new species have an aedeagus with an unusual plate-like apex that is similar to that of males of *P. mahyoubi* from Yemen. This newly described species differs mostly in having pale legs with a dark brown hind tibia, only the fore femur armed with ventral spines and a distinctly curved gonostylus. Males of *P. mahyoubi* have a nearly straight gonostylus, dark brown legs, and all femora have ventral spines.

***Palpomyia nakali* Boorman & van Harten, 2002**

Palpomyia nakali Boorman & van Harten, 2002: 459 (♂, figure, description, Oman).

Diagnosis

Scutum and abdomen pale orange-brown. Legs pale, non-banded, without obvious darker bands; fore femur with 3 spines. Male genitalia with gonocoxite and gonostylus long, slender; parameres fused, with slender apical portion; aedeagus broadly triangular, probably with apical cap (Boorman & van Harten 2002).

Distribution

Oman.

***Palpomyia schmidti* Goetghebuer, 1934**

Palpomyia schmidti Goetghebuer, 1934: 36 (♀, Iraq, Iran).

Palpomyia miki Goetghebuer, 1934: 91 (♀, Hungary).

Palpomyia miki – Remm 1976: 175 (♀, ♂, Russia).

Palpomyia schmidti – Szadziewski *et al.* 2009: 195 (♀, ♂, Iraq).

Diagnosis

Body pale brown. Legs yellowish; fore femur with 5–8 spines, mid and hind femora with 1–3 spines; mid and hind tibiae with numerous large setae; tarsomere 1 of mid leg with medial spines. Male gonocoxite with long ventral apodeme; gonostylus evenly bent, with pointed apex; parameres separate with swollen apices; aedeagus broadly triangular. Female with simple claws; with 2 ovoid seminal capsules with short necks (Szadziewski *et al.* 2009).

Material examined

TURKEY: 1 ♀, Aydiklar, 21 Jul. 2004, R. Dobosz leg. (CEIUG).

ALWIN-KOWNACKA A. *et al.*, Palpomyiini and Sphaeromiini biting midges from the Middle East

Distribution

Azerbaijan, Europe (Hungary, Spain, Slovakia, Ukraine), Iran, Iraq, Kazakhstan, Mongolia, Russia (S Siberia), Tajikistan, Turkey (Szadziewski *et al.* 2009).

Palpomyia tibialis group

Palpomyia tibialis (Meigen, 1818)

Ceratopogon tibialis Meigen, 1818: 82 (♀, Europe).

Palpomyia atripectus Kieffer, 1919: 101 (♀, Hungary, Serbia).

Palpomyia laticollis Goetghebuer, 1922: 55 (♂, Belgium).

Palpomyia nipponica Tokunaga, 1939: 297 (♂, ♀, Japan).

Palpomyia tibialis – Kieffer 1901: 156 (combination, in key).

Diagnosis

Dark brown species. Fore and mid legs pale brown to yellow; coxae brown; hind femur dark brown on distal $\frac{1}{4}$ (♀) or $\frac{1}{2}$ (♂), hind tibia dark. Fore femur with 6–9, mid femur with 0–2 and hind femur with 1–3 ventral spines; fifth tarsomere of hind leg with 2 rows of large setae. Male gonocoxite long, basally expanded, with small ventral bulge on distal portion covered with small setae; gonostylus triangular, with short, stout setae on ventral surface; parameres fused only on extreme basal portion, apices elongate, slender, pointed, recurved. Female with 2 ovoid seminal capsules with distinct necks.

Distribution

North America (Grogan & Wirth 1979; Borkent & Grogan 2009), Azerbaijan, N China, Europe (Austria, Belarus, Belgium, Czech Republic, Denmark, Estonia, Finland, France, Germany, Hungary, Netherlands, Norway, Poland, Romania, N and NW Russia, Serbia, Slovakia, Sweden, Ukraine and former Yugoslavia), Georgia, Japan and Tajikistan. From the Middle East known from Turkey (Remm 1988; Szadziewski *et al.* 2013; Turgut & Kilic 2015).

Genus *Phaenobezzia* Haeselbarth, 1965

Type species

Probezzia pistiae Ingram & Macfie, 1921; by original designation.

Diagnosis

Large, slender, nearly bare midges. Legs slender; femora unarmed; female fifth tarsomeres with long, stout, sharp setae with bent tips. Female claws equal, evenly curved or bent at base, each with small basal inner tooth. Wing with 1 radial cell; costa extending 0.80–0.90 of wing length. Gonostylus very small, rarely absent (de Meillon & Wirth 1991).

Phaenobezzia spekei (Macfie, 1939)

Nilobezzia spekei Macfie, 1939: 101 (♀, description, Uganda).

Phaenobezzia spekei – Boorman & van Harten 2002: 459 (♀ and ♂ recorded, diagnosis of ♀, Yemen).

Diagnosis

Scutum dark brown, scutellum paler, with 8 bristles. Fore leg yellow; tibia darker towards tip; tarsomeres 3–5 dark; mid leg similarly colored but tarsus darker; hind leg with yellow femur, dark tibia and tarsus with bristles. Female fifth tarsomeres with 3–4 pairs of sharp, black, spine-like setae with bent tips; 2

large seminal capsules with long necks (Boorman & van Harten 2002). Boorman & van Harten (2002) reported males from Yemen, but they were not diagnosed or illustrated.

Distribution

Uganda, Yemen.

Tribe Sphaeromiini Newman, 1834

Key to genera of Middle East Sphaeromiini

- | | |
|---|-------------------------------------|
| 1. Tibiae with stout bristles | 2 |
| – Tibiae without stout bristles | 3 |
| 2. Parameres fused, H-shaped; female claws with short basal external tooth | <i>Nilobezzia</i> Kieffer, 1921 |
| – Parameres fused, Y-shaped; female claws with short basal inner tooth | <i>Sphaeromias</i> Curtis, 1829 |
| 3. Wing with one radial cell | <i>Macropeza</i> Meigen, 1818 |
| – Wing with two radial cells | 4 |
| 4. Parameres separate; female claws equal on fore leg, unequal on mid and hind legs | <i>Johannsenomyia</i> Malloch, 1915 |
| – Parameres fused; female claws equal on all legs | <i>Homohelea</i> Kieffer, 1917 |

Genus *Homohelea* Kieffer, 1917

Type species

Palpomyia adjuncta Kieffer, 1913; by original designation.

Diagnosis

Large, grayish, stout-bodied midges. Legs stout; femora with 4–8 stout, black ventral spines; fifth tarsomeres of female with 2 pairs of batonnets; female claws long, equal, curved, on fore leg with basal inner teeth. Wings with 2 radial cells and long costa that extends to near wing tip. Male genitalia short, stout; gonostylus stout, with slender hooked tip; aedeagus with high basal arch and stout tip; parameres fused except basally (de Meillon & Wirth 1991). Female abdomen without ventral setal tufts on sternite VIII; 2 seminal capsules.

Homohelea telmatoscopa (Ingram & Macfie, 1921)

Schizodactylus telmatoscopus Ingram & Macfie, 1921: 353 (♂, ♀, description, figures, Ghana).

Homohelea telmatoscopa – de Meillon & Wirth 1981: 542 (key to sub-Saharan species of *Homohelea*); Boorman & van Harten 2002: 455 (♀, Yemen).

Diagnosis

Thorax uniformly dark brown; small prothoracic lobes present. Flagellum dark brown. Palpus very small; third segment not swollen, sensory pit rudimentary. Femora with 10–12 ventral spines; femora and tibiae dark brown, tarsomeres 1–4 paler; tarsomeres 5 dark brown, with 2 pairs of dark batonnets; female claws long and equal, fore claws with a large basal inner tooth. Female with 2 ovoid seminal capsules with short necks (Ingram & Macfie 1921).

ALWIN-KOWNACKA A. et al., Palpomyiini and Sphaeromiini biting midges from the Middle East

Distribution

Ghana, Yemen.

Genus ***Johannsenomyia*** Malloch, 1915

Type species

Johannsenomyia halteralis Malloch, 1915; by original designation.

Diagnosis

Slender midges. Legs long, slender, apices of femora clubbed, especially on fore femur; femora usually without ventral spines; tarsomeres 5 of females with numerous pairs of batonnets, males with 2–3 pairs of batonnets; female claws unequal on mid and hind legs, longer claw slightly curved, each claw with short, stout basal inner tooth. Wings with 2 radial cells (rarely 1); costa moderately long (female costal ratio about 0.80). Male tergite IX rounded distally; aedeagus broad, with cap-like tip; parameres separate, basal portion curved, apical portion broad, flattened (de Meillon & Wirth 1991). Female abdomen petiolate, narrow at base, sternite VIII without setal tufts; 2 large seminal capsules.

Johannsenomyia imparunguis (Becker, 1903)

Ceratopogon imparunguis Becker, 1903: 72 (♀, description, Egypt).

Ceratolophus imparunguis – Kieffer 1906: 60 (combination).

Palpomyia imparunguis – Kieffer 1917: 364 (combination).

Allohelea imparunguis – Kieffer 1925: 263 (combination).

Johannsenomyia imparunguis – Szadziewski 1984: 189 (combination, ♀, redescription).

Diagnosis

Scutellum and scutum blackish-brown. Wings pale; second radial cell 4 times as long as first. Halteres with proximal half yellow, distal half blackish-brown. Legs slender, unarmed; female with 2 equal fore claws, claws with 1 large inner basal tooth, mid and hind legs with single long claw and 3 basal teeth as in fore claws; fore and hind tarsomeres 5 with 5 pairs of batonnets. Females with 2 large, oval seminal capsules with short necks and a distinct seminal duct (Szadziewski 1984).

Distribution

Egypt.

Genus ***Macropeza*** Meigen, 1818

Type species

Macropeza albatarsis Meigen, 1818; by original designation.

Diagnosis

Large midges. Wings narrow, with 1 radial cell; female costa extends to wing tip, male radial cell shorter. Legs slender; femora unarmed; tarsomeres 5 of female with several pairs of batonnets; female claws short, curved, equal on all legs, with a short basal outer tooth. Male genitalia elongate, slender; gonocoxite elongate, gonostylus short; aedeagus short to elongate with rounded to truncate apex; parameres separate, each with heavily sclerotized, ventrally recurved, hook-like process (de Meillon & Wirth 1991). Female abdomen with pair of lateral setal tufts on sternite VIII; 2 large seminal capsules.

***Macropeza nuda* (Becker, 1903)**

Macroptilum nudum Becker, 1903: 77 (♂, ♀, description, Egypt).

Macropeza nuda – Wirth & Ratanaworabhan 1972: 216 (combination). — Szadziewski 1986: 85 (♀, redescription, Egypt).

Diagnosis

Blackish-brown midges. Scutellum with 2 lateral, 2 submedian setae. Wing with brownish radial cell. Legs brown, except tarsomeres 1–2 of fore and mid legs and tarsomeres 1–4 of hind leg paler; hind tibial comb with 7 spines; tarsomeres 1 of all legs with 2 pairs of spines; female claws stout, nearly equal, with short basal outer teeth, male claws simple. Female with 2 elongate seminal capsules with very short necks (Szadziewski 1986).

Distribution

Egypt.

Genus ***Nilobezzia*** Kieffer, 1921

Type species

Nilobezzia armata Kieffer, 1921; by original designation.

Diagnosis

Large robust midges. Wing often whitish at base, with distal or marginal infuscation; 1 or 2 radial cells; costa moderately short (female costal ratio about 0.80). Legs slender; femora slender to moderately swollen distally, females with short ventral spines, males with long, slender setae; hind tibia with long spine-like setae; female tarsomeres 5 with numerous ventral batonnets; female claws large, equal and bent at base, nearly straight distally, with short basal outer teeth. Male genitalia rotated and bent ventrally so that sternal surface is oriented caudally; gonocoxite and gonostylus poorly developed, usually fused, gonostylus short, thumb-like, immovable; aedeagus with short basal arch and cap-like tip; parameres with bulbous fused basal portion, distal halves elongate, tightly adpressed, tips slightly bulbous (de Meillon & Wirth 1991). Female abdomen with 2 small, slender sclerites and pair of setal tufts on sternite VIII; 2 seminal capsules.

Remarks

We are unable to distinguish *Nilobezzia nilotica* from *N. nigritibialis* and were not able to borrow the holotypes of either species or have access to them. Moreover, the original description of *N. nilotica*, by Kieffer (1925), is very poor and not helpful in distinguishing either species.

***Nilobezzia nilotica* (Kieffer, 1925)**

Parrotia nilotica Kieffer, 1925: 263 (♂, Egypt).

Diagnosis

Black species. Legs brown, tarsi whitish, distal tarsomeres black, femora unarmed.

Distribution

Egypt.

ALWIN-KOWNACKA A. et al., Palpomyiini and Sphaeromiini biting midges from the Middle East

Remarks

The genus *Parrotia* Kieffer, 1923 was placed as subgenus within *Nilobezzia* by Lane (1958: 25).

Nilobezzia nigritibialis (Ingram & Macfie, 1921)

Dicrobezzia nigritibialis Ingram & Macfie, 1921: 371 (♂, ♀, descriptions, figures, Ghana).

Nilobezzia nigritibialis – Clastrier & Wirth 1961: 212 (combination). — Boorman & Harten 2002: 455 (♀, description, Oman, Yemen).

Diagnosis

Very dark brown midges. Legs with femora and tibiae dark brown; tarsomeres 1–4 pale brown, tarsomeres 5 dark brown; female claws large, equal, with basal outer teeth; female tarsomeres 5 armed with 8–10 black slender batonnets. Male with moderately well developed gonocoxite, broad basally, gonostylus conical and reduced; aedeagus with high basal arch and broad, cap-like apex (Boorman & Harten 2002). Female with 2 subspherical, unequal seminal capsules.

Distribution

Ghana, Oman, Yemen (Boorman & Harten 2002).

Genus *Sphaeromias* Curtis, 1829

Type species

Sphaeromias albomarginatus Curtis, 1829; by original designation.

Diagnosis

Large, grayish pollinose midges. Legs slender; femora with numerous spines; hind tibia with long dorsal spine-like setae; female tarsomeres 5 with numerous slender batonnets; female claws large, curved, equal, with long, slender, sharp basal inner teeth. Wings with 2 radial cells; female costa long, extending nearly to wing tip. Male gonocoxite and gonostylus elongate; aedeagus with broad, cap-like apex; parameres fused, Y-shaped, distal portion becoming increasingly broader distally, apex rounded, setose (de Meillon & Wirth 1991). Female abdomen without pair of sclerites or setal tufts on sternite VIII; 2 seminal capsules.

Sphaeromias pictus (Meigen, 1818)

Ceratopogon pictus Meigen, 1818: 80 (♀, Germany).

Ceratopogon punctatus Meigen, 1830: 264 (♀, probably Germany).

Ceratopogon elegans Winnertz, 1852: 58 (♀, Poland).

Xylocrypta miricornis Kieffer, 1919: 77 (♂, Hungary).

Sphaeromias pictus – Goetghebuer 1934: 59 (combination; = *C. elegans*).

Diagnosis

Body brown, scutellum slightly paler. Legs brown, mid and hind legs darkest; fore tibia with distinct pale band at apex (poorly developed on mid tibia); tarsomeres 1–4 yellow, with more or less brown tips, tarsomeres 5 brown; fore femur slightly swollen, with 8–9 ventral spines, mid femur slender, with 5–6 spines, hind femur with 8–10 spines; tarsomeres 5 with 6 massive ventral batonnets; female claws equal, long, curved, with basal inner teeth. Female with 2 sub-equal, rounded seminal capsules with short necks, and distinct seminal duct.

Table 1. Zoogeographical elements among the Middle East Palpomyiini and Sphaeromiini.

Afrotropical	
<i>Bezzia melanoflava</i>	Senegal; Yemen
<i>Phaenobezzia spekei</i>	Uganda; Yemen
<i>Homohelea telmatoscopa</i>	Ghana; Yemen
<i>Nilobezzia nigritibialis</i>	Ghana; Oman, Yemen
Saharo-Arabian	
<i>Bezzia libanensis</i> sp. nov.	Lebanon
<i>Bezzia mellori</i>	Oman
<i>Bezzia naseri</i>	Yemen
<i>Bezzia sharjahi</i> sp. nov.	United Arab Emirates, Yemen
<i>Palpomyia buettikeri</i>	Saudi Arabia
<i>Palpomyia ebejeri</i>	Oman, United Arab Emirates
<i>Palpomyia mahyoubi</i>	Yemen
<i>Palpomyia nakali</i>	Oman
<i>Palpomyia freidbergi</i> sp. nov.	Israel
<i>Johannsenomyia imparunguis</i>	Egypt
<i>Macropeza nuda</i>	Egypt
<i>Nilobezzia nilotica</i>	Egypt
Palearctic	
<i>Bezzia albicornis</i>	Algeria, Afghanistan, Europe, South Africa; Egypt, Israel, Lebanon, United Arab Emirates, Yemen
<i>Bezzia fuliginata</i>	Afghanistan, Azerbaijan, Europe, Georgia, Kazakhstan, Turkmenistan, Tajikistan, Uzbekistan; Israel, Lebanon
<i>Bezzia flavigornis</i>	Azerbaijan, Central Russia, Europe, Kyrgyzstan; Israel
<i>Bezzia kuhetiensis</i>	Azerbaijan, Ukraine; Turkey
<i>Bezzia pachypygia</i>	Tajikistan, Turkmenistan; Oman, United Arab Emirates
<i>Palpomyia schmidti</i>	Azerbaijan, Europe, Kazakhstan, Mongolia, S Russia, Tajikistan; Iran, Iraq, Turkey
<i>Palpomyia flavipes</i>	Armenia, Georgia, Kazakhstan, Europe; Turkey, Israel
<i>Palpomyia serripes</i>	Armenia, Georgia, Europe, Tunisia; Turkey
<i>Sphaeromias pictus</i>	China, Europe, Japan, Kazakhstan, Kyrgyzstan, Russia; Israel
Holarctic (Palearctic + Nearctic)	
<i>Palpomyia tibialis</i>	North America, Azerbaijan, N China, Europe, Georgia, Japan, N and NW Russia, Tajikistan; Turkey

Material examined

ISRAEL: 1 ♀, Golan Aniam, 18 May 1983, A. Freidberg leg. (TAU); 2 ♀♀, Hula, 1 Jul. 1993, A. Freidberg leg. (TAU); 1 ♀, Berekhat Ya'ar, 14 May 2003, A. Freidberg leg. (TAU); 2 ♀♀, same collection data except 23 May 2003 (TAU); 1 ♀, En HaHoresh, 21 May 2005, A. Freidberg leg. (TAU).

Distribution

Belarus, Belgium, Great Britain, China, Czech Republic, Estonia, Finland, France, Germany, Hungary, Italy, Japan, Kazakhstan, Kyrgyzstan, Latvia, Lithuania, Poland, Russia (central, northwest and southern regions), Slovakia, Ukraine (Remm 1988; Szadziewski *et al.* 2013). We provide the first records for the Middle East, from Israel.

ALWIN-KOWNACKA A. *et al.*, Palpomyiini and Sphaeromiini biting midges from the Middle East

Discussion

In the Middle East the tribe Palpomyiini is represented by 20 species in three genera. Nine are known only from the Middle East or the Saharo-Arabian Region (Table 1). Nine species occur in the Palearctic Region, including several species also reported from the Nearctic Region, whereas only two species have been recorded from the Afrotropical Region. The tribe Sphaeromiini is represented by only six species in five genera in the Middle East. Three species are recorded from the Saharo-Arabian Region, two from the Afrotropics, and one from the Palearctic.

Acknowledgements

We are grateful to Dr. Amnon Freidberg (TAU) who kindly sent us materials from Israel. Special thanks are directed to Dr Antonius van Harten for collecting and making available to us specimens from the United Arab Emirates and Yemen, and to Professor Dany Azar (Lebanese University) for his help during field research. This is a contribution to the cooperative agreement between the Lebanese University and University of Gdańsk in the project “Origin and Evolution of Terrestrial Biodiversity of Europe and the Middle East”.

We are also grateful to Dr. William L. Grogan Jr and an anonymous referee for reviewing the manuscript and providing valuable comments.

References

- Alwin-Kownacka A., Szadziewski R. & Szwedo J. 2016. Biting midges of the tribe Ceratopogonini (Diptera: Ceratopogonidae) from the Middle East, with keys and descriptions of new species. *Zootaxa* 4079 (5): 551–572. <https://doi.org/10.11646/zootaxa.4079.5.3>
- Becker T. 1903. Ägyptische Dipteren. *Mitteilungen aus dem zoologischen Museum in Berlin* 2: 67–195.
- Boorman J. & van Harten A. 2002. Some Ceratopogonidae (Insecta: Diptera) from the Arabian Peninsula, with particular reference to the Republic of Yemen. *Fauna of Arabia* 19: 427–462.
- Borkent A. 2016. World species of biting midges (Diptera: Ceratopogonidae). Available from <http://www.inhs.illinois.edu/research/FLYTREE/CeratopogonidaeCatalog.pdf> [accessed 6 Jul. 2016].
- Borkent A. & Grogan Jr W.L. 2009. Catalog of the New World biting midges north of Mexico (Diptera: Ceratopogonidae). *Zootaxa* 2273: 1–48.
- Clastrier J. 1958. Notes sur les Cératopogonidés. IV. Cératopogonidés d’Afrique Occidentale française. *Archives de l’Institut Pasteur d’Algérie* 36: 192–258.
- Clastrier J. 1962. Notes on Ceratopogonidae. XVI. Species of the genus *Bezzia* Kieffer or related species of the Palearctic Region. *Archives de l’Institut Pasteur d’Algérie* 40: 53–125.
- Clastrier J. & Wirth W.W. 1961. Notes on Ceratopogonidae. XIII. Ceratopogonidae of the Ethiopian region. *Archives de l’Institut Pasteur d’Algérie* 39: 190–240.
- Edwards, F.W. 1926. On the British biting midges (Diptera, Ceratopogonidae). *Transactions of the Royal Entomological Society of London* 74: 389–426. <https://doi.org/10.1111/j.1365-2311.1926.tb02249.x>
- Goetghebuer M. 1920. Ceratopogoninae de Belgique. *Mémoires du Musée royal d’Histoire naturelle de Belgique* 8 (3): 1–116. Available from <http://biodiversitylibrary.org/page/42372753> [accessed 4 Apr. 2017].
- Goetghebuer M. 1922. Étude critique des *Ceratopogon* de la collection Meigen conservée au Muséum national d’Histoire naturelle de Paris. *Bulletin de la Société entomologique de Belgique* 4: 50–59.

European Journal of Taxonomy 318: 1–30 (2017)

- Goetghebuer M. 1934. A. Die Imagines. Heleidae (Ceratopogonidae). 13a. In: Lindner E. (ed.) *Die Fliegen der Palaearktischen Region*: 1–94. E. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart.
- Grogan Jr W.L. & Wirth W.W. 1975. A revision of the genus *Palpomyia* Meigen of northeastern North America (Diptera: Ceratopogonidae). *University of Maryland Agricultural Experiment Station Miscellaneous Publication* 875: 1–49.
- Grogan Jr W.L. & Wirth W.W. 1979. The North American predaceous midges of the genus *Palpomyia* Meigen (Diptera: Ceratopogonidae). *Memoirs of the Entomological Society of Washington* 8: 1–125.
- Ingram A. & Macfie J.W.S. 1921. West African Ceratopogonidae. *Annals of Tropical Medicine and Parasitology* 15: 313–377.
- Kieffer J.J. 1901. Synopse des représentants européens du groupe *Ceratopogon*, avec description d'espèces nouvelles. *Bulletin de la Société d'Histoire naturelle de Metz* 9: 143–165. Available from <http://biodiversitylibrary.org/page/33770412> [accessed 3 Apr. 2017].
- Kieffer J.J. 1906. Chironomidae. In: Wytsman P. (ed.) *Genera Insectorum* 42: 1–78. Verteneuil & Desmet, Brussels.
- Kieffer J.J. 1917. Chironomides d'Australie conservés au Musée national hongrois de Budapest. *Annales Musei Nationalis Hungarici* 15: 175–228.
- Kieffer J.J. 1918. Chironomides d'Afrique et d'Asie conservés au Museum national hongrois de Budapest. *Annales Musei Nationalis Hungarici* 16: 31–139.
- Kieffer J.J. 1919. Chironomides d'Europe conservés au Musée national hongrois de Budapest. *Annales Musei Nationalis Hungarici* 17: 1–160.
- Kieffer J.J. 1922. Nouveaux chironomides piqueurs habitant le Sleswig-Holstein. *Annales de la Société scientifique de Bruxelles* 41: 230–238. Available from <http://biodiversitylibrary.org/page/44916070> [accessed 4 Apr. 2017].
- Kieffer J.J. 1925. Chironomides d'Egypte (Dipt.). *Bulletin de la Société royale entomologique d'Egypte* 8: 244–313.
- Krzywiński J. 1995. *Rewizja Systematyczna Drapieżnych Kuczmanów Plemienia Palpomyiini Polski* (Diptera: Ceratopogonidae). PhD Thesis (manuscript). Uniwersytet Gdańsk, Gdańsk, Poland.
- Lane J. 1958. On Neotropical *Bezzia*. *Revista Brasileira de Entomologia* 8: 25–36.
- Macfie J.W.S. 1939. Ceratopogonidae. In: *Ruwenzori Expedition 1934–35* 1: 81–102. British Museum of Natural History.
- Macfie J.W.S. 1944. A new species of *Homobezzia* (Diptera, Ceratopogonidae) from Egypt. *Proceedings of the Royal Entomological Society of London B* 13: 125–126. <https://doi.org/10.1111/j.1365-3113.1944.tb00770.x>
- Meigen J.W. 1804. *Klassifikation und Beschreibung der Europäischen Zweiflügeligen Insekten* (Diptera Linn.). Vol. 1. XXX, Braunschweig.
- Meigen J.W. 1818. *Systematische Beschreibung der bekannten Europäischen Zweiflügeligen Insekten*. Part 1. Beaufort Sohn, Aachen, Germany. Available from <http://biodiversitylibrary.org/page/12741784> [accessed 4 Apr. 2017].
- Meigen J.W. 1830. *Systematische Beschreibung der bekannten Europäischen Zweiflügeligen Insekten*. Part 6. Schulzische Buchhandlung, Hamm, Germany. Available from <http://biodiversitylibrary.org/page/12754233> [accessed 4 Apr. 2017].

ALWIN-KOWNACKA A. et al., Palpomyiini and Sphaeromiini biting midges from the Middle East

- Meigen J.W. 1838. *Systematische Beschreibung der bekannten Europäischen Zweiflügeligen Insekten*. Part 7 or Supplement. Schulzische Buchhandlung, Hamm, Germany. Available from <http://biodiversitylibrary.org/page/12841287> [accessed 4 Apr. 2017].
- Meillon B. de & Wirth W.W. 1981. Subsaharan Ceratopogonidae (Diptera). VI. New species and records of South African biting midges collected by A.L. Dyce. *Annals of the Natal Museum* 24: 525–561.
- Meillon B. de & Wirth W.W. 1987. Subsaharan Ceratopogonidae (Diptera). XII. New species and records, mainly from South Africa. *Journal of the Entomological Society of Southern Africa* 50: 35–74.
- Meillon B. de & Wirth W.W. 1991. The genera and subgenera (excluding *Culicoides*) of the Afrotropical biting midges (Diptera: Ceratopogonidae). *Annals of the Natal Museum* 32: 27–147.
- Remm H. 1967. On the fauna of Ceratopogonidae (Diptera) in the Caucasus. *Tartu Riikliku Ülikooli Toimetised* 194: 3–37.
- Remm H. 1972. Novye vidy mokrecov iz Južnoj Sibiri. *Tartu Riikliku Ülikooli Toimetised* 293: 62–90.
- Remm H. 1973. Beiträge zur Ceratopogoniden-Fauna Ungarns (Diptera). *Folia Entomologica Hungarica* 26: 349–357.
- Remm H. 1974. A review of species of the genus *Bezzia* Kieffer (Diptera, Ceratopogonidae) from the fauna of the USSR. *Entomologicheskoe Obozrenie* 53: 429–442.
- Remm H. 1976. A synopsis of the *Palpomyia* of the USSR (Diptera, Ceratopogonidae). *Proceedings of the Estonian Academy of Sciences* 65: 172–197.
- Remm H. 1988. Family Ceratopogonidae. In: Soos Á. & Papp L. (eds) *Catalogue of Palaearctic Diptera* 3: 11–110. Akadémiai Kiadó, Budapest.
- Spinelli G.R., Grogan Jr W.L. & Ronderos M.M. 2009. A revision of the Patagonian predaceous midges of the genus *Palpomyia* Meigen (Diptera: Ceratopogonidae). *Insect Systematics & Evolution* 40: 43–70. <https://doi.org/10.1163/187631209X416705>
- Staeger R.C. 1839. Systematisk fortægnelse over de i Danmark hidtil fundne Diptera. *Naturhistorisk Tidsskrift* 2: 549–600.
- Strobl G. 1900. Spanische Dipteren. XI. Theil. *Wiener Entomologische Zeitung* 19: 169–174.
- Szadziewski R. 1984. Redescriptions of three species of the biting midges Ceratopogonidae (Diptera) described by Becker from Egypt. *Polish Journal of Entomology* 54: 183–194.
- Szadziewski R. 1986. Redescriptions and notes on some Ceratopogonidae (Diptera). *Polish Journal of Entomology* 56: 3–103.
- Szadziewski R., Dominiak P. & Lewańczyk A. 2009. Redescriptions of *Atrichopogon horni* Kieffer, 1925 from Sri Lanka and *Palpomyia schmidti* Goetghebuer, 1934 from Iraq (Diptera: Ceratopogonidae). *Polish Journal of Entomology* 78: 193–199.
- Szadziewski R., Gwizdalska-Kentzer M. & Gilka W. 2011. Order Diptera, family Ceratopogonidae. In: Sheikh Tahnoon Bin Zayed Al Nahyan (patron) & van Harten A. (ed.) *Arthropod Fauna of the United Arab Emirates* 4: 636–653. Dar Al Ummah Printing, Publishing, Distribution & Advertising, Abu Dhabi, United Arab Emirates.
- Szadziewski R., Borkent A. & Dominiak P. 2013. Ceratopogonidae. In: Beuk P. & Pape T. (eds) *Fauna Europaea: Diptera Nematocera*. Fauna Europaea v. 2.6.2. Available from <http://www.faunaeur.org> [accessed 4 Apr. 2017].
- Tokunaga M. 1939. Japanese biting midges of *Bezzia* and *Palpomyia* (Ceratopogonidae, Diptera). *Tenthredo* 2: 273–313.

European Journal of Taxonomy 318: 1–30 (2017)

- Tokunaga M. 1966. Some nematocerous Diptera of the north-east of Afghanistan. *Report of the Kyoto University Scientific Expedition to Karakoram and Hindukush* 8: 273–286.
- Turgut F. & Kilic A.Y. 2015. The Ceratopogonidae (Insecta: Diptera) fauna of the Central Black Sea Region in Turkey. *Turkish Journal of Zoology* 39 (6): 1071–1089. <https://doi.org/10.3906/zoo-1410-33>
- Walker F. 1856. *Insecta Britannica. Diptera*. Vol. 3. Reeve and Benham, London.
- Winnertz J. 1852. Beitrag zur Kenntniss der Gattung *Ceratopogon* Meigen. *Linnaea Entomologica* 6: 1–80.
- Wirth W.W. & Ratanaworabhan N.C. 1972. Notes on the genus *Macropeza* Meigen and description of a new species from Florida (Diptera: Ceratopogonidae). *Florida Entomologist* 55: 213–217.
- Zetterstedt J.W. 1855. *Diptera Scandinaviae Disposita et Descripta*. Vol. 12: 4547–4942. Available from <http://biodiversitylibrary.org/page/9906725> [accessed 4 Aprt. 2017].
- Zilahi-Sebess G. 1940. Magyarország Heleidái. *Folia Entomologica Hungarica* 5: 10–133.

Manuscript received: 8 February 2016

Manuscript accepted: 5 September 2016

Published on: 15 May 2017

Topic editor: Gavin Broad

Desk editor: Danny Eibye-Jacobsen

Printed versions of all papers are also deposited in the libraries of the institutes that are members of the EJT consortium: Muséum national d'Histoire naturelle, Paris, France; Botanic Garden Meise, Belgium; Royal Museum for Central Africa, Tervuren, Belgium; Natural History Museum, London, United Kingdom; Royal Belgian Institute of Natural Sciences, Brussels, Belgium; Natural History Museum of Denmark, Copenhagen, Denmark; Naturalis Biodiversity Center, Leiden, the Netherlands; Museo Nacional de Ciencias Naturales-CSIC, Madrid, Spain; Real Jardín Botánico de Madrid CSIC, Spain.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [European Journal of Taxonomy](#)

Jahr/Year: 2017

Band/Volume: [0318](#)

Autor(en)/Author(s): Alwin-Kownacka Alicja, Szadziewski Ryszard, Szwedo Jacek

Artikel/Article: [Predatory midges of the tribes Palpomyiini and Sphaeromiini \(Diptera: Ceratopogonidae\) from the Middle East, with keys and descriptions of new species 1-30](#)