



Research article

[urn:lsid:zoobank.org:pub:AA0E7BC2-9407-4EF1-A48D-EE65D0C19C08](https://zoobank.org/pub/AA0E7BC2-9407-4EF1-A48D-EE65D0C19C08)

The identification of the species of the ‘*Spilogona contractifrons* species-group’ and the ‘*Spilogona nitidicauda* species-group’ (Diptera, Muscidae) based on morphological and molecular analysis

Vera S. SOROKINA^{1,*} & Elena V. SHAIKEVICH²

¹Siberian Zoological Museum, Institute of Systematics and Ecology of Animals, Russian Academy of Sciences, Siberian Branch, Frunze str. 11, Novosibirsk 630091, Russia.

²Laboratory of Insect Genetics, N.I. Vavilov Institute of General Genetics, Russian Academy of Sciences, Gubkin Street 3, Moscow 119991, Russia.

*Corresponding author: sorokinavs@mail.ru

²Email: elenashaikevich@mail.ru

¹[urn:lsid:zoobank.org:author:394AB897-8128-49F4-806D-4DF36557C48A](https://zoobank.org/author/394AB897-8128-49F4-806D-4DF36557C48A)

²[urn:lsid:zoobank.org:author:39F9E7C8-6761-46C0-BEDE-81F396FBD5AC](https://zoobank.org/author/39F9E7C8-6761-46C0-BEDE-81F396FBD5AC)

Abstract. Muscid species of the ‘*Spilogona contractifrons* species-group’ (*Spilogona alticola* (Malloch, 1920), *S. arctica* (Zetterstedt, 1838), *S. contractifrons* (Zetterstedt, 1838), *S. orthosurstyla* Xue & Tian, 1988) and of the ‘*Spilogona nitidicauda* species-group’ (*S. nitidicauda* (Schnabl, 1911), *S. hissarensis* Hennig, 1959, *S. imitatrix* (Malloch, 1921), *S. platyfrons* Sorokina, 2018) are notoriously difficult to distinguish. In this paper, their morphological features are analysed, images of the male head, frons and abdomen of all the species are given, and the male terminalia are figured. The study of extensive material has shown that all the morphologically recognised species in each of these groups are valid species. An identification key is provided for both groups of species. To confirm the morphological differences, genetic differences in the cytochrome oxidase I gene of flies of the ‘*Spilogona contractifrons* species-group’ and of the ‘*Spilogona nitidicauda* species-group’ were analysed. It is shown that members of both groups of species have not only distinguishing morphological characters but also fixed substitutions in the DNA sequences. Since a low interspecific polymorphism is known in the Muscidae Latreille, 1802, the revealed genetic distances confirm the existence of separate species or subspecies in each of the groups studied.

Keywords. *COI* gene, DNA barcoding, flies, identification key, male terminalia.

Sorokina V.S. & Shaikevich E.V. 2018. The identification of the species of the ‘*Spilogona contractifrons* species-group’ and the ‘*Spilogona nitidicauda* species-group’ (Diptera, Muscidae) based on morphological and molecular analysis. *European Journal of Taxonomy* 484: 1–26. <https://doi.org/10.5852/ejt.2018.484>

Introduction

The genus *Spilogona* Schnabl, 1911 is one of the most speciose genera of the Muscidae Latreille, 1802 and is dominant both in the number of species and in the number of individuals in cold habitats, such

as the high-mountain and latitudinal tundra (Sorokina 2012a, 2012b, 2013a, 2013b, 2017; Sorokina & Khruleva 2012).

This genus has been most comprehensively studied in the Nearctic Region, in particular in the territory of northern Canada and Alaska (Huckett 1965). The number of arctic species of *Spilogona* known from the Nearctic Region is 88. The tundra zones of the Palaearctic Region have not been studied for a long time, and accordingly the number of known species of *Spilogona* for these territories is significantly less (66) (Hennig 1959a, 1959b, 1959c). Recent papers on this genus have increased the number of Palaearctic species to 90, whilst the number of Nearctic species has decreased to 86 (Sorokina 2018).

It should be noted that the changes in the number of species of *Spilogona* is due not only to the description of new species, but also to the synonymization of some Nearctic and Palaearctic species. The reason for the latter is that most species of *Spilogona* have a circumpolar or arctic-montane distribution, but this fact was not previously taken into account by researchers when describing new species. Moreover, the genus *Spilogona* consists of small to medium-sized, dark-coloured muscids covered in greyish to blackish-brown dusting, frequently revealing paired dark markings, which makes them very similar to each other. Furthermore, the high number of species encountered in arctic situations in combination with an often poor state of preservation of specimens collected in traps, preserved in alcohol, etc., make species identification of samples of *Spilogona* a difficult task even for the specialist. There are therefore still many problems in clarifying the taxonomic status of the various known species of *Spilogona*.

This paper is dedicated to flies of the ‘*Spilogona contractifrons* species-group’ and the ‘*Spilogona nitidicauda* species-group’. Members of these species groups are some of the most dominant in the tundra zone. Like other species of the genus, adults of these groups of species are predaceous on other small and soft-bodied insects close to running water. One of their important roles in nature is their known activity as predators on the adults and larvae of biting black flies (Simuliidae) and on other small insects (Werner & Pont 2006a, 2006b). However, adults of these species live at the same time and in the same biotopes and have very similar morphological characters, and this causes difficulty with their identification and therefore confusion in the results of field research. Moreover, there are different opinions as to the number of existing species in each of these groups of species.

The ‘*Spilogona contractifrons* species-group’ includes four morphologically very similar species: *S. contractifrons* (Zetterstedt, 1838), *S. alticola* (Malloch, 1920), *S. arctica* (Zetterstedt, 1838) and *S. orthosurstyla* Xue & Tian, 1988. The first three species are widely distributed, and consequently have given rise to discussion for many years. Opinions have varied about the presence of only two, or even one species, but not three. Collin (1930) recorded just two species, *S. contractifrons* and *S. arctica* from Greenland. Hennig (1959a: 288) and Pont (1986: 166) regarded *S. arctica* as a synonym of *S. contractifrons*. *Spilogona alticola* was previously known only from North America. In his monograph, Huckett (1965: 238) included all three as valid species but added, under the entry for *S. contractifrons*, “I am doubtful whether *alticola* Malloch can be regarded specifically as different to *contractifrons*”. Later, Michelsen (2006: 118) reinstated *S. arctica* as a valid species. In a recent paper by the first author of the present paper, four species of this group of species were reported in the Altai Mountains (Sorokina 2018).

The ‘*Spilogona nitidicauda* species-group’ included three morphologically similar species: *S. nitidicauda* (Schnabl, 1911), *S. hissarensis* Hennig, 1959 and *S. imitatrix* (Malloch, 1921). Recently, a fourth species (*S. platyfrons* Sorokina, 2018) of this group has been described (Sorokina 2018). *Spilogona hissarensis* is known only from the type series.

In the genus *Spilogona*, species limits are occasionally ambiguous and species identification based on morphological characters is difficult. In such cases, a molecular-genetic analysis can be applied. Species

identification and species delimitation are more accurate when based not only on morphological but also on molecular-genetic characters. Unfortunately, there are very few papers devoted to the DNA barcoding of the Muscidae and most of them have used the sequence data to perform phylogenetic analyses (Savage *et al.* 2004; Schuehli *et al.* 2004, 2007; Kutty *et al.* 2008, 2014; Haseyama *et al.* 2015). Nevertheless, there is one paper in which the relevance of DNA barcoding as a taxonomic tool in muscid fly identification has been established (Renaud *et al.* 2012).

The aim of this paper is to analyse genetic differences in the cytochrome oxidase I gene of flies of the '*Spilogona contractifrons* species-group' and the '*Spilogona nitidicauda* species-group', to confirm the morphological differences.

Material and methods

The material of *Spilogona* used in this study, including type material, is deposited in the Siberian Zoological Museum of the Institute of Systematics and Ecology of Animals, Russian Academy of Sciences, Siberian Branch, Novosibirsk (SZMN), the Zoological Museum of the Moscow State University, Moscow (ZMUM), the Zoological Institute, Saint Petersburg (ZISP), and the Canadian National Collection of Insects, Ottawa (CNC).

Morphological analyses

Specimens were examined using an Altami PSO745-T microscope for external morphological characters. For the dissection of the male and female terminalia, the end of abdomen was removed and boiled in a 10% KOH solution for 15–20 s. After dissection and study, the abdomen and terminalia were washed and then stored in microvials of glycerine pinned directly underneath the specimens. Pictures were made with the camera Canon EOS 600D. Illustrations were made in ink and then edited using Adobe Photoshop CS.

Molecular-genetics analyses

The molecular-genetic analysis was based on the study of the *COI* gene, the most standardized short genetic marker in an organism's DNA that is used as a taxonomic DNA barcoding method (Ratnasingham & Hebert 2007). Total DNA was extracted from samples of *Spilogona* using the DIA™ DNA Prep kit (Isogen, Russia). Polymerase chain reaction (PCR) of the cytochrome c oxidase subunit I (*COI*) locus was conducted with an Encyclo PCR kit (Evrogen, Russia) using TY-J-1460 (Simon *et al.* 1994) and COIR (Shaikevich 2007) primers. The following amplification conditions were used: initial denaturation at 94°C (5 min); 35 cycles: denaturing at 94°C (30 s), annealing at 55°C (30 s), elongation at 72°C (40 s); and final elongation at 72°C (10 min). The PCR fragments were purified from agarose gel with a Clean-Up Extraction Kit (Evrogen, Russia) and were sequenced using the BigDye Termination kit 3.1 (Applied Biosystems, USA). All sequences were deposited in the GenBank under accessions numbers MH413049–MH413064.

Specimens for genetic analysis were taken from different populations in the territories of Russia. The institution code, the voucher and the GenBank accessions number of specimens are indicated within brackets. A total of 16 specimens of 6 species of *Spilogona* was examined:

Spilogona alticola

RUSSIA: 1 ♂, Altai Republic, Kosh-Agach district, Ukok plateau, environs of Kal'dzhin-Kul'-Bas lake, 49°19' N, 87°26' E, 2400–2600 m a.s.l., 20 Jul. 2006, V. Sorokina leg. (SZMN SA13 MH413052); 1 ♂, Altai Republic, Kosh-Agach district, Ukok plateau, environs of Kal'dzhin-Kul'-Bas lake, 49°19' N, 87°26' E, 2400–2600 m a.s.l., 20 Jul. 2006, V. Sorokina leg. (SZMN SA14 MH413053).

Spilogona arctica

RUSSIA: 1 ♂, Altai Republic, Kosh-Agach district, Severo-Chuyskiy Range, valley of Aktru River, 50°06' N, 87°48' E, 2064 m a.s.l., 21 Jul. 2013, V. Sorokina leg. (SZMN SA9 MH413051); 1 ♂, Krasnoyarsk Krai, Taimyr Peninsula, 12.5 km to south from Dixon, the mouth of the river Lemberova, 73°24' N, 80°39' E, 23 Jul. 2012, A. Barkalov leg. (SZMN, SA1 MH413049); 1 ♂, Nenets AO, lower part of the river Pechora, 68°55' N, 53°50' E, 11 Jul. 2008, A. Ozerov leg. (SZMN SA4 MH413050).

Spilogona contractifrons

RUSSIA: 1 ♂, Altai Republic, Kosh-Agach district, Ukok plateau, Rodonoviy spring, upper part of Zhumaly river, 49°27' N, 88°03' E, 2410 m a.s.l., 4 Jul. 2005, T. Novgorodova leg. (SZMN SC22 MH413054); 1 ♂, Kurgan region, environs Kurgan, 55°30' N, 66°19' E, 17 Jun. 2015, V. Sorokina leg. (SZMN SC25 MH413055); 1 ♂, Nenets AO, Naryan Mar, 67°38' N, 53°00' E, 8 Jul. 2008, N. Vikhrev leg. (SZMN SC29 MH413056); 1 ♂, Nenets AO, Pakhancheskaya Bay, 68°29' N, 57°26' E, 26 Jul.–5 Aug. 2015, O. Makarova and M. Bizin leg. (SZMN SC30 MH413057).

Spilogona imitatrix

RUSSIA: 1 ♂, Altai Republic, Ongudai district, Terektinskiy mountain range, upper part of Bolshoi Yaloman River, 50°28' N, 86°19' E, 2200–2300 m a.s.l., 3 Aug. 2007, V. Sorokina leg. (SZMN SI35 MH413061).

Spilogona nitidicauda

RUSSIA: 1 ♂, Altai Republic, Kosh-Agach district, Ukok plateau, environs of Muzdy-Bulak lake, 49°17' N, 87°39' E, 2450 m a.s.l., 2 Jul. 2005, V. Sorokina leg. (SZMN SN38 MH413062); 1 ♂, Chukotka AO, Beringovskii district, 40 km SSW Beringovskii, 62°43' N, 178°55' E, 22 Jul. 2012, A. Stekolshchikov leg. (SZMN SN40 MH413063); 1 ♂, Chukotka AO, Beringovskii district, 40 km SSW Beringovskii, 62°43' N, 178°55' E, 22 Jul. 2012, A. Stekolshchikov leg. (SZMN SN41 MH413064).

Spilogona platyfrons

RUSSIA: 1 ♂, Altai Republic, Ulagan district, Shapshal'skiy mountain range, upper part of Shui River, ~50°33' N, 89°50' E, 2550–2900 m a.s.l., 24 Jul. 2007, V. Sorokina leg. (SZMN SP32 MH413058); 1 ♂, Altai Republic, Ust-Koksa district, 47 km W of Ust-Koksa, Krasnaya Mountains, 50°04' N, 85°13' E, 2000–2100 m a.s.l., 16 Jul. 2015, V. Sorokina leg. (SZMN SP33 MH413059); 1 ♂, Altai Republic, Ust-Koksa district, 47 km W of Ust-Koksa, Krasnaya Mountain, 50°04' N, 85°13' E, 2000–2100 m a.s.l., 16 Jul. 2015, V. Sorokina leg. (SZMN SP34 MH413060).

To investigate the genetic relationships of the isolates of *Spilogona*, Maximum Likelihood phylogenetic trees based on the *COI* gene nucleotide data were reconstructed; values lower than 50% were eliminated. A matrix of pairwise distances were estimated using the Maximum Composite Likelihood (MCL) approach. The analysis involved 38 nucleotide sequences and there was a total of 408 positions in the final dataset. Evolutionary analyses were conducted in MEGA6 (Tamura *et al.* 2013).

The following abbreviations are used in the text: *a* = anterior; *acr* = acrostichal; *ad* = anterodorsal; *av* = anteroventral; *d* = dorsal; *dc* = dorsocentral; *p* = posterior; *pd* = posterodorsal; *ph* = posthumeral; *post* = postsutural; *ps* = presutural; *p_v* = posteroventral; *v* = ventral; AO = Autonomous Okrug; ML = Maximum Likelihood.

The citation of some original labels is given in quotation marks, but interpretations of these data are given in square brackets.

Results

Taxonomic analysis of 'Spilogona contractifrons species-group'

Flies of this group of species can be distinguished as follows: scutum with 3 *post dc* setae, mid tibia without *pv* and with 1 *ad* seta, fore tibia with 1 *p* seta, hind femur with a row of *pv* setae, mid femur with anterior preapical seta, haltere yellow, facial edge not projecting beyond level of profrons, prementum shining, scutellum without downwardly-directed preapical setulae on upper border of declivities, abdomen plump, katapisternal setae arranged in a more or less equilateral triangle like *Coenosia* Meigen, 1826.

Four species of this group are currently recognised: the widespread *S. alticola*, *S. arctica*, *S. contractifrons*, and the East Asian *S. orthosurstyla*. Research for the current paper has included a detailed study of extensive material from the territory of Russia, from both tundra zones and mountain areas, and personal study of the bulk of Hockett's collection deposited in the Canadian National Collection, Ottawa, Canada.

The study of extensive material has confirmed the existence of four species in this group. Despite the difficulties in the recognition of species of this group, there are several helpful diagnostic characters: the shape of male sternite 5, the shape of male terminalia, and the colour of the scutum (Figs 1–2, Table 1). The colour of the scutum can be variable in different populations but the shape of the male sternite 5 is constant and is clearly visible in dry specimens (Fig. 1 G–I).

At present it is impossible to separate females of these species. Males can be distinguished by the following couplets:

1. Scutum completely dark brown (mountain populations) or light grey with indistinct stripes (arctic tundra populations); sternite 5 with a narrow median notch (Fig. 1E, H), with a wavy inner edge; in lateral view hypandrium with a protuberance; surstylus sharply curved at middle (Fig. 1K) *S. arctica* (Zetterstedt, 1838)
 - Scutum grey or brownish dusted with more or less distinct dark stripes; sternite 5 with a rather wide median notch; hypandrium without a distinct protuberance; shape of surstylus not as above 2
2. Sternite 5 with a wide rectangular median notch, inner edge more or less straight (Fig. 1F, I); scutum with distinct dark stripes; surstylus short, gently curved, about as long as cerci (Fig. 1L) *S. alticola* (Malloch, 1920)
 - Sternite 5 not as above; scutum with indistinct dark stripes, often diffuse; surstylus longer than cerci 3
3. Sternite 5 with an extended and pointed caudal margin (Fig. 2C); surstylus straight, curved only at apex (Fig. 2A) *S. orthosurstyla* Xue & Tian, 1988
 - Sternite 5 with a rounded caudal margin (Fig. 1D, G); surstylus not straight, weakly curved at middle (Fig. 1J) *S. contractifrons* (Zetterstedt, 1838)




Fig. 1. Males of *Spilogona* Schnabl, 1911. **A–C.** Male head and scutum, anterior view. **A.** *S. contractifrons* (Zetterstedt, 1838). **B.** *S. arctica* (Zetterstedt, 1838). **C.** *S. alticola* (Malloch, 1920). **D–F.** Sternite 5. **D.** *S. contractifrons*. **E.** *S. arctica*. **F.** *S. alticola*. **G–I.** Abdomen, dorsal view. **G.** *S. contractifrons*. **H.** *S. arctica*. **I.** *S. alticola*. **J–L.** Terminalia, lateral view. **J.** *S. contractifrons*. **K.** *S. arctica*. **L.** *S. alticola*. Scale bars: 1 mm.




Fig. 2. *Spilogona orthosurstyla* Xue & Tian, 1988. **A.** Terminalia, lateral view. **B.** Terminalia, dorsal view. **C.** Sternite 5. Scale bars: 0.25 mm.

Table 1. Diagnostic characters of males of ‘*Spilogona contractifrons* species-group’.

	<i>S. alticola</i> (Malloch, 1920)	<i>S. arctica</i> (Zetterstedt, 1838)	<i>S. contractifrons</i> (Zetterstedt, 1838)	<i>S. orthosurstyla</i> Xue & Tian, 1988
Colour of scutum	grey and brown dusted with distinct dark stripes (Fig. 1C)	completely dark brown (mountains) or grey dusted (arctic tundra) with indistinct dark stripes (Fig. 1B)	grey or brownish-grey dusted with brown stripes, sometimes diffuse (Fig. 1A)	brown dusted with indistinct dark stripes
Shape of sternite 5	with a wide median notch, inner edge more or less straight, with a rounded caudal margin (Fig. 1F, 1I)	with a narrow median notch, inner edge wavy, with a rounded caudal margin (Fig. 1E, 1H)	with a fairly wide median notch, the inner edge strongly curved, with a rounded caudal margin (Fig. 1D, 1G)	with a fairly wide median notch, the inner edge not curved, with extended and pointed caudal margin (Fig. 2C)
Shape of surstylus	short, gently curved (Fig. 1L)	sharply curved at middle (Fig. 1K)	weakly curved at middle (Fig. 1J)	straight, curved only at apex (Fig. 2A)
Shape of hypandrium	without processes, or with a small tubercle, narrow	with protuberance (Fig. 1K)	without protuberance	without protuberance

Family Muscidae Latreille, 1802
Subfamily Coenosiinae Verrall, 1888
Tribe Limnophorini Villeneuve, 1902
Genus *Spilogona* Schnabl, 1911

Spilogona alticola (Malloch, 1920)
Fig. 1C, F, I, L

Type locality

USA: New Hampshire.

Material examined

CANADA: 2 ♂♂, Manitoba, 4 km S Churchill, Beach Bay, 58°45' N, 94°08' W, Malaise trap, 17–21 Jul. 2007, A. Renaud leg. (SZMN); 1 ♀, same data as for preceding except: pan traps, 25–29 Jul. 2007 (SZMN); 1 ♀, Manitoba, Goose Creek Rd., Pump House, 58°38' N, 94°14' W, Malaise trap, 07–11 Aug. 2007, A. Renaud leg. (SZMN).

RUSSIA: 14 ♂♂, Altai Republic, Kosh-Agach district, Ukok plateau, environs of Muzdy-Bulak lake, 49°17' N, 87°39' E, 2400–2900 m a.s.l., sweep in vegetation, 26 Jun. 2005, 1 Jul. 2005, A. Barkalov and V. Sorokina leg. (SZMN); 2 ♂♂, Altai Republic, Kosh-Agach district, Ukok plateau, Rodonovyi spring, upper part of Zhumaly river, 49°27' N, 88°03' E, 2410 m a.s.l., sweep in vegetation, 24 Jun. 2005, V. Sorokina leg. (SZMN); 13 ♂♂, Altai Republic, Kosh-Agach district, Ukok plateau, environs of

SOROKINA V.S. & SHAIKEVICH E.V., Morphological and molecular analysis of *Spilogona*

Kal'dzhin-Kul'-Bas lake, 49°19' N, 87°26' E, 2400–2600 m a.s.l., sweep in vegetation, 20 Jul. 2006, V. Sorokina and T. Novgorodova leg. (SZMN); 1 ♂, Chukotka AO, lower part of the Anadyr river, 64°50' N, 175°58' E, 5 m a.s.l., 20 Jul. 2013, A. Barkalov leg. (SZMN).

Distribution

Holarctic Region; in the Palaearctic, from Southern Siberia to the north of the Russian Far East. Arcto-alpine species.

Spilogona arctica (Zetterstedt, 1838)

Fig. 1B, E, H, K

Type locality

GREENLAND.

Material examined

CANADA: 1 ♂, Manitoba, Goose Creek Rd., Pump House, 58°38' N, 94°14' W, Malaise trap, 10–14 Jul. 2007, A. Renaud leg. (SZMN).

GREENLAND: 1 ♂, Cape Farewell, “Pamiagdruk, Anordiuitsok” [59°46' N, 43°55' W], 15 Jul. 1970, J. Böcher leg. (SZMN); 1 ♀, same data as for preceding except: 1 Aug. 1970 (SZMN).

RUSSIA: 6 ♂♂, Altai Republic, Turochak district, 8 km S of Obogo village, south slope of Archa Mountain, [~51°26' N, 87°19' E], 1250 m a.s.l., 25 Jun. 2003, V. Sorokina leg. (SZMN); 2 ♂♂ 9 ♀♀, Altai Republic, Shebalino district, environs of Mukhorcherga village, Belog-Mukhorcherga Mountain, 51°20' N, 85°19' E, 1548–1854 m a.s.l., 22 Jul. 2011, V. Sorokina leg. (SZMN); 1 ♂, Altai Republic, Seminsky Pass, 51°03' N, 85°35' E, 1650 m a.s.l., 27–30 Jun. 2016, N. Vikhrev leg. (ZMUM); 2 ♂♂ 4 ♀♀, Altai Republic, Kosh-Agach district, upper part of Naryn-Gol River, 49°49' N, 89°32' E, 2520 m a.s.l., 15–19 Jul. 2009, V. Sorokina and T. Novgorodova leg. (SZMN); 5 ♂♂ 5 ♀♀, Altai Republic, Kosh-Agach district, south slope of Yuzhno-Chuiskiy range, valley of Tara River, 49°39' N, 88°13' E, 2200–2600 m a.s.l., 1–4 Jul. 2006, 11 Jul. 2009, V. Sorokina leg. (SZMN); 15 ♂♂ 1 ♀, Altai Republic, Kosh-Agach district, Ukok plateau, Rodonovyi spring, upper part of Zhumaly river, 49°27' N, 88°03' E, 2410 m a.s.l., 24 Jun. 2005, 4 Jul. 2005, 23–24 Jul. 2006, V. Sorokina, A. Barkalov and T. Novgorodova leg. (SZMN); 32 ♂♂ 7 ♀♀, Altai Republic, Kosh-Agach district, 6–8 km NW Maitobe Mountain, 49°34' N, 87°43' E, 2400–2800 m a.s.l., 6–9 Jul. 2006, V. Sorokina and T. Novgorodova leg. (SZMN); 18 ♂♂ 7 ♀♀, Altai Republic, Kosh-Agach district, environs of Muzdy-Bulak lake, 49°17' N, 87°39' E, 2400–2900 m a.s.l., 26 Jun.–2 Jul. 2005, V. Sorokina and A. Barkalov leg. (SZMN); 2 ♂♂ 6 ♀♀, Altai Republic, Kosh-Agach district, environs of Kal'dzhin-Kul'-Bas lake, 49°19' N, 87°26' E, 2400–2600 m a.s.l., 17–21 Jul. 2006, V. Sorokina and T. Novgorodova leg. (SZMN); 11 ♂♂ 5 ♀♀, Altai Republic, Kosh-Agach district, Severo-Chuyskiy Range, valley of Aktru river, 50°06' N, 87°48' E, 2014–2064 m a.s.l., 21 Jul. 2013, V. Sorokina and T. Novgorodova leg. (SZMN); 1 ♀, Altai Republic, Kosh-Agach district, south slope of Kurayskiy mountain ridge, 50°18' N, 87°51' E, 2736 m a.s.l., 19 Jul. 2013, V. Sorokina leg. (SZMN); 1 ♂, Altai Republic, Ulagan district, upper part of Ioldu river, 50°50' N, 88°57' E, 2000 m a.s.l., 18 Jun. 2005, V. Sorokina leg. (SZMN); 10 ♂♂ 3 ♀♀, Altai Republic, Ulagan district, environs of Dzhulukul' lake, 50°28' N, 89°46' E, 2200 m a.s.l., 21 Jul. 2007, V. Sorokina leg. (SZMN); 1 ♀, Altai Republic, Ongudai district, Terektinskiy range, upper part of Bolshoi Yaloman river, 50°28' N, 86°19' E, 2200–2300 m a.s.l., 3 Aug. 2007, V. Sorokina leg. (SZMN); 4 ♂♂ 6 ♀♀, Altai Republic, Ust'-Koksa district, Nizhne-Multinskoe lake, 50°01' N, 85°49' E, 1633 m a.s.l., 13 Jul. 2013, V. Sorokina leg. (SZMN); 4 ♂♂ 2 ♀♀, Altai Republic, Ust'-Koksa district, 47 km W Ust'-Koksa, Krasnaya Mountain, 50°04' N, 85°13' E, 2000–2100 m a.s.l., 16 Jul. 2015, V. Sorokina leg. (SZMN); 3 ♂♂ 6 ♀♀, Chukotka AO, Beringovskiy district, 40 km SSW of Beringovskiy, 62°43' N, 178°55' E, 22, 23, 26, 30 Jul. 2012,

A. Stekolshchikov leg. (SZMN); 8 ♂♂ 8 ♀♀, Chukotka AO, lower part of the Anadyr river, 64°26' N, 175°53' E, 5 m a.s.l., 18, 24–25 Jul. 2013, A. Barkalov leg. (SZMN); 7 ♂♂ 2 ♀♀, Chukotka AO, 30 km NW Krasneno, Anadyr river, 64°46' N, 174°8' E, 6–19 Jul. 2014, A. Barkalov and V. Zinchenko leg. (SZMN); 1 ♂, “Chita region” [Zabaykalsky Krai], bank of Bukukunskoe lake, 49°42' N, 111°5' E, 1880 m a.s.l., 3 Aug. 1991, V. Pekin leg. (SZMN); 1 ♂, “Chita region” [Zabaykalsky Krai], Sokhondo Nature Reserve, 49°41' N, 111°5' E, 1350–1400 m a.s.l., 2 Jun. 1991, V. Pekin and D. Logunov leg. (SZMN); 7 ♂, Krasnoyarsk Kray, environs of Norilsk, Talnakh, 69°21' N, 87°45' E, birch forest, 10, 12 Jul. 1967, K. Gorodkov leg. (ZISP); 4 ♂ 1 ♀, same data as for preceding except: in village, 22 Jul. 1973; 1 ♂, Krasnoyarsk Kray, upper part of Pura river [left tributary of Pyasina river], 10 km NW of Berasmu lake [71°51' N, 87°30' E], willow in bank of river, 21 Jul. 1967, K. Gorodkov leg. (ZISP); 6 ♂ 1 ♀, Krasnoyarsk Kray, Taymyr Peninsula, 90 km S of Tareya village, Pyasina river, [72°25' N, 90°56' E], willow in bank of river, 23 Jul. 1967, K. Gorodkov leg. (ZISP); 4 ♂♂ 1 ♀, Krasnoyarsk Kray, Taymyr Peninsula, 12.5 km S of Dikson, 73°24' N, 80°39' E, 26 Jul. 2012, A. Barkalov leg. (SZMN); 8 ♂♂ 4 ♀, Krasnoyarsk Kray, Taymyr Peninsula, 20 km NW of Khatanga, Ary-Mas station, 72°03' N, 101°56' E, 14 m a.s.l., 10–22 Jul. 2010, A. Barkalov leg. (SZMN); 2 ♀♀, Krasnoyarsk Kray, Taymyr Peninsula, 114 km S of Khatanga, Kotuy river, 71°02' N, 103°00' E, 2–5 Jul. 2010, A. Barkalov leg. (SZMN); 2 ♂♂, Magadan region, Ust'-Omchug, 61°08' N, 149°39' E, 29 Jun. 1971, K. Gorodkov leg. (ZISP); 4 ♂♂, same data as for preceding except: larch woodlands, 1 Jul. 1971; 2 ♀♀, same data as for preceding except: bog, 1 Aug. 1971, K. Gorodkov and Chelnokov leg. (ZISP); 1 ♂, Magadan region, 20 km S of Gizhiga village, “Chaibukha aerodrome”, [~61°50' N, 160°32' E], 2 Sep. 1987, K. Gorodkov leg. (ZISP); 1 ♂, Magadan region, environs of Magadan, Marchekan, 59°32' N, 150°47' E, 28 Aug. 1969, K. Gorodkov leg. (ZISP); 1 ♂ 1 ♀, Magadan region, the motorway R-504, Yablonevyy Pass, 120 km NNE of Magadan, 60°35' N, 151°32' E, 900 m a.s.l., 17 Jul. 2014, N. Vikhrev leg. (ZMUM); 6 ♂♂, Nenets AO, lower part of the Pechora river, 68°20' N, 53°18' E, 11 Jul. 2008, A. Ozerov leg. (2 ♂♂ SZMN, 4 ♂♂ ZMUM); 2 ♀♀, Yamalo-Nenets AO, 73 km NE of Labytnangi, Longot'egan river, 67°19' N, 66°43' E, 175 m a.s.l., 6, 22 Jul. 2015, A. Barkalov and V. Zinchenko leg. (SZMN).

SWEDEN: 3 ♂♂ 1 ♀, Abisko, Nuolia Mountain, Ridonjira, heath, 29 Jul. 1995, A. Pont leg. (SZMN).

USA: 1 ♂ 1 ♀, Alaska, Aleutian Islands, Chowiet Island, 56°02' N, 156°44' W, 107 m a.s.l., sweep in grass meadow, 10 Jul. 2009, A. Renaud leg. (SZMN).

Distribution

Holarctic Region; in the Palaearctic, from Greenland and Northern Europe to the Russian Far East. Arcto-alpine species.

Spilogona contractifrons (Zetterstedt, 1838)

Fig. 1A, D, G, J

Type locality

SWEDEN: “Lapland”.

Material examined

CANADA: 1 ♂, Manitoba, Goose Creek Rd., Pump House, 58°38' N, 94°14' W, Malaise trap, 17–21 Jul. 2007, A. Renaud leg. (SZMN); 1 ♂, Manitoba, 4 km S of Churchill, Beach Bay, 58°45' N, 94°08' W, sweep in vegetation, 24 Jul. 2007, A. Renaud leg. (SZMN); 1 ♀, Manitoba, Churchill, 12 km E on Launch road, 58°45' N, 93°59' W, Malaise trap, 3–7 Jul. 2007, A. Renaud leg. (SZMN); 1 ♀, Manitoba, 24 km E of Churchill, Ramsay Creek, 58°44' N, 93°47' W, Malaise trap, 25–29 Jul. 2007, A. Renaud leg. (SZMN).

SOROKINA V.S. & SHAIKEVICH E.V., Morphological and molecular analysis of *Spilogona*

RUSSIA: 1 ♂, Altai Republic, Kosh-Agach district, Ukok plateau, environs of Kal'dzhin-Kul'-Bas lake, 49°19' N, 87°26' E, 2450 m a.s.l., 21 Jul. 2006, V. Sorokina leg. (SZMN); 2 ♂♂, Altai Republic, Kosh-Agach district, Rodonovyi spring, upper part of Zhumaly river, 49°27' N, 88°03' E, 2410 m a.s.l., 4 Jul. 2005, 23–24 Jul. 2006, T. Novgorodova and V. Sorokina leg. (SZMN); 1 ♂, Altai Republic, Kosh-Agach district, Severo-Chuyskyi Ridge, valley of Aktru River, 50°06' N, 87°48' E, 2064 m a.s.l., 21 Jul. 2013, V. Sorokina leg. (SZMN); 2 ♀♀, "Chita region" [Zabaykalsky Krai], 6 km N, of Teli Mountain, Toreyskie lakes, 50°06' N, 115°41' E, steppe, 6 Jul. 2011, V. Dubatolov leg. (SZMN); 1 ♂, Kurgan region, environs of Kurgan, 55°30' N, 66°19' E, sweep in grass meadow, 17 Jun. 2015, V. Sorokina leg. (SZMN); 1 ♂, Magadan region, Ust'-Omchug, 61°08' N, 149°39' E, 1 Jul. 1971, Gorodkov leg. (ZISP); 1 ♀ 1 ♂, Magadan region, 20 km S of Gizhiga village, "Chaibukha aerodrome" [~61°50' N, 160°32' E], valley of Chaibukha River, 7 km from mouth of river, 2 Sep. 1987, Gorodkov leg. (ZISP); 1 ♀, Magadan region, Aborigen field station, 61°20' N, 149°23' E, 500 m a.s.l., 5–10 Aug. 1990, M. Wood leg. (SZMN); 10 ♂♂ 3 ♀♀, Magadan region, environs of Sokol village, 59°54' N, 150°45' E, 11–19 Jul. 2014, N. Vikhrev leg. (1 ♀ 1 ♂ in SZMN, rest in ZMUM); 1 ♂, Magadan region, 92 km NE of Magadan, valley of Donyshko river, 60°20' N, 151°23' E, 570 m a.s.l., 17 Jul. 2014, N. Vikhrev leg. (ZMUM); 1 ♂, Magadan region, Koni Peninsula, environs of Cape Ploskyi cordon, valley of Khindzha river, 59°09' N, 151°38' E, 29 Jun. 2016, N. Tridrikh leg. (SZMN); 2 ♀♀, same data as for preceding except: birch copse on the slope, 22 Aug. 2017; 1 ♀, Magadan region, Koni Peninsula, Cape Skalistyi, valley of Skalistaya river, 59°07' N, 151°23' E, yellow plate traps, 23 Aug. 2017, N. Tridrikh leg. (SZMN); 1 ♂ 2 ♀♀, Nenets AO, Khaypudyrskaya Bay, "Chuba", 68°16' N, 59°57' E, yellow plate trap, meadow, 6–16 Aug. 2015, O. Makarova and Bizin leg. (SZMN); 2 ♂♂ 2 ♀♀, Nenets AO, Pakhancheskaya Bay, 68°29' N, 57°26' E, yellow plate trap, tundra, 26 Jul.–5 Aug. 2015, O. Makarova and Bizin leg. (SZMN); 1 ♂ 1 ♀, Nenets AO, Bolvanskaya Bay, 68°05' N, 54°47' E, yellow plate trap, anthropogenic meadow, 18–25 Jul. 2015, O. Makarova and Bizin leg. (SZMN); 1 ♂ 4 ♀♀, Novosibirsk region, Zdvinsk district, environs of Shirokaya Kurya village [54°34' N, 78°08' E], field station, 8 Jul. 2002, Zavarukhina leg. (SZMN).

SWEDEN: 1 ♂ 1 ♀, "Vällistefjäll" [Jämtland, Välliste Mountain, 63°16' N, 13°07' E, 700–800 m a.s.l.], forest zone, 23 Jul. 1980, A. Pont leg. (SZMN).

Distribution

Holarctic Region; in the Palaearctic, from Northern Europe to the Russian Far East. Arcto-boreal-mountain species.

Spilogona orthosurstyla Xue & Tian, 1988

Fig. 2A–C

Type locality

CHINA: Hebei.

Material examined

RUSSIA: 1 ♂, Altai Republic, Turochak district, 30 km SW of Iogach, 51°31' N, 87°23' E, 1600 m a.s.l., 21 Jun. 2003, V. Sorokina leg. (SZMN).

Distribution

East Palaearctic: China (Hebei, Shanxi), Russia (Altai Mts). East Asian mountain species.

Remark

In view of the distribution of other species of this group, we can assume that this species also lives in the arctic zone.

Table 2. Estimates of evolutionary divergence between studied species of *Spilogona* Schnabl, 1911. The number of base substitutions per site from averaging over all sequence pairs between groups are shown. The analysis involved 16 nucleotide sequences.

	1	2	3	4	5
1 <i>S. alticola</i> (Malloch, 1920)					
2 <i>S. contractifrons</i> (Zetterstedt, 1838)	0.006				
3 <i>S. arctica</i> (Zetterstedt, 1838)	0.018	0.018			
4 <i>S. imitatrix</i> (Malloch, 1921)	0.046	0.039	0.039		
5 <i>S. nitidicauda</i> (Schnabl, 1911)	0.057	0.051	0.053	0.016	
6 <i>S. platyfrons</i> Sorokina, 2018	0.055	0.049	0.049	0.012	0.004

Molecular-genetic analysis of ‘Spilogona contractifrons species-group’

Since in our collection there is only one specimen of *S. orthosurstyla* and it is not in good condition, the molecular analysis was performed only on the other three species of this group.

Genetic distances between species are shown in the ML tree (Fig. 3) and in Table 2. Regardless of the geographic origin of the specimens, all three studied species in this group have identical *COI* gene sequences within the species. At the same time, at least five nucleotide substitutions (single nucleotide polymorphisms (SNPs)) vary between *S. alticola* and *S. contractifrons*. *Spilogona arctica* differs from *S. alticola* with 13 SNP and from *S. contractifrons* with 12 SNP; one substitution results in amino acid change (M112L).

The genetic distance between *S. alticola* and *S. contractifrons* is 0.6%, and between each of these species and *S. arctica* is 1.8%. All three species formed a distinct cluster on the ML tree with high (98–99%) statistical support in which the associated taxa clustered together (Fig. 3).

The following results were obtained after comparing the sequences of the studied specimens with the annotated sequences in Genbank. *Spilogona arctica* from the Altai, Taimyr and from the Nenets AO is identical with *S. arctica* from Sweden. All of them are distinguished by a single substitution of C438T from *S. arctica* from Canada and Greenland ([Supplementary file](#)). *Spilogona contractifrons* from the Altai, Nenets AO, Kurgan and Naryan-Mar is identical in the *COI* gene with *S. contractifrons* from Norway and Sweden. But *S. alticola* from the Altai clustered with *S. contractifrons* from Canada (Fig. 3).

Taxonomic analysis of ‘Spilogona nitidicauda species-group’

Flies of this group of species can be distinguished as follows: scutum with 4 *post dc* setae, anepisternum with interspatial setae, facial edge not projecting beyond level of profrons, hind femur without *pv* setae, scutellum without downwardly-directed preapical setulae on upper border of declivities, abdomen flattened with large trapezoidal marks on tergite 3.

Four species of this group are currently recognised: *S. nitidicauda*, *S. hissarensis*, *S. platyfrons* and *S. imitatrix*. *Spilogona hissarensis* was studied only from the original type-series. The other species of this group were studied with material from the territory of Russia and from Hockett’s collection deposited in the Canadian National Collection, Ottawa, Canada.

Despite the difficulty in distinguishing the species of this group, there are several helpful diagnostic characters: the width of frons and parafacial, presence of pollen on the prementum, the colour of the scutum and abdomen, the shape of male terminalia (Figs 4–5). Additional diagnostic characters are given in Table 3.




Fig. 3. Maximum Likelihood phylogenetic tree shows evolutionary relationships of species of *Spilogona* Schnabl, 1911.




Fig. 4. Males of *Spilogona* Schnabl, 1911. **A–C.** Head, lateral view. **A.** *S. imitatrix* (Malloch, 1921). **B.** *S. nitidicauda* (Schnabl, 1911). **C.** *S. platyfrons* Sorokina, 2018. **D–F.** Frons, anterior view. **D.** *S. imitatrix*. **E.** *S. nitidicauda*. **F.** *S. platyfrons*. **G–I.** Abdomen, dorsal view. **G.** *S. imitatrix*. **H.** *S. nitidicauda*. **I.** *S. platyfrons*. Scale bars: 1 mm.




Fig. 5. Males of *Spilogona* Schnabl, 1911. **A–C.** Terminalia, lateral view. **A.** *S. imitatrix* (Malloch, 1921). **B.** *S. nitidicauda* (Schnabl, 1911). **C.** *S. platyfrons* Sorokina, 2018. **D–F.** Cercal plate, posterior view. **D.** *S. imitatrix*. **E.** *S. nitidicauda*. **F.** *S. platyfrons*. **G–I.** Sternite 5. **G.** *S. imitatrix*. **H.** *S. nitidicauda*. **I.** *S. platyfrons*. Scale bars: 0.25 mm.

Table 3. Diagnostic characters of the ‘*Spilogona nitidicauda* species-group’.

	<i>S. nitidicauda</i> (Schnabl, 1911)	<i>S. hissarensis</i> Hennig, 1959	<i>S. imitatrix</i> (Malloch, 1921)	<i>S. platyfrons</i> Sorokina, 2018
Males				
Frons	fronto-orbital plates separated by a distinct frontal vitta (Fig. 4E)	fronto-orbital plates separated by a very narrow frontal vitta	fronto-orbital plates touching in upper part or separated by a very narrow frontal vitta (Fig. 4D)	fronto-orbital plates touching for greater part of length, frons very narrow (Fig. 4F)
Distance between eye margins	as wide as distance across posterior ocelli	narrower than distance across posterior ocelli	narrower than distance across posterior ocelli	as wide as diameter of anterior ocellus
Premetum	shining	shining	shining	shining, with a patch of light dust centrally on underside
Notopleuron	bare	bare	with some hairs in upper part, but not around posterior seta	bare
Interspatial setae	one	two	one, two or three	one or two
Scutum (in frontal view)	anterior half of presutural region with grey dust, sometimes without distinct dust	presutural region without grey dust	anterior half of presutural region with grey dust, sometimes without distinct dust	whole presutural region with whitish-grey dust
Halteres	yellow	dark brown	yellow	yellow, sometimes brownish-yellow
Mid tibia	1 <i>ad</i> and sometimes <i>v</i> present	without <i>ad</i> and <i>v</i>	without <i>ad</i> and <i>v</i>	without <i>ad</i> and <i>v</i>
Mid femora	2–4 stout <i>pv-v</i>	5–6 stout <i>pv-v</i>	2–4 stout <i>pv-v</i>	5–6 stout <i>pv-v</i>
Hind femora (apical part)	3–4 <i>av</i>	5–6 <i>av</i>	usually 3–4 <i>av</i>	5–6 <i>av</i>
Abdomen	marks separated (Fig. 4H)	marks narrowly separated	usually marks of tergite 3 connected, sometimes narrowly separated (Fig. 4G)	marks narrowly separated (Fig. 4I)
Sternite 5	processes without dense setulae (Fig. 5H)	processes without dense setulae	processes with dense setulae (Fig. 5G)	processes without dense setulae (Fig. 5I)
Cercal plate	oval in shape (Fig. 5E)	triangular in shape	Trapezoid in shape (Fig. 5D)	triangular in shape (Fig. 5F)
Females				
Premetum	shining	unknown	shining	shining, with patch of light dust centrally on underside
Notopleuron	bare		with some hairs in upper part, but not around posterior seta	usually bare, sometimes with several hairs in upper part
Interspatial setae	one		usually two or one	usually one or two
Mid femora	1 stout <i>pv-v</i> in basal ¼ and 1 extra hair-like seta directly at base		1 long stout <i>pv-v</i> in basal ¼ and 1–2 extra stout or hair-like setae directly at base	row of 3–5 hair-like <i>pv-v</i> in basal 1/2 (one seta stouter)

SOROKINA V.S. & SHAIKEVICH E.V., Morphological and molecular analysis of *Spilogona*

The species of this group can be separated by the following couples:

Males

1. In frontal view, presutural region of scutum without grey dust; haltere dark brown*S. hissarensis* Hennig, 1959
 – In frontal view, presutural region of scutum with grey dust; haltere yellow or brownish-yellow 2
2. Fronto-orbital plates touching along the greater part of their length, narrower than diameter of anterior ocellus; distance between eye margins on upper part of frons about as wide as diameter of anterior ocellus; prementum shining but with a patch of light dust centrally on underside; presutural region mostly with whitish-grey dust; mid femur with 5–6 long and stout *pv-v* in basal half; hind femur with 5–6 *av* at apex*S. platyfrons* Sorokina, 2018
 – Fronto-orbital plates touching only in upper part or separated by a narrow frontal vitta; distance between eye margins on upper part of frons as wide as or narrower than the distance between posterior ocelli; prementum completely shining, without any dust; extreme anterior half of presutural region with grey dust; mid femur with 2–4 stout *pv-v* in basal half; hind femur with 3–4 *av* at apex 3
3. Notopleuron with some hairs in upper part, but not around posterior seta; fronto-orbital plates touching in upper part or separated by a very narrow frontal vitta; mid tibia without *ad* and *v*; dark marks on tergite 3 usually connected or narrowly separated; caudal margin of sternite 5 with dense setulae*S. imitatrix* (Malloch, 1921)
 – Notopleuron completely bare; fronto-orbital plates usually separated by a distinct frontal vitta, sometimes touching; mid tibia with 1 *ad* and sometimes a short *v* present; tergites with well separated marks; caudal margin of sternite 5 without dense setulae*S. nitidicauda* (Schnabl, 1911)

Females

1. Prementum of proboscis shining but with a patch of light dust centrally on underside; mid femur with a row of 3–5 hair-like *pv-v* in basal ½, sometimes one of them stouter*S. platyfrons* Sorokina, 2018
 – Prementum of proboscis completely shining; mid femur with 1 stout *pv-v* in basal ¼ and 1–2 additional stout or hair-like setae directly at base 2
2. Notopleuron completely bare; anepisternum with 1 interspatial seta; mid tibia usually with *ad*; mid femur with 1 stout *pv-v* in basal ¼ and 1 additional hair-like seta*S. nitidicauda* (Schnabl, 1911)
 – Notopleuron with some hairs in upper part; usually anepisternum with 2 interspatial setae, rarely with one; mid tibia without *ad*; mid femur with 1 long stout *pv-v* in basal ¼ and 1–2 additional stout or hair-like setae*S. imitatrix* (Malloch, 1921)

Spilogona hissarensis Hennig, 1959

Type locality

TAJIKISTAN: Varzob.

Material examined

Holotype

TAJIKISTAN: 1 ♂, “u. Kondara, 1100 m, d. Varzoba, Tadzh. 24.vii.38, leg. Gussakovskii” [Varzob village, ~8 km N of Dushanbe, 33°39' N, 68°47' E, 1100 m a.s.l., 24 Jul. 1938, Gussakovskii leg.] (ZISP).

Paratype

TAJIKISTAN: 2 ♂♂, “u. Kvak-Ruidasht, Gissar. khr. Tadzsh., 16.vi.38., Gussakovskii” [Gissar Mountains, Rudaki district, 16 Jun. 1938, Gussakovskii leg.] (ZISP).

Distribution

Central Palaearctic: Tajikistan.

Remark

Mountain species, but probably living in the arctic tundra too.

Spilogona imitatrix (Malloch, 1921)

Figs 4A, D, G, 5A, D, G

Type locality

CANADA: Labrador, Nain.

Material examined

CANADA: 2 ♀♀, Manitoba, 24 km E of Churchill, Pistol range, 58°45' N, 93°56' W, sweep in vegetation, 12 Jul. 2007, A. Renaud leg. (SZMN).

RUSSIA: 1 ♂, Altai Republic, Kosh-Agach district, upper part of Naryn-Gol River, 49°49' N, 89°32' E, 2520 m, 15–18 Jul. 2009, T. Novgorodova leg. (SZMN); 10 ♂♂ 6 ♀♀, Altai Republic, Kosh-Agach district, 7 km NW of Kuray, south slope of Kurayskii mountain ridge, 50°18' N, 87°51' E, 2588 m a.s.l., 17, 19 Jul. 2013, V. Sorokina leg. (SZMN); 5 ♂♂ 1 ♀, Altai Republic, Kosh-Agach district, Ukok plateau, Rodonovyi spring, upper part of Zhumaly River, 49°27' N, 88°03' E, 2410 m a.s.l., 24 Jun. 2005, V. Sorokina leg. (SZMN); 6 ♂♂ 2 ♀♀, Altai Republic, Kosh-Agach district, 4 km NW of Kal'dzhin-Kul'-Bas lake, 49°19' N, 87°26' E, 2600 m a.s.l., 17–21 Jul. 2006, V. Sorokina leg. (SZMN); 1 ♂ 1 ♀, Altai Republic, Kosh-Agach district, 6–8 km NW of Maitobe Mountain, 49°34' N, 87°43' E, 2400–2700 m a.s.l., 8–10 Jul. 2006, T. Novgorodova leg. (SZMN); 6 ♂♂ 1 ♀, Altai Republic, Ongudai district, Terektinskiy range, upper part of Bolshoi Yaloman River, 50°28' N, 86°19' E, 2200–2300 m a.s.l., 3 Jul. 2007, V. Sorokina leg. (SZMN); 3 ♂♂ 5 ♀♀, Altai Republic, Ulagan district, environs of Dzhulukul' lake, 50°28' N, 89°46' E, 2200 m a.s.l., 21 Jul. 2007, V. Sorokina leg. (SZMN); 44 ♂♂ 23 ♀♀, Altai Republic, Ulagan district, Shapshal'skiy mountain range, 50°32' N, 89°48' E, 2550–2878 m a.s.l., on flowers of *Schultzia crinita* Spreng., 23–25 Jul. 2007, V. Sorokina and A. Barkalov leg. (SZMN); 1 ♂, Altai Republic, Ulagan district, Kurayskiy range, 50°20' N, 87°45' E, 2500–2700 m a.s.l., 3 Jul. 2008, A. Barkalov leg. (SZMN).

Distribution

Holarctic Region; in the Palaearctic only from southern Siberia. Arcto-alpine species.

Spilogona nitidicauda (Schnabl, 1911)

Figs 4B, E, H, 5B, E, H

Type locality

RUSSIA: Yamalo-Nenetsk AO.

SOROKINA V.S. & SHAIKEVICH E.V., Morphological and molecular analysis of *Spilogona***Material examined****Syntype**

RUSSIA: 5 ♂♂ 5 ♀♀, “Karskaja tundra, s. Tobol’s. gub., 21.vii.09, Zaizev” [Yamalo-Nenetskiy AO, east slope of Polar Ural, mouth of the River Kara, ~69°02’N, 64°35’E, 21 Jul. 1909, Zaizev leg.]. 1 ♂, holotype of *Limnophora (Spilogona) nitidicauda* var. *depressa* Schnabl, same data as for preceding (ZISP).

Other material examined

RUSSIA: 9 ♂♂ 2 ♀♀, Altai Republic, Kosh-Agach district, Ukok plateau, environs of Muzdy-Bulak lake, 49°17’ N, 87°39’ E, 2450 m a.s.l., 26 Jun. 2005, 1–2 Jul. 2005, A. Barkalov and V. Sorokina leg. (SZMN); 24 ♂♂ 8 ♀♀, Altai Republic, Kosh-Agach district, Ukok plateau, environs of Kal’dzhin-Kul’-Bas lake, 49°19’ N, 87°26’ E, 2400–2600 m a.s.l., 17–21 Jul. 2006, V. Sorokina and T. Novgorodova leg. (SZMN); 4 ♂♂ 1 ♀, Altai Republic, Kosh-Agach district, Ukok plateau, 8 km NW of Maitobe Mountain, 49°34’ N, 87°43’ E, 2400–2500 m a.s.l., 8–10 Jul. 2006, V. Sorokina leg. (SZMN); 12 ♂♂, Altai Republic, Kosh-Agach district, Ukok plateau, Rodonovyi spring, upper part of Zhumaly River, 49°27’ N, 88°03’ E, 2410 m a.s.l., 24 Jun. 2005, 4 Jul. 2005, 23–24 Jul. 2006, V. Sorokina and T. Novgorodova leg. (SZMN); 1 ♂ 1 ♀, Altai Republic, Kosh-Agach district, upper part of Naryn-Gol River, 49°49’ N, 89°32’ E, 2520 m a.s.l., 15–19 Jul. 2009, V. Sorokina and T. Novgorodova leg. (SZMN); 1 ♂, Altai Republic, Kosh-Agach district, Severo-Chuyskiy Ridge, valley of Aktru River, 50°06’ N, 87°48’ E, 1865–2064 m a.s.l., 21 Jul. 2013, T. Novgorodova leg. (SZMN); 2 ♀♀, Altai Republic, Ongudai district, Terektinskiy range, upper part of Bolshoi Yaloman River, 50°28’ N, 86°19’ E, 2200–2300 m a.s.l., 3 Jul. 2007, V. Sorokina leg. (SZMN); 1 ♂, Altai Republic, Ulagan district, Kurayskiy range, 50°20’ N, 87°45’ E, 2500–2800 m a.s.l., 29–30 Jun. 2008, A. Barkalov leg. (SZMN); 5 ♂♂ 16 ♀♀, Altai Republic, Ulagan district, environs of Dzhulukul’ lake, 50°28’ N, 89°46’ E, 2200 m a.s.l., 21 Jul. 2007, V. Sorokina leg. (SZMN); 2 ♂♂ 5 ♀♀, Altai Republic, Ulagan district, Shapshal’skiy mountain range, upper part of Shuy River, 50°32’ N, 89°48’ E, 2550–2878 m a.s.l., 23–25 Jul. 2007, V. Sorokina leg. (SZMN); 1 ♂ 1 ♀, Altai Republic, Ust-Koksa district, 47 km W Ust-Koksa, Krasnaya Mountain, 50°04’ N, 85°13’ E, 2000–2100 m a.s.l., 16 Jul. 2015, V. Sorokina leg. (SZMN); 15 ♂♂ 3 ♀♀, Chukotka AO, Beringovskii district, 40 km SSW of Beringovskii, 62°43’ N, 178°55’ E, yellow plate trap, 22–23, 26 Jul. 2012, A. Stekolshchikov leg. (SZMN); 1 ♀, Chukotka AO, 30 km NW of Krasneno, Anadyr river, 64°46’ N, 174°8’ E, 15 Jul. 2014, A. Barkalov leg. (SZMN); 1 ♂, Krasnoyarsk Kray, Taymyr Peninsula, Kuria village, upper part of Pyasina river [70°16’ N, 88°30’ E], alder forest near village, 13 Jul. 1967, K. Gorodkov leg. (ZISP); 1 ♀, Krasnoyarsk Kray, Taymyr Peninsula, 114 km S of Khatanga, Kotuy river, 71°02’ N, 103°00’ E, 2–5 Jul. 2010, A. Barkalov leg. (SZMN); 1 ♂ 2 ♀♀, Nenets AO, Khaypudyrskaya Bay, “Chuba”, 68°16’ N, 59°57’ E, yellow plate trap, meadow, 6–16 Aug. 2015, O. Makarova and Bizin leg. (SZMN); 1 ♂ 1 ♀, Nenets AO, Bolvanskaya Bay, 68°05’ N, 54°47’ E, yellow plate trap, anthropogenic meadow, 18–25 Jul. 2015, O. Makarova and Bizin leg. (SZMN); 1 ♀, Nenets AO, Pakhancheskaya Bay, 68°29’ N, 57°26’ E, 26 Jul.–5 Aug. 2015, O. Makarova and Bizin leg. (SZMN); 2 ♂♂, Yamalo-Nenets AO, 73 km NE of Labytnangi, Longot’egan river, 67°19’ N, 66°43’ E, 175 m a.s.l., 2, 10 Jul. 2015, A. Barkalov leg. (SZMN).

SWEDEN: 1 ♂, Abisko, Nuolja, meadow below Topstation, 19 Jul. 1987, A. Pont leg. (SZMN).

Distribution

Holarctic Region; in the Palaearctic, from North Scandinavia, Russia and China. Arcto-alpine species.

***Spilogona platyfrons* Sorokina, 2018**
Figs 4C, F, I, 5C, F, I

Type locality

RUSSIA: Altai Mountains.

Material examined

Holotype

RUSSIA: 1 ♂, Altai Republic, Ulagan district, Shapshal'skiy mountain range, upper part of Shui River, ~50°33' N, 89°50' E, 2550–2900 m a.s.l., 25 Jul. 2007, V. Sorokina leg. (SZMN).

Paratypes

RUSSIA: 34 ♂♂ 21 ♀♀, Altai Republic, Ulagan district, Shapshal'skiy mountain range, upper part of Shui River, ~50°33' N, 89°50' E, 2550–2900 m a.s.l., 25 Jul. 2007, V. Sorokina leg. (2 males and 2 females in ZMUM & ZISP, the rest in SZMN); 1 ♀, Altai Republic, Kosh-Agach district, upper part of Naryn-Gol River, 49°49' N, 89°32' E, 2520 m a.s.l., 16–19 Jul. 2009, V. Sorokina leg. (SZMN); 1 ♂ 1 ♀, Altai Republic, Kosh-Agach district, Ukok plateau, environs of Muzdy-Bulak lake, 49°17' N, 87°39' E, 2400–2900 m a.s.l., 27 Jun., 1 Jul. 2005, V. Sorokina leg. (SZMN); 6 ♂♂, Altai Republic, Kosh-Agach district, Ukok plateau, 6–8 km NW of Maitobe Mountain, 49°34' N, 87°43' E, 2420–2550 m a.s.l., 6–10 Jul. 2006, V. Sorokina and T. Novgorodova leg. (SZMN); 6 ♂♂ 4 ♀♀, Altai Republic, Kosh-Agach district, Ukok plateau, environs of Kal'dzhin-Kul'-Bas lake, 49°19' N, 87°26' E, 2400–2600 m a.s.l., 18, 20–21 Jul. 2006, V. Sorokina and T. Novgorodova leg. (SZMN); 17 ♂♂ 3 ♀♀, Altai Republic, Kosh-Agach district, Ukok plateau, south slope of Yuzhno-Chuiskiy range, valley of Tara River, 49°39' N, 88°13' E, 2175–2200 m a.s.l., 1–4 Jul. 2006, 11 Jul. 2009, leg. V. Sorokina leg. (SZMN); 6 ♂♂ 1 ♀, Altai Republic, Kosh-Agach district, Ukok plateau, upper part of Zhumaly river, Rodonovyi spring, 49°16' N, 88°30' E, 2300 m a.s.l., 24 Jun. 2005, V. Sorokina leg. (SZMN); 6 ♂♂ 1 ♀, Altai Republic, Ongudai district, Terektinskiy range, upper part of Bolshoi Yaloman River, 2200–2300 m a.s.l., 3 Aug. 2007, V. Sorokina leg. (SZMN); 1 ♂, Altai Republic, Ulagan district, Kurayskiy range, 50°20' N, 87°45' E, 2500–2800 m a.s.l., 29–30 Jun. 2008, A. Barkalov leg. (SZMN); 13 ♂♂ 20 ♀♀, Altai Republic, Ulagan district, environs of Dzhulukul' lake, 50°28' N, 89°46' E, 2200 m a.s.l., 21–24 Jul. 2007, V. Sorokina leg. (SZMN); 18 ♂♂ 3 ♀♀, Altai Republic, Ust-Koksa district, 47 km W of Ust-Koksa, Krasnaya Mountain, 2000–2100 m a.s.l., 50°04' N, 85°13' E, 16 Jul. 2015, V. Sorokina leg. (SZMN); 6 ♂♂ 4 ♀♀, Tyva Republic, Morgun-Taiginskii kozhuun, environs of Mugur-Aksy, ~50°22' N, 90°25' E, 2000–2100 m a.s.l., 22 Jul. 1993, A. Barkalov leg. (SZMN).

Distribution

Palearctic Region: southern Siberia.

Remark

Mountain species, but probably living in the arctic tundra too.

Molecular-genetic analysis of 'Spilogona nitidicauda species-group'

Since only the old type specimens of *S. hissarensis* are known, the molecular analysis was performed on the other three species of this group.

DNA variability is higher within taxa of the '*Spilogona nitidicauda* species-group' in comparison with taxa of the '*Spilogona contractifrons* species-group'. Intraspecies DNA variability was found in *S. nitidicauda* and *S. platyfrons* from different localities. The isolate SN38 of *Spilogona nitidicauda* from the Altai (Muzdy-Bulak lake) differs from the isolates SN40 and SN41 of *S. nitidicauda* from

Chukotka AO (Beringovskii) by a synonymous substitution of C186T. Both isolates from Chukotka have identical DNA. The SP32 of *S. platyfrons* from the Shapshalsky ridge (Altai) differs from the SP33 of *S. platyfrons* from the Krasnaya Mountain (Altai) by substitutions of T636C ([Supplementary file](#)). *Spilogona nitidicauda* can be distinguished from *S. platyfrons* by 3–4 SNPs. *Spilogona imitatrix* differs from *S. platyfrons* with 6 SNP, and from *S. nitidicauda* with 7 SNP.

The genetic distance between *S. nitidicauda* and *S. platyfrons* is 0.4%, between *S. platyfrons* and *S. imitatrix* is 1.2%, and between *S. nitidicauda* and *S. imitatrix* is 1.6%. All three species formed distinct clusters on the ML tree, but the differences between *S. nitidicauda* and *S. platyfrons* are insignificant statistically.

The following results were obtained after comparing the sequences of the studied specimens with the annotated sequences in GenBank. Unfortunately, there were no DNA sequences of *S. nitidicauda* and *S. platyfrons* in GenBank nor in the BOLD database, so comparison with our results was impossible. But there were many DNA sequences of *S. imitatrix* from Canada.

Spilogona imitatrix from the Altai is identical with several individuals of *S. imitatrix* from Canada (KR615453 and KR622696). However, several individuals from GenBank identified as *S. imitatrix* have a very variable DNA, as can be seen on the ML tree (Fig. 3). Such a high polymorphism may be due to errors in morphological identification.

Discussion

The analysis of genetic distances based on the differences in the cytochrome oxidase I gene in members of the ‘*Spilogona contractifrons* species-group’ and the ‘*Spilogona nitidicauda* species-group’ confirmed that there are not only morphological differences but also genetic differences between closely related species.

Our results with the DNA analysis of the *COI* gene have demonstrated that *S. alticola*, *S. contractifrons* and *S. arctica* are not synonymous because the genetic differences between individuals of these species persist in geographically distant locations (Fig. 6). The DNA of *S. arctica* is identical in individuals from different regions of Russia, Sweden, Greenland and Canada. Individuals of *S. contractifrons* from Russia are identical with *S. contractifrons* from Sweden and Norway, and all of them form a distinct cluster of “European” or “Palearctic” *S. contractifrons*. It should be emphasised that the specimens of *S. arctica* and *S. contractifrons* from Russia were identical with specimens from the type localities (Sweden and Greenland) but show differences from specimens of these species from Canada.

In turn, specimens from Canada designated in GenBank as *S. contractifrons* were identical with specimens from the Altai identified by us as *S. alticola*. The latter species was described from specimens from USA (New Hampshire) and was later found in different areas of Northern Canada (Huckett 1965). The reasons for this result could be either the incorrect identification of the specimens deposited in GenBank, or a rejection of the existence of *S. alticola* species by Canadian researchers and a designation of such specimens as *S. contractifrons*.

In the ‘*Spilogona nitidicauda* species-group’ there are stable nucleotide differences between *S. nitidicauda* and *S. platyfrons* from different places and years of collection despite the morphological similarity. Although the genetic distance between *S. nitidicauda* and *S. platyfrons* is only 0.4%, *S. imitatrix* differs from them significantly. The DNA distance between *S. imitatrix* and *S. nitidicauda* is 1.6%, and that between *S. imitatrix* and *S. platyfrons* is 1.2% which demonstrates the separate species status of *S. imitatrix*. In Muscidae, ranges of 0.18% and 3.01% have been reported as averages for the means and maxima of *COI* intraspecific distances, respectively (Renaud *et al.* 2012). Limits of species with a

distance to the nearest neighbour of $< 2\%$ in 137 species of Muscidae were supported by morphological characters and confirmed that the minimum interspecific distance in Muscidae ranges well below 2% for many species (Renaud *et al.* 2012).

Spilogona imitatrix was described from specimens from Labrador (Canada) and was later found in different regions of Canada and USA (Alaska). Recently this species was found in the Palaearctic Region for the first time, here known only in the Altai Mountains (Sorokina 2018). The DNA of *S. imitatrix* from the Altai (Terektinsky ridge) was identical with the DNA of several individuals of *S. imitatrix* from the Northwest Territories (Canada). There is a rather large number of the sequences of this species




Fig. 6. Localities of the species of *Spilogona* Schnabl, 1911 used in the DNA analysis. Symbols denote the species, whilst the colour shows the same DNA sequences. A. '*S. contractifrons* species-group'. B. '*S. nitidicauda* species-group'.

from Canada in GenBank. Several DNA sequences of *S. imitatrix* from Canada were randomly taken for genetic matching with the specimen of this species from the Altai. It transpired that one sequence (KR628150) clustered with *S. platyfrons* from the Altai, whilst the other sequences of Canadian specimens (# HM389123, # HM389124) differed from all of them by six nucleotide substitutions (Fig. 3). Such differences between DNA sequences of one species can be explained by errors in the initial morphological identification. This has been proved by the work of the first author of this paper (VSS) with the Muscidae collection in the CNC, where the following species were found under the name *S. imitatrix*, and vice versa: *S. alticola*, *S. arctica*, *S. deflorata* (Holmgren, 1872), *S. incauta* (Huckett, 1932), *S. nitidicauda*, *S. trigonifera* (Zetterstedt, 1838), *S. sororcula* (Zetterstedt, 1845), *S. trilineata* (Huckett, 1932) and another *Spilogona* sp.

These misidentifications are not surprising, since most members of the genus *Spilogona* are difficult to distinguish, even for muscid specialists. Therefore, in addition to a genetic analysis it is important to have an idea of the most stable morphological characters in each group. According to our results with this genus, the following can be considered as the most stable morphological characters in *Spilogona*: presence of *pv/v* setae on mid tibia, presence of *pv* setae on hind femur, presence of dust on the prementum, shape of the sternite 5, presence interspatial seta(e) on anepisternum, number of *post dc* setae on the scutum (3 or 4), presence on the scutellum of downwardly-directed preapical setulae on upper border of declivities. The last character was proposed and used only by Huckett (1965) for the separation of species into groups in his key to the species of *Spilogona*, in addition to the other well-known characters. Other researchers (Hennig 1959a, 1959b, 1959c; Gregor et al. 2002) used well-established characters for identification, specifically the number of *post dc* setae on the scutum (3 or 4), the elongation of the face, and the presence of *pv* setae on hind femur. The character suggested by Huckett (downwardly-directed preapical setulae on the upper border of declivities of scutellum) is quite difficult to see, which is probably why it has not been generally used by the authors of the keys. However, in our opinion, this is one of the important characters for the identification of the species of *Spilogona*. To demonstrate this, several outgroup species (*S. aerea* (Fallén, 1825), *S. bifimbriata* Huckett, 1965, *S. baltica* (Ringdahl, 1918) and *S. tornensis* (Ringdahl, 1926)) were included in the genetic analysis. These species have the same characters (3 and 4 *post dc* setae, mid tibia without *v* seta, hind femur with and without *pv* setae, face not projecting) as in the species groups analysed above, but, unlike them, these species have scutellum with downwardly-directed preapical setulae on the upper border of declivities. The results of the genetic analysis showed a distinct distance between the cluster of the ‘*Spilogona contractifrons* species-group’ and the ‘*Spilogona nitidicauda* species-group’ on the one hand, the members of which have scutellum without downwardly-directed preapical setulae on the upper border of declivities, and on the other hand the cluster of species having these setulae on the scutellum (Fig. 3).

The members of the ‘*Spilogona contractifrons* species-group’ and the ‘*Spilogona nitidicauda* species-group’ thus have not only distinguishing morphological characters but also fixed substitutions in the DNA sequences. Since a low interspecific polymorphism is known in the Muscidae, the revealed genetic distances confirm the existence of separate species or subspecies in each of the groups studied.

Acknowledgments

We would especially like to thank Adrian Pont (Oxford, UK) for discussion of this manuscript and for checking the English. The research presented in this paper was supported by the Federal Fundamental Scientific Research Programme for 2013–2020, the project No. VI.51.1.5(AAAA–A16–116121410121–7) and the Russian Federation Budget Project No. AAAA–A16–116111610180–3.

References

- Collin J.E. 1930. A revision of the Greenland species of the anthomyid genus *Limnophora* sens. lat. (Diptera), with figures of the male genitalia of these and many other Palaearctic species. *Transactions of the Entomological Society of London* 78: 255–281. <https://doi.org/10.1111/j.1365-2311.1930.tb00387.x>
- Gregor F., Rozkošný R., Barták M. & Vaňhara J. 2002. The Muscidae (Diptera) of Central Europe. *Folia Facultatis Scientiarum Naturalium Universitatis Masarykianae Brunensis, Biologia* 107: 1–280.
- Haseyama K.L.F., Wiegmann B.M., Almeida E.A.B. & de Carvalho C.J.B. 2015. Say goodbye to tribes in the new house fly classification: a new molecular phylogenetic analysis and an updated biogeographical narrative for the Muscidae (Diptera). *Molecular Phylogenetics and Evolution* 89: 1–12. <https://doi.org/10.1016/j.ympev.2015.04.006>
- Hennig W. 1959a. Muscidae [Part, Lieferung 204]. In: Lindner E. (ed.) *Die Fliegen der palaearktischen Region* 63b: 233–288. Schweizerbart, Stuttgart.
- Hennig W. 1959b. Muscidae [Part, Lieferung 205]. In: Lindner E. (ed.) *Die Fliegen der palaearktischen Region* 63b: 289–336. Schweizerbart, Stuttgart.
- Hennig W. 1959c. Muscidae [Part, Lieferung 207]. In: Lindner E. (ed.) *Die Fliegen der palaearktischen Region* 63b: 337–384. Schweizerbart, Stuttgart.
- Huckett H.C. 1965. The Muscidae of Northern Canada, Alaska and Greenland (Diptera). *Memoirs of the Entomological Society of Canada* 42: 5–369. <https://doi.org/10.4039/entm9742fv>
- Kutty S.N., Pape T., Pont A.C., Wiegmann B.M. & Meier R. 2008. The Muscoidea (Diptera: Calyptratae) are paraphyletic: evidence from four mitochondrial and four nuclear genes. *Molecular Phylogenetics and Evolution* 49: 639–652. <https://doi.org/10.1016/j.ympev.2008.08.012>
- Kutty S.N., Pont A.C., Meier R. & Pape T. 2014. Complete tribal sampling reveals basal split in Muscidae (Diptera), confirms saprophagy as ancestral feeding mode, and reveals an evolutionary correlation between instar numbers and carnivory. *Molecular Phylogenetics and Evolution* 78: 349–364. <https://doi.org/10.1016/j.ympev.2014.05.027>
- Michelsen V. 2006. Annotated catalogue of the Anthomyiidae, Fanniidae, Muscidae and Scathophagidae (Diptera: Muscoidea) of Greenland. *Steenstrupia* 29: 105–126.
- Pont A.C. 1986. Family Muscidae. In: Soós Á. & Papp L. (eds) *Catalogue of Palaearctic Diptera* 11: 57–215. Akadémiai Kiadó, Budapest.
- Ratnasingham S. & Hebert P.D.N. 2007. BOLD: The Barcode of Life Data System (www.barcodinglife.org). *Molecular Ecology Notes* 7: 355–364. <https://doi.org/10.1111/j.1471-8286.2006.01678.x>
- Renaud A.K., Savage J. & Adamowicz S.J. 2012. DNA barcoding of Northern Nearctic Muscidae (Diptera) reveals high correspondence between morphological and molecular species limits. *BMC Ecology* 12 (1): 1–15. <https://doi.org/10.1186/1472-6785-12-24>
- Savage J., Wheeler T.A. & Wiegmann B.M. 2004. Phylogenetic analysis of the genus *Thricops* Rondani (Diptera: Muscidae) based on molecular and morphological characters. *Systematic Entomology* 29: 395–414. <https://doi.org/10.1111/j.0307-6970.2004.00252.x>
- Schuehli G.S.E., de Carvalho C.J.B. & Wiegmann B.M. 2004. Regarding the taxonomic status of *Ophyra* Robineau-Desvoidy (Diptera: Muscidae): a molecular approach. *Zootaxa* 712: 1–12.
- Schuehli G.S.E., de Carvalho C.J.B. & Wiegmann B.M. 2007. Molecular phylogenetics of the Muscidae (Diptera: Calyptratae): new ideas in a congruence context. *Invertebrate Systematics* 21: 263–278. <https://doi.org/10.1071/IS06026>

SOROKINA V.S. & SHAIKEVICH E.V., Morphological and molecular analysis of *Spilogona*

- Shaikovich E.V. 2007. PCR-RFLP of the *COI* gene reliably differentiates *Cx. pipiens*, *Cx. pipiens* form *molestus* and *Cx. torrentium* of the *pipiens* Complex. *European Mosquito Bulletin* 23: 25–30.
- Simon C., Frati F., Beckenbach A., Crespi B., Liu H. & Flook P. 1994. Evolution, weighting, and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved polymerase chain reaction primers. *Annals of the Entomological Society of America* 87 (6): 651–701. <https://doi.org/10.1093/aesa/87.6.651>
- Sorokina V.S. 2012a. Fauna of Muscidae (Diptera) of the Altai Mountains. *Proceedings of the Russian Entomological Society* 83 (1): 193–222. [In Russian, with English abstract.]
- Sorokina V.S. 2012b. The Muscidae (Diptera) of the Russian tundra zones. Report 1. *Caucasian Entomological Bulletin* 8 (2): 328–332. [In Russian, with English abstract.]
- Sorokina V.S. 2013a. Fauno-genetic connections of Muscoidea (Diptera) of the high altitude of the Altai Mts and the latitudinal tundra of Taymyr Peninsula. In: Babin V.G., Tabakaev U.V., Dolgovykh S.V., Yurkova N.A., Aleynikova V.N. & Bondarenko A.V., *Biodiversity, ecological issues of Gorny Altai and its neighboring regions: present, past and future*: 115–118. Russia, Gorno-Altai. [In Russian, with English summary.]
- Sorokina V.S. 2013b. Muscidae from above the tree-line in the Altai Mountains. In: Babin V.G., Tabakaev U.V., Dolgovykh S.V., Yurkova N.A., Aleynikova V.N. & Bondarenko A.V., *Biodiversity, ecological issues of Gorny Altai and its neighboring regions: present, past and future*: 118–122. Russia, Gorno-Altai. [In Russian, with English summary.]
- Sorokina V.S. 2017. The Muscoidea Flies (Diptera) of the northern territories of Russia. *Euroasian Entomological Journal* 16(1): 44–56. [In Russian, with English summary.]
- Sorokina V.S. 2018. Eleven new species of *Spilogona* Schnabl, 1911 (Diptera, Muscidae) from the Altai Mountains of Russia, with key to species. *Zootaxa* 4410 (2): 201–250. <https://doi.org/10.11646/zootaxa.4410.2.1>
- Sorokina V.S. & Khruleva O.A. 2012. Details of species composition and distribution of house-flies (Diptera, Muscidae) of the Wrangel Island, Russia. *Euroasian Entomological Journal* 11: 553–564. [In Russian with English abstract.]
- Tamura K., Stecher G., Peterson D., Filipowski A. & Kumar S. 2013. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Molecular Biology and Evolution* 30 (12): 2725–2729. <https://doi.org/10.1093/molbev/mst197>
- Werner D. & Pont A.C. 2006a. The feeding and reproductive behavior of the Limnophorini (Diptera: Muscidae). *Studia Dipterologica, Supplement, Halle/Saale* 14: 79–114.
- Werner D. & Pont A.C. 2006b. New results on Diptera predators in the blackfly plague areas of Central Europe and the Caucasus. *Acta Entomologica Serbica, Supplement*: 131–140.

European Journal of Taxonomy 484: 1–26 (2018)

Manuscript received: 13 September 2018

Manuscript accepted: 15 October 2018

Published on: 11 December 2018

Topic editor: Gavin Broad

Desk editor: Alejandro Quintanar

Printed versions of all papers are also deposited in the libraries of the institutes that are members of the *EJT* consortium: Muséum national d'Histoire naturelle, Paris, France; Meise Botanic Garden, Belgium; Royal Museum for Central Africa, Tervuren, Belgium; Natural History Museum, London, United Kingdom; Royal Belgian Institute of Natural Sciences, Brussels, Belgium; Natural History Museum of Denmark, Copenhagen, Denmark; Naturalis Biodiversity Center, Leiden, the Netherlands; Museo Nacional de Ciencias Naturales-CSIC, Madrid, Spain; Real Jardín Botánico de Madrid CSIC, Madrid, Spain; Zoological Research Museum Alexander Koenig, Bonn, Germany.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [European Journal of Taxonomy](#)

Jahr/Year: 2018

Band/Volume: [0484](#)

Autor(en)/Author(s): Sorokina Vera S., Shaikevich Elena V.

Artikel/Article: [The identification of the species of the 'Spilogona contractifrons species-group' and the 'Spilogona nitidicauda species-group' \(Diptera, Muscidae\) based on morphological and molecular analysis 1-26](#)