



## Research article

[urn:lsid:zoobank.org:pub:5B9DC474-1F26-4732-B7B8-32C285FE0099](https://zoobank.org/pub:5B9DC474-1F26-4732-B7B8-32C285FE0099)

# Taxonomic revision of the subterranean genus *Virpazaria* Gittenberger, 1969 (Gastropoda, Spelaeodiscidae)

Zoltán FEHÉR<sup>1,\*</sup>, Tamás DELI<sup>2</sup>, Zoltán P. ERŐSS<sup>3</sup> & Romilda LIKA<sup>4</sup>

<sup>1</sup> Department of Zoology, Hungarian Natural History Museum,  
Baross u. 13, Budapest, H-1088, Hungary.

<sup>2</sup> Móricz Zsigmond u. 2, Gyomaendrőd, H-5500, Hungary.

<sup>3</sup> Bem u. 36., Budapest, H-1151, Hungary.

<sup>4</sup> Department of Biology, Faculty of Natural Sciences, Tirana University,  
Bulevardi Zogu i Parë, Tiranë, AL-1001, Albania.

\* Corresponding author: [feher.zoltan@nhmus.hu](mailto:feher.zoltan@nhmus.hu)

<sup>2</sup> Email: [virpazaria@gmail.com](mailto:virpazaria@gmail.com)

<sup>3</sup> Email: [eross peter@hotmail.com](mailto:eross peter@hotmail.com)

<sup>4</sup> Email: [romildalika@gmail.com](mailto:romildalika@gmail.com)

<sup>1</sup> [urn:lsid:zoobank.org:author:E801EC76-8B1E-450B-993E-BBBE57C00EA9](https://zoobank.org/author:E801EC76-8B1E-450B-993E-BBBE57C00EA9)

<sup>2</sup> [urn:lsid:zoobank.org:author:A33DCDF3-6293-4B34-B1DD-330211905D74](https://zoobank.org/author:A33DCDF3-6293-4B34-B1DD-330211905D74)

<sup>3</sup> [urn:lsid:zoobank.org:author:0B2C74A7-0001-4E25-984C-88CB0C705313](https://zoobank.org/author:0B2C74A7-0001-4E25-984C-88CB0C705313)

<sup>4</sup> [urn:lsid:zoobank.org:author:9662552B-B203-43FF-B427-4330080E1CBA](https://zoobank.org/author:9662552B-B203-43FF-B427-4330080E1CBA)

**Abstract.** *Virpazaria* Gittenberger, 1969 is distributed in the Balkan Peninsula (Albania, Montenegro and Croatia) and inhabits the shallow subterranean habitat (MSS) on limestone base rock. Reviewing historical and recently collected material, two species, *Virpazaria* (*Virpazaria*) *gittenbergeri* Fehér & Erőss sp. nov. and *Virpazaria* (*Virpazaria*) *pesici* Fehér & Deli sp. nov., are introduced as new to science. The conservation status of the new species are assessed using IUCN criteria. Two taxa, *Virpazaria* (*Virpazaria*) *pageti alexanderi* Reischütz & Subai, 2012 and *Virpazaria* (*Aemiliella*) *ripkeni pastorpueri* Reischütz *et al.*, 2011, are synonymized with their nominate subspecies. Some new distribution records, as well as geological and geomorphological data about the known locations for *Virpazaria*, are presented.

**Keywords.** Balkan, shallow subterranean habitat (MSS), new taxa, microendemic species, limestone habitat.

Fehér Z., Deli T., Erőss Z.P. & Lika R. 2019. Taxonomic revision of the subterranean genus *Virpazaria* Gittenberger, 1969 (Gastropoda, Spelaeodiscidae). *European Journal of Taxonomy* 558: 1–25.  
<https://doi.org/10.5852/ejt.2019.558>

## Introduction

*Virpazaria* Gittenberger, 1969 is a subterranean land snail genus occurring around the transboundary Lake Skadar (in Albania and Montenegro) as well as in the northern part of Kotor Bay (in Montenegro)

and Croatia). Previously, *Virpazaria* was believed to be an obligate troglobiont, but later findings have demonstrated that – similarly to many other land snail genera in the Balkans such as *Agardhiella* Hesse, 1923, *Balkanodiscus* Riedel & Urbański, 1964, *Gyalina* Andreae, 1902, *Klemmia* Gittenberger, 1969, *Pholeoteras* Sturany, 1904, *Sciocochlea* C.R. Boettger, 1935, *Spelaeodiscus* Brusina, 1886, *Troglomitrea* Negrea & Riedel, 1968, *Tsoukatosia* Gittenberger, 2000 and many species in *Platyla* Moquin-Tandon, 1856 or *Vitrea* Fitzinger, 1833 – it rather prefers superficial underground compartments (also known as the ‘Mesovoid Shallow Substratum’ or ‘Milieu Souterrain Superficiel’, widely abbreviated as MSS; see, e.g., Juberthie *et al.* 1980, 1981; Camacho 1992; Culver & Pipan 2014; Mammola *et al.* 2016). It is still disputed whether such a bedrock MSS habitat is characteristically different from caves or it can be considered as just the extension of the cave system, and therefore whether its fauna is composed of troglobionts of wider ecological tolerance or there are certain specialized elements exclusive of MSS (see Mammola *et al.* 2016 for a review). Either way, caves are interconnected with the bedrock’s crevice network, giving a reasonable explanation for occasional cave records of this genus.

Extensive field surveys applying the ‘scratch & flotate’ sampling method (Páll-Gergely *et al.* 2018) significantly increased the number of known populations and the amount of available material of subterranean gastropods in the past decade. In this paper we revise the genus *Virpazaria*, describe new taxa, present new distribution records and try to draw taxonomy-related conclusions. As little is known about the ecology of MSS-dwelling gastropods (Arndt & Subai 2013; Rendoš *et al.* 2014) and practically nothing about their habitat preferences in terms of the type, age and morphology of bedrock, we present data about the geology and geomorphology of the sampled habitats.

## Material and methods

The new samples were collected between 2015 and 2018. Sampling was done using the ‘scratch & flotate’ approach. First, we looked for superficial fissures, cracks and holes in limestone cliffs from where it was possible to scratch out fine granulate material by applying long and narrow handrakes. Shells were then separated by flotation in water.

For non-type material listed in the ‘Material examined’ sections, the number of specimens in each lot is not generally specified, as there is often uncertainty regarding how the specimens from a collecting event have been distributed between official institutions and the private collections of the collectors.

Albanian and former Yugoslavian geological maps (MMKS 1983; Mikinčić 1953) were used to identify the types and ages of the bedrock at the georeferenced sites.

Digital pictures were made using a Nikon DS-Ri2 camera attached to a Nikon SMZ25 stereo microscope and then combined into focus stack images with Nikon’s NIS software. ImageJ 1.49 scientific image analyzing software (Schneider *et al.* 2012) was used for measurements. We measured shell height, shell width, aperture height and aperture width (see Páll-Gergely *et al.* 2018: fig. 1), and counted the number of ribs on the last whorl, as well as the number of whorls ( $\pm 0.25$ ) according to Kerney & Cameron (1979: 13).

## Abbreviations used:

|        |   |                                                     |
|--------|---|-----------------------------------------------------|
| CNHM   | = | Croatian Natural History Museum, Zagreb, Croatia    |
| DT     | = | Collection Tamás Deli, Gyomaendrőd, Hungary         |
| ER     | = | Collection Zoltán Péter Erőss, Budapest, Hungary    |
| HNHM   | = | Hungarian Natural History Museum, Budapest, Hungary |
| NHMW   | = | Naturhistorisches Museum Wien, Vienna, Austria      |
| NHMW-E | = | Edlauer Collection in NHMW                          |

NHMW-K = Klemm Collection in NHMW

NMBE = Natural History Museum of Bern, Bern, Switzerland

RMNH = National Museum of Natural History, Leiden, the Netherlands

SMF = Senckenberg Forschungsinstitut und Naturmuseum, Frankfurt am Main, Germany

ad. = adult

fr. = fragment

H = shell height

H<sub>a</sub> = aperture height

W = shell width

W<sub>a</sub> = aperture width

## Results

Class Gastropoda Cuvier, 1795

Subclass Heterobranchia Burmeister, 1837

Order Stylommatophora A. Schmidt, 1855

Superfamily Pupilloidea W. Turton, 1831

Family Spelaeodiscidae Steenberg, 1925

Genus *Virpazaria* Gittenberger, 1969

### Type species

*Virpazaria adrianae* Gittenberger, 1969, by original designation.

### Diagnosis

Shell discoid, dextral, ribbed, colourless (weathered specimens are opaque white), peristome attached, thickened, usually with knob- or tooth-like prominences on palatal and/or basal sides. Peristome insertions are connected; this parietal connection varies from a swollen bulge to a lamelliform flap; in some species this flap might cover a significant part of the aperture in frontal view. Aperture somewhat narrow and moon-shaped.

### Remarks

According to Gittenberger (1969, 1975) as well as Reischütz & Reischütz (2009), *Virpazaria* is distinguished from the conchologically similar genus *Spelaeodiscus* by the continuous peristome, i.e., the presence of a parietal bulge or flap that connects the peristome insertions. *Spelaeodiscus virpazarioides* Páll-Gergely & Fehér, 2018 also possesses a callous parietal structure, but it is much smaller in proportion to the size of the whole aperture. Also, the narrower, crescent-shaped aperture is a good feature to tell *Virpazaria* apart from related genera.

Due to their hidden life, in most of the cases only empty shells are found. So far, there is one reported case when *Virpazaria*, more specifically two individuals of *V. deelemanorum*, were found alive (Gittenberger 1975). Having no pigmented eyes, they are presumably blind, as is usual for subterranean animals. Comparing their genitals to those of *Spelaeodiscus*, *Klemmia* and *Aspasita* Westerlund, 1889 (Bole 1965; Hudec 1965; Gittenberger 1975; Subai & Dedov 2008), one might discover differences in terms of proportional lengths of certain reproductive organs, and there are certain structures that are called differently by different authors (e.g., the same short protrusion at the border of penis and epiphallus is referred to as flagellum by Gittenberger (1975) and penial caecum by Schileyko (1998)). However, the overall arrangement of the genitalia is similar in these four genera, and no genus-specific differences could be found on the basis of the available literature.

# A key to species of *Virpazaria*

1. Two lamellae, one palatal and one basal, reach deeply (more than  $\frac{1}{10}$  of whorl) into the aperture ...  
.....*V. (Aemiliella) ripkeni* Gittenberger, 1969  
– No deeply reaching lamellae in the aperture, just knob(s) or on elongated thickening parallel with  
the peristome .....2
2. In frontal view, the crescent-shaped hole of the aperture is well visible .....3  
– In frontal view, the hole of the aperture is almost or entirely covered .....5
3. One knob and a basal thickening on the peristome .....*V. (V.) gittenbergeri* Fehér & Eröss sp. nov.  
– Two knobs on the peristome .....4
4. Shell width greater than 5 mm .....*V. (V.) pesici* Fehér & Deli sp. nov.  
– Shell width less than 5 mm .....*V. (V.) deelemanorum* Gittenberger, 1969
5. Upper side of the shell is entirely flat in frontal view .....*V. (V.) stojaspali* Reischütz *et al.*, 2009  
– Shell is conical in frontal view .....6
6. No knobs on the peristome .....7  
– At least one knob on the peristome .....8
7. Very narrow umbilicus .....*V. (V.) aspectulabeatidis* Reischütz *et al.* in Reischütz & Reischütz, 2009  
– Wide, perspectival umbilicus .....*V. (V.) pageti* Gittenberger, 1969
8. Less than three knobs on the peristome .....9  
– Three knobs on the peristome .....10
9. An elongated basal thickening and a detached palatal knob on the peristome .....  
.....*V. (V.) backhuysi* Gittenberger, 1969  
– Palatal knob is fused to the basal thickening of the peristome .....  
.....*V. (V.) gittenbergeri* Fehér & Eröss sp. nov.
10. In bottom-view the umbilicus seems wider than  $\frac{1}{4}$  of the shell width, in frontal view the  
shell is flat conical ( $H:W < 2:3$ ) .....*V. (V.) adrianae* Gittenberger, 1969  
– In bottom-view the umbilicus seems narrower than  $\frac{1}{4}$  of the shell width, in frontal view the  
shell is conical ( $H:W \approx 2:3$ ) .....*V. (V.) nicoleae* Reischütz & Reischütz, 2012

## *Virpazaria (Virpazaria) adrianae* Gittenberger, 1969

Fig. 1A–F

*Virpazaria (Virpazaria) adrianae* Gittenberger, 1969: 298–299, fig. 4.

*Virpazaria (Virpazaria) adrianae* – Gittenberger 1975: 268. — Welter-Schultes 2012: 211, unnumbered fig.

## Diagnosis

Discoid, shouldered shell, barely raised spire, wide umbilicus. Aperture crescent-shaped, most of it but the sinus region covered by a parietal flap in frontal view. Three wide knobs at basal–palatal peristome.

## Material examined

### Holotype

MONTENEGRO • W of Virpazar, Velika Jama, at Mt. Soko near Dupilo; Dabović leg.; “*Trissexodon kušćeri*” [original label]; NHMW-E 57441.

### Paratypes

MONTENEGRO • >30 spec.; same collection data as for holotype; NHMW-E 57441a • 1 spec.; same collection data as for holotype; SMF123347 • 2 spec.; same collection data as for holotype; RMNH 53303 • 1 spec.; same collection data as for holotype; NHMW-K 29619 • 2 spec.; same locality as holotype; Kuščer leg.; Apr.1936; SMF 53886.

### Other material

MONTENEGRO • 6 spec.; “Grotte Grbočica” [Pečina Grbočica]; 10 Jul. 1936; Dabović leg.; [on label indicated as paratype]; NHMW-E 48634 • 3 spec.; “Grotte Grbočica” [Pečina Grbočica]; 10 Jul. 1936; Dabović leg.; [on label indicated as paratype]; NHMW-E 48092 • 1 spec.; Vrelo kod Jabukovog dola; Dabović leg.; [on label indicated as paratype]; NHMW-E 48486 • 3 spec.; [without locality]; “*Trissexodon montenegrina* n. sp. Smmlg. Kuščer, Smmlg. Edlauer”; ex. Kuščer; NHMW 79007 • many spec.; Sokopečina; ex. Edlauer; NHMW 79008 • 7 spec.; Pečina na ostrvu Vranjini [cave on Vranjina Island]; Kuščer leg.; NHMW 78990 • Pečina na ostrvu Vranjini [cave on Vranjina Island]; Kuščer leg.; NHMW-K 64035 • 3.5 km N of Virpazar, on road to Rijeka Crnojevića, Vidikovac Observation Point; 42.2547° N, 19.0896° E; 150 m alt.; 25 May 2015; T. Deli, Z. Eröss and Z. Fehér leg.; HNHM 103419 • W of Virpazar, 1.2 km from Dupilo, S side of Mt Tor; 42.2527° N, 19.0662° E; 85 m alt.; 20 Oct. 2017; T. Deli, Z. Eröss, A. Hunyadi and B. Páll-Gergely leg.; DT • W of Virpazar, 1.2 km from Dupilo, S side of Mt Tor; 42.2527° N, 19.0662° E; 85 m alt.; 20 Oct. 2017; T. Deli, Z. Eröss, A. Hunyadi and B. Páll-Gergely leg.; ER • NNW of Virpazar, 0.4 km W of Poseljani, on road to Rijeka Crnojevića; 42.3060° N, 19.0483° E; 130 m alt.; 20 Oct. 2017; T. Deli, Z. Eröss, A. Hunyadi and B. Páll-Gergely leg.; DT • NNW of Virpazar, 0.4 km W of Poseljani, on road to Rijeka Crnojevića; 42.3060° N, 19.0483° E; 130 m alt.; 20 Oct. 2017; T. Deli, Z. Eröss, A. Hunyadi and B. Páll-Gergely leg.; ER • Začir, Pečina u Pečkom Brdu, above village; 20 Jul. 1972; L. Pintér, P. Subai and A. Szigethy leg.; HNHM 72071 • Komarno, Jabukov Do Cave; 3 Jun. 1978; W.J.M. Maassen leg.; NMBE 539837.

### Dimensions

H 1.6–1.9 mm; W 3.0–3.5 mm.

### Distribution

This is the most frequent species of the genus. Its range is northwest of Lake Skadar in Montenegro. The following locations are mentioned in the literature: Velika Jama near Dupilo, Pečina Grbočica near Trnovo, Pečina u Pečkom Brdu near Začir, Pečina Ostrin Vranjini northeast of Virpazar, Babotuša Spilja near Trnovo and Jabukov do Pečina near Komarno (Gittenberger 1969, 1975). Recently discovered new sites fall within the boundaries of the previously known range. All georeferenced records (and probably the non-georeferenced ones too) came from Lower Triassic limestone (Fig. 2).

*Virpazaria (Virpazaria) aspectulabeatidis* Reischütz *et al.* in Reischütz & Reischütz, 2009

*Virpazaria aspectulabeatidis* A. Reischütz, P.L. Reischütz & Subai in A. Reischütz & P.L. Reischütz, 2009: 56, figs 4–5.

*Virpazaria aspectulabeatidis* – Subai 2009: 109, fig. 9. — Welter-Schultes 2012: 211.

### Diagnosis

Discoid, shouldered shell, low but visibly raised spire. 60–65 ribs on body whorl. Aperture very narrow, crescent-shaped, entirely covered by a parietal flap in frontal view. In bottom view, very narrow umbilicus, parietal flap and palatal peristome form like two straight parallel lines.


## Material examined

### Holotype

MONTENEGRO • Rumija Mts, rocks above Bes; Jul. 2008; A. Reischütz, N. Reischütz and P.L. Reischütz leg.; NHMW 106689.

### Paratypes

MONTENEGRO • 1 spec.; Mt Kroništár, S slope, approx. 10 km E of Đuravci (in direction to Ostros, S of Lake Skadar); 350 m alt.; 16 Sep. 2006; P. Subai leg.; SMF 333501 • 1 spec.; Mt Kroništár, S slope,



**Fig. 1.** Shells of species of *Virpazaria* Gittenberger, 1969. **A–F.** *Virpazaria (Virpazaria) adrianae* Gittenberger, 1969, holotype (NHMW-E 57441). **G–L.** *Virpazaria (Virpazaria) backhuysi* Gittenberger, 1969, holotype (NHMW-E 48856). Scale bar = 1 mm.

approx. 10 km E of Đuravci (in direction to Ostros, S of Lake Skadar); 350 m alt.; 16 Sep. 2006; P. Subai leg.; NMBE 539842.

#### Other material

MONTENEGRO • Rumija Mts, Besa, Krone i Besit, 2 km S of Donji Murići junction on Virpazar–Ostros road; 320 m alt.; 42.1424° N, 19.2197° E; 19 Oct. 2017; T. Deli, Z. Erőss, A. Hunyadi and B. Páll-Gergely leg.; DT.

#### Dimensions

H 2.1–2.3 mm; W 3–3.5 mm.

#### Distribution

This species has a very restricted range west of Lake Shkodër at the foot of the Rumija Mountains in Montenegro (Reischütz & Reischütz 2009; Subai 2009). Apart from the type locality there is another literature record, namely “Đuravci, 10 km to Ostros” (Subai 2009). None of the published distribution records were georeferenced; thus, it is difficult to say how far these locations are from one another. The site where we found this species is likely identical to, or very close to, the type locality and its bedrock type is Upper–Middle Jurassic limestone. At all three sites, this species was reported to co-occur with *V. stojaspali* (Fig. 2).

#### *Virpazaria (Virpazaria) backhuysi* Gittenberger, 1969

Fig. 1G–L

*Virpazaria (Virpazaria) backhuysi* Gittenberger, 1969: 299–301, fig. 5.

*Virpazaria (Virpazaria) backhuysi* – Welter-Schultes 2012: 212.

#### Diagnosis

Small, discoid and shouldered shell, low but raised spire, wide umbilicus. Callous thickening along entire basal peristome, a smaller knob on palatal peristome. High and thick parietal flap reaching almost up to angular junction. Aperture crescent-shaped, all of it but the sinus region covered by a parietal flap in frontal view.

#### Material examined

##### Holotype

MONTENEGRO • Virpazar, “Cp.360”; Dabović leg.; NHMW-E 48856.

#### Dimensions

H 1.5 mm; W 2.5 mm (holotype).

#### Distribution

Only the holotype specimen is known, which came from the Edlauer Collection and was probably collected in the 1930s. On the label only “Virpazar Cp.360” is stated. Virpazar is a small historical town with a fortress. In past decades, various malacologists have sampled in and around Virpazar, including E. Gittenberger, H. Nordsieck, A. and P.L. Reischütz, P. Subai and the authors of the present paper, but no one has been able to refind this enigmatic species. This either means that *V. backhuysi* has a very narrow range and/or is a locally very rare species, or that we searched in all the wrong places. In case of old museum material, the possibility of a past mix-up can never be excluded (see also the doubt regarding the real origin of the type series of *V. ripkeni*), but it is also feasible that the label information refers to

the area around Virpazar in a broader sense. Due to this uncertainty we cannot conclude anything about its preferred bedrock type.

*Virpazaria (Virpazaria) deelemanorum deelemanorum* Gittenberger, 1975

*Virpazaria (Virpazaria) deelemanorum* Gittenberger, 1975: 268–269, figs 3 (shell), 4–5 (genitals), table I.1.

*Virpazaria (Virpazaria) deelemanorum* – Welter-Schultes 2012: 212 (in part).

**Diagnosis**

Discoid shell, hardly emerging spire, body whorl silhouette symmetrically rounded. Wide umbilicus. Aperture crescent-shaped, mostly visible in frontal view. Two – more or less equal-sized – knobs on palatal peristome. Callous thickening on peristome between its columellar junction and upper palatal knob.


**Material examined**

**Holotype**

MONTENEGRO • E of Podgorica, Mt. Fundina, Čafa Pješatica Cave; A.P.B. Deeleman leg.; RMNH 55031.

**Paratypes**

MONTENEGRO • 1 spec.; same collection data as for holotype; includes genital preparation RMNH 806a,b; RMNH 55032 • 1 spec.; same locality as for holotype; E. Gittenberger leg.; RMNH 55033 • 1 spec.; Pećina od Zavora near Peuta, ENE of Podgorica; Deeleman-Reinhold leg.; RMNH 55034 [not seen].



**Fig. 2.** Distribution of species of *Virpazaria* Gittenberger, 1969 in the western part of the Balkan Peninsula.

### Dimensions

H 2.1–2.3 mm; W 3.7–4.2 mm.

*Virpazaria (Virpazaria) deelemanorum dhorai* Reischütz *et al.*, 2010

Fig. 3A–E

*Virpazaria dhorai* A. Reischütz, N. Reischütz & P. L. Reischütz, 2010: 29, figs 1–2.

*Virpazaria deelemanorum* – Welter-Schultes 2012: 212 (in part).

*Virpazaria (Virpazaria) deelemanorum dhorai* – Gittenberger 2015: 27, figs 9–10.

### Diagnosis

Compared to nominate subspecies, shell somewhat smaller and lower palatal knob palpably larger.

### Material examined

#### Holotype

ALBANIA • rocks between Xhajë and Makaj, along Koplik–Bogë road; Jul. 2009; A. Reischütz, N. Reischütz and P.L. Reischütz leg.; NHMW 107224.

#### Other material

ALBANIA • Shkodër district, Drisht, right bank of Kiri River opposite fortress hill; 42.1304° N, 19.609° E; 90 m alt.; 28 Jun. 2016; Eröss, Fehér, Szekeres and Grego leg.; HNHM 103421 • Shkodër District, Drisht, right bank of Kiri River opposite fortress hill; 42.1304° N, 19.609° E; 90 m alt.; 28 Jun. 2016; Eröss, Fehér, Szekeres and Grego leg.; NHMW 110430/MN/0987 • Malësia district, 0.5 km NE of Xhajë; 42.3306° N, 19.6057° E; 650 m alt.; 4 Jul. 2015; Eröss, Fehér and Grego leg.; HNHM 103423 • Malësia district, 0.5 km NE of Xhajë; 42.3306° N, 19.6057° E; 650 m alt.; 4 Jul. 2015; Eröss, Fehér and Grego leg.; NHMW 110430/MN/0988.

### Dimensions


H 1.8–2.25 mm; W 3.4–4.2 mm.

### Remarks

Gittenberger (2015) mentioned that the protoconch of *V. d. dhorai* is coarsely granular and that the teleoconch has some irregular radial lines and a very fine microsculpture of incised spiral lines. This, however, is discernible on fresh specimens. Only a few specimens of the nominate form (the types) are known and all are too weathered to observe anything about their shell microsculpture; it is also difficult to conclude whether it is a consistent feature that the palatal knobs are equal-sized in the nominate form. Gittenberger (2015) was somewhat uncertain about the systematic position of the two taxa, but due to the limited amount of material of *V. d. deelemanorum*, he kept them as subspecies.

### Distribution

The range of this species can be found east-northeast of the Shkodër Basin (Albania and Montenegro) (Gittenberger 1975). The nominate subspecies is known from two locations at Mt Fundina (northeast of Podgorica in Montenegro), ca 5–6 km from each other, namely Pecina Cafa Pesatica and Pecina od Zavora near Peuta. The subspecies *dhorai* was found east of the Shkodër Basin, in Bogë Valley near Xhajë and in the Kiri River Valley near Drisht. The newly discovered latter site extends the known range of this species southwards and indicates that further research in this area might reveal additional locations and a larger range.



**Fig. 3.** Shells of species of *Virpazaria* Gittenberger, 1969. **A–E.** *Virpazaria (Virpazaria) deelemanorum dhorai* Reischütz *et al.*, 2010. **A–D.** Holotype (NHMW 107224). **E.** Drisht (HNHM 103421). **F–L.** *Virpazaria (Virpazaria) stojaspali* Reischütz *et al.*, 2009. **F–J.** Štegváš Pass (HNHM 103441). **K–L.** Holotype (NHMW 106690). Scale bar = 1 mm.

The occurrences are associated with three different bedrock types, namely Xhajë on Upper Jurassic and Driht on Middle Triassic, whereas the two non-georeferenced sites east of Podgorica are probably on Upper Cretaceous limestone.

*Virpazaria (Virpazaria) gittenbergeri* Fehér & Eröss sp. nov.  
[urn:lsid:zoobank.org:act:C987B11A-418B-48B2-A5D5-B71966765600](https://zoobank.org/urn:lsid:zoobank.org:act:C987B11A-418B-48B2-A5D5-B71966765600)

Fig. 4A–E

*Spelaeodiscus* sp. (juv.) – Subai 2009: 111.

### Differential diagnosis

This species can be distinguished from congeners by its small size and by its having one knob on the palatal peristome, and from species of the related *Spelaeodiscus*, which have one knob/tooth on the peristome by its inferiorly broader rounded body whorl and narrower aperture.

### Etymology

This species is dedicated to and named after Edmund Gittenberger (Leiden, the Netherlands), a prominent malacologist who did pioneering work on the subterranean terrestrial gastropods of the Balkans. The name is used as the genitive of a noun of male gender.

### Material examined

#### Holotype

MONTENEGRO • Rumija Mts, 2 km S of Štegvaš Pass on Ostros–Ulcinj road (4 km S of Arbnež); 42.0582° N, 19.3535° E; 350 m alt.; 28 May 2015; T. Deli, Z. Eröss and Z. Fehér leg.; HNHM 103411.

#### Paratypes

MONTENEGRO • 20 spec.; same collection data as for holotype; HNHM 103413 • 20 spec.; same collection data as for holotype; NHMW 111671 • 20 spec., 38 fr.; same collection data as for holotype; DT • 8 ad., 8 juv.; same locality as for holotype; 16 Jul. 2017; Z. Eröss and Z. Fehér leg.; HNHM 103451 • 8 ad., 9 juv.; same locality as for holotype; 16 Jul. 2017; Z. Eröss and Z. Fehér leg.; NHMW 111674 • 16 ad., 9 juv.; same locality as for holotype; 16 Jul. 2017; Z. Eröss and Z. Fehér leg.; ER • 2 ad., 1 juv.; SW of Štegvaš Pass, 2 km S of Arbnež on Ostros–Ulcinj road; 42.0636° N, 19.3710° E; 480 m alt.; 16 Jul. 2017; Z. Eröss and Z. Fehér leg.; HNHM 103445 • 3 ad., 1 juv.; same collection data as for preceding; NHMW 111673 • 3 ad., 1 juv.; same collection data as for preceding; ER.

#### Other material


ALBANIA • Shkodër district, Mt Tarabosh, Vallas, near mosque; 42.0506° N, 19.4162° E; 190 m alt.; 27 May 2015; T. Deli, Z. Eröss and Z. Fehér leg.; HNHM 103415 • same collection data as for preceding; NHMW 110430/MN/0986 • same collection data as for preceding; DT.

### Dimensions

Holotype: H 1.55 mm; W 2.5 mm; Ha 1.0 mm; Wa 1.1 mm. Paratypes (HNHM 103413, n=5): H 1.4–1.6 mm; W 2.35–2.5 mm. Vallas population (HNHM 103415, n=5): H 1.25–1.5 mm; W 2.2–2.4 mm.

### Description

Shell discoid, spire barely raised, consists of  $4\frac{1}{4}$ – $4\frac{3}{4}$  regularly increasing whorls, of which  $1\frac{1}{2}$ –2 are protoconch whorls. Fresh shells glassy translucent, older shells whitish. Protoconch grained. Teleoconch whorls equidistantly ribbed: 59–69 ribs on body whorl and fine, regular lines in between.



**Fig. 4.** Shells of species of *Virpazaria* Gittenberger, 1969. **A–E.** *Virpazaria (Virpazaria) gittenbergeri* Fehér & Erőss sp. nov., holotype (HNHM 103411). **F–J.** *Virpazaria (Virpazaria) pesici* Fehér & Deli sp. nov., holotype (HNHM 103403). Scale bars = 1 mm (note that the latter species is less magnified).

In bottom view, umbilicus wide,  $\frac{1}{4}$ – $\frac{1}{5}$  of total shell width, a small part of it overlaid by columellar part of aperture. In frontal view, silhouette of body whorl asymmetrically rounded (shouldered): broader curved in bottom, more angulated on upper side and near suture horizontally flattened. Suture barely depressed. Aperture narrow crescent-shaped. Instead of a flap, with an outstretched swelling (a thickened and emerged callus) on parietal side. One knob on palatal peristome, ca at one-third of way between its angular and columellar junctions. Peristome somewhat reflected between columellar junction and palatal knob, appearing thickened in frontal view. Above knob, peristome simple, delineating an apertural sinus. In side view, parietal swelling oblique (inferior part stands forward) and aperture sigmoidally inclined (inferior part stands backward).

### Distribution

This species is known from the southeastern part of the Rumija Mts (also known as Mt Tarabosh). The Štegvaš Pass (Montenegro) and Vallas (Albania) populations are within a distance of 5 km of one another. Both sites are on Upper–Middle Triassic limestone. At the type locality, this species was found together with *V. stojaspali* and *V. pesici*.

### Conservation status

To our present knowledge, this is a narrow-range endemic species with an area of occurrence (AOO) smaller than 20 km<sup>2</sup>. However, there is no reason to suppose that AOO, extent of occurrence (EOO), number of locations, number of subpopulations or the number of mature individuals are declining or extremely fluctuating. Therefore, it might be assessed as Near Threatened (NT).

### Remarks

The Vallas population differs from the typical form by its denser ribbing (76–88 ribs on the body whorl).

*Virpazaria (Virpazaria) nicoleae* Reischütz & Reischütz, 2012  
Fig. 5A–E

*Virpazaria nicoleae* A. Reischütz & P.L. Reischütz, 2012: 17–18, 3 unnumbered figs.

### Diagnosis

Small, shouldered, low conical shell, with conically raised spire, moderately wide umbilicus. Aperture crescent-shaped, most of it apart from sinus region covered by a strong, prominent parietal flap in frontal view. Callous thickening on peristome with a high palatal tooth, very strong basal tooth and an additional lower callous denticle between basal tooth and columellar insertion of peristome.

### Material examined

#### Holotype

MONTENEGRO • 2 km S of Vitoja Spring, near Lake Skadar; Mar. 2010; Reischütz leg.; NHMW 108113.

#### Other material

MONTENEGRO • Podgorica Municipality, 1 km S of Izvor Vitoja, near Lake Shkodra; 42.3196° N, 19.3693° E; 30 m alt.; 5 Jul. 2015; Eröss, Fehér and Grego leg.; HNHM 103425 • same collection data as for preceding; NHMW 110430/MN/0989.

### Dimensions

H 1.7–1.9 mm; W 2.6–2.75 mm.

### Distribution

This species is known only from the type locality. The baserock where it was found is Lower Cretaceous (Neocomian) limestone.

***Virpazaria (Virpazaria) pageti pageti*** Gittenberger, 1969  
Fig. 5F–M

*Virpazaria (Virpazaria) pageti* Gittenberger, 1969: 301–302, fig. 6.

*Virpazaria pageti alexanderi* P.L. Reischütz & Subai in Reischütz *et al.*, 2012: 32, fig. 1.

*Virpazaria (Virpazaria) pageti* – Gittenberger 1988: 424. — Welter-Schultes 2012: 212, unnumbered fig.

*Virpazaria cf. pageti* – Reischütz *et al.* 2011: 58, fig. 3.

*Virpazaria pageti pageti* – Reischütz *et al.* 2012: 33, fig. 3.

### Diagnosis

Discoid shell, barely raised spire, body whorl silhouette rounded. Aperture narrow crescent-shaped. Peristome not thickened, without teeth or knobs. Parietal flap sharp-edged, directly connected with palatal peristome, detached from penultimate whorl (looks like a deformed tube).

### Material examined

#### **Holotype** (*V. pageti*)

MONTENEGRO • “Bocche di Cattaro, Sutorina Mündung (Genist)” [Kotor Bay, in debris at mouth of Sutorina rivulet]; “*Trissexodon Käufeli* Typus”; Käufel leg.; NHMW-E 47864.

#### **Holotype** (*V. p. alexanderi*)

CROATIA • Kotor Bay, Prevlaka Peninsula, near Prevlaka Village, between Osti rt and border station at Njivice; Mar. 2010; A., N. and P.L. Reischütz leg.; NHMW 108106.

#### **Paratypes** (*V. pageti*)

MONTENEGRO • 1 spec.; same collection data as for holotype; NHMW-E 47864a.

#### **Paratypes** (*V. p. alexanderi*)

CROATIA • 1 spec.; same collection data as for holotype; NMBE 539847 • 2 ad., 15 juv.; Reischütz private collection [not seen].

#### **Other material**

CROATIA • Prevlaka Peninsula, 1.5 km from Croatian border station to Fort Prevlaka; 42.4181° N, 18.5132° E [ $\approx$  type locality of *V. alexanderi*]; 60 m alt.; 17 Jul. 2018; Z. Eröss, T. Fehér and Z. Fehér leg.; HNHM 104391.

MONTENEGRO • Igalo, W of Herceg Novi, S of Sutorina-Njivice-Igalo junction, limestone rock cavity; 19 Sep. 2005; P. Subai leg.; HNHM 102250 • Prevlaka Peninsula, Igalo (near Herceg-Novi), W side of bay at Žvinje junction; 42.4455° N, 18.5050° E; 50 m alt.; 29 May 2015; T. Deli, Z. Eröss and Z. Fehér leg.; HNHM 103427 • same collection data as for preceding; NHMW 110430/MN/0990 • beneath Žvinje on serpentine road; 42.4460° N, 18.5029° E; 160 m alt.; 17 Jul. 2018; Z. Eröss, T. Fehér and Z. Fehér leg.; HNHM 104390 • Njivice, ca 2 km from Igalo junction; 42.4316° N, 18.5170° E; 60 m alt.; 18 Jul. 2017; Z. Eröss and Z. Fehér leg.; HNHM 104395.

### Dimensions


H 1.3–1.9 mm; W 2.1–3.2 mm.

*Virpazaria (Virpazaria) pageti kleteckii* Štamol & Subai in Reischütz *et al.*, 2012

*Virpazaria pageti kleteckii* Štamol & Subai in Reischütz *et al.*, 2012: 32, fig. 2.

**Diagnosis**

Distinguished from nominate form by smaller, non-detached aperture, which is less covered by parietal flap in frontal view.



**Fig. 5.** Shells of species of *Virpazaria* Gittenberger, 1969. **A–E.** *Virpazaria (Virpazaria) nicoleae* Reischütz & Reischütz, 2012, holotype (NHMW 108113). **F–M.** *Virpazaria (Virpazaria) pageti* Gittenberger, 1969. **F–I.** Between Fort Prevlaka and the border station (HNHM 104391). **J–M.** Igalo (HNHM 102250). Scale bar = 1 mm.

## Material examined

### Holotype

CROATIA • Konavle Region, Gruda, Vilina Cave; UTM: BN81; 6 Apr. 2010; R. Ozimec leg.; CNHM 10393.

### Paratypes

CROATIA • 2 spec.; same collection data as for holotype; Croatian Biospeleological Society collection [not seen] • 1 spec.; in Dubravka village, ca 5 km NE of Gruda; Mar. 2011; A. and P.L. Reischütz leg.; Reischütz private collection [not seen].

## Dimensions

H 1.5–1.65 mm; W 2.75–3.0 mm.

## Distribution and remarks

This species (including its subspecies) has a restricted range north of Kotor Bay in Croatia and Montenegro (Gittenberger 1969; Reischütz *et al.* 2012). The type lot of the nominate form was collected from debris at the mouth of the Sutorina River. Later, this species was found in rock crevices at the eastern side of Prevlaka Peninsula, between Njivice and Igalo very near the type locality (see Gittenberger 1988). At another point of the Prevlaka, ca 3.5 km as the crow flies from the former site, Reischütz *et al.* (2012) reported further shells of this species (four adults and 15 juveniles). They were distinguished from the nominate form by their smaller shell, finer and denser ribbing, smaller parietal flap, as well as the non-detached aperture, and were described as *Virpazaria pageti alexanderi*. In 2018, we were able to obtain a larger sample of *V. p. alexanderi* from the type locality that provided a good opportunity to study size extremes and the variability of character states within this population. In our sample (HNHM 104391), shell width ranged between 2.1 and 3.0 mm, shell height between 1.3 and 1.9 mm, number of whorls between  $3\frac{3}{4}$  and  $4\frac{1}{4}$ , number of ribs on the last whorl between 48 and 83 and we found specimens with detached and non-detached apertures as well. The values and character states given by Gittenberger (1988) for the nominate form and those given by Reischütz *et al.* (2012) for *V. p. alexanderi* all fall within these ranges. This alone might be a good reason for treating them as synonyms, which is only reinforced once their geographical distribution is also taken into consideration. The two type localities are not only close to each other, but are located along the foot of the same limestone hill, which is composed of the same Upper Cretaceous geological formation. Having no geographical or geological barrier in between, it is likely that the two sites contain the same subterranean population and an extensive sampling along the Prevlaka Peninsula might detect this species at further locations.

*Virpazaria pageti kleteckii* was found at two locations somewhat north of the Prevlaka; namely at Vilina Spilja near Gruda and at Dubravka ca 5 km NE of Gruda. The original description was based on four specimens only (three from Vilina Spilja and one from Dubravka). Until further available material makes it possible to test the stability of distinguishing features, *V. p. kleteckii* must be accepted as a distinct subspecies.

*Virpazaria (Virpazaria) pesici* Fehér & Deli sp. nov.

urn:lsid:zoobank.org:act:248898E8-C732-44A5-A599-620ACC478962

Fig. 4F–J

## Differential diagnosis

This species can be distinguished from all congeners by its large size; additionally, from *V. deelemanorum*, which also has two knobs on the peristome, by the smaller knobs and more reduced parietal flap; from

*Spelaeodiscus unidentatus* Bole, 1961 and *S. albanicus* (A.J. Wagner, 1914), which have small knobs on their peristome, by the crescent-shaped aperture and the presence of a parietal flap.

### Etymology

This species is dedicated to and named after Vladimir Pešić (Podgorica, Montenegro) a prominent acarologist and hydrobiologist who significantly contributed to our knowledge of the mollusc fauna of Montenegro. The name is used as the genitive of a noun of male gender.

### Material examined

#### Holotype

MONTENEGRO • Rumija Mts., 2 km S of Štegvaš Pass on Ostros–Ulcinj road, 4 km S of Arbnež; 42.0582° N, 19.3535° E; 350 m alt.; 28 May 2015; T. Deli, Z. Erőss and Z. Fehér leg.; HNHM 103403.

#### Paratypes

MONTENEGRO • 30 ad., 20 fr.; same collection data as for holotype; HNHM 103405 • 23 ad., 18 fr.; same collection data as for holotype; NHMW 111668 • 32 ad., 32 fr.; same collection data as for holotype; DT • 3 fr.; same locality as for holotype; 16 Jul. 2017; Z. Erőss and Z. Fehér leg.; ER • 1 fr.; SW of Štegvaš Pass, 2 km S of Arbnež on Ostros–Ulcinj road; 42.0636° N, 19.3710° E; 480 m alt.; 16 Jul. 2017; Z. Erőss and Z. Fehér leg.; HNHM 103407 • 1 fr.; same collection data as for preceding; NHMW 111669 • 2 fr.; same collection data as for preceding; ER • 1 spec.; 2 km S of Donji Murići junction on Virpazar–Ostros road; 42.1424° N, 19.2197° E; 320 m alt.; 16 Jul. 2017; Z. Erőss and Z. Fehér leg.; HNHM 103409 • 1 spec.; same collection data as for preceding; NHMW 111670 • 2 spec.; same collection data as for preceding; ER.

### Dimensions

Holotype: H 3.15 mm; W 5.8 mm; Ha 2.65 mm; Wa 2.7 mm. Paratypes (HNHM 103405, n=10): H 3.0–3.3 mm; W 5.1–6.0 mm.

### Description

Shell discoid, spire barely raised, consisting of  $4\frac{2}{3}$ –5 regularly increasing whorls, of which  $1\frac{1}{2}$ – $1\frac{3}{4}$  are protoconch whorls. Protoconch smooth. Teleoconch whorls densely and equidistantly ribbed (91–110 ribs on last whorl), no riblets, only some irregular and indistinct lines visible between ribs at 50× magnification.

In bottom view, umbilicus wide,  $\frac{1}{4}$  of total shell width, a small part of it overlayed by columellar part of aperture. In frontal view, silhouette of body whorl more or less regularly rounded: bottom side slightly broader curved. Suture slightly but perceptibly depressed. Aperture crescent-shaped, parietal flap small, thin and weakly attached to shell (often missing from weathered old shells). Two knobs on peristome: a wider basal and a higher (rather tooth-like) palatal. Palatal knob situated somewhat lower than one-third way between angular and columellar junctions of peristome. Peristome somewhat reflected between columellar junction and palatal knob, appearing thickened in frontal view. Above palatal knob, peristome simple. In side view, parietal flap oblique (inferior part stands forward) and aperture sigmoidally inclined (inferior part stands backward).

### Distribution

So far, this species has only been found on the southern side beneath Štegvaš Pass in the Rumija Mts, on Upper–Middle Triassic limestone. It was found there together with *V. stojaspali* and *V. gittenbergeri* sp. nov.

### Conservation status

To our present knowledge this species is very rare, all known records belonging to the same subpopulation and AOO is smaller than 20 km<sup>2</sup>. At the same time, there is no reason to suppose that AOO, EOO, number of locations, number of subpopulations or the number of mature individuals are declining or extremely fluctuating. Therefore, it might be assessed as Near Threatened (NT).

### *Virpazaria (Virpazaria) stojaspali* Reischütz *et al.*, 2009

Fig. 3F–L

*Virpazaria stojaspali* A. Reischütz, P.L. Reischütz & Subai in A. Reischütz & P.L. Reischütz 2009: 58, figs 6–7.

*Virpazaria stojaspali* – Subai 2009: 109, fig. 10. – Welter-Schultes 2012: 212 (in part).

### Diagnosis

Disc-shaped shell, entirely flat on topside. Protoconch relatively large and somewhat prominent. Body whorl silhouette shouldered, umbilicus wide. Aperture very narrow crescent-shaped, entirely covered by a parietal flap in frontal view. Peristome thickened, without knobs.

### Material examined

#### Holotype

MONTENEGRO • Rumija Mts, rocks above Bes; Jul. 2008; A. Reischütz, N. Reischütz and P.L. Reischütz leg.; NHMW 106690.

#### Paratypes

MONTENEGRO • 1 spec.; Mt. Kroništár, S slope, approx. 10 km E of Đuravci, in direction to Ostros, S of Lake Skadar; 350 m alt.; 21 Sep. 2005; P. Subai leg.; SMF 333502 • 29 spec.; same collection data as for preceding; NMBE 539839 • 1 spec.; same locality as for preceding; 16 Sep. 2006; P. Subai leg.; HNHM 96784 • 26 spec.; same collection data as for preceding; NMBE 539838 • 4 spec.; near preceding locality, on W slope; 26 Sep. 2006; P. Subai and M. Szekeres leg.; NMBE 539840.

#### Other material

MONTENEGRO • Donji Murići, above village, at junction of a minor road to Besa; 42.1539° N, 19.2155° E; 200 m alt.; 16 Jul. 2017; Z. Erőss and Z. Fehér leg.; HNHM 103443 • Besa, Krone i Besit, 2 km S of Donji Murići junction on Virpazar–Ostros road; 42.1424° N, 19.2197° E; 320 m alt.; 16 Jul. 2017; Z. Erőss and Z. Fehér leg.; HNHM 103216 • same locality as for preceding; 19 Oct. 2017; T. Deli, Z. Erőss, A. Hunyadi and B. Páll-Gergely leg.; DT • same collection data as for preceding; ER • SE of Virpazar, 7.3 km S (SE) of Đuravci, 1.6 km NNW of Tejani; 42.1134° N, 19.2231° E; 455 m alt.; 19 Oct. 2017; T. Deli, Z. Erőss, A. Hunyadi and B. Páll-Gergely leg.; DT • same collection data as for preceding; ER • Rumija Mts, 2 km S of Arbnež, on Ostros–Ulcinj road, SW of Štegvaš Pass; 42.0636° N, 19.3710° E; 480 m alt.; 16 Jul. 2017; Z. Erőss and Z. Fehér leg.; HNHM 103447 • Rumija Mts, 4 km S of Arbnež, on Ostros–Ulcinj road, 2 km S of Štegvaš Pass; 42.0582° N, 19.3535° E; 350 m alt.; 28 May 2015; T. Deli, Z. Erőss and Z. Fehér leg.; HNHM 103441 • same collection data as for preceding; NHMW 110430/MN/0997 • same locality as for preceding; 16 Jul. 2017; Z. Erőss and Z. Fehér leg.; HNHM 103449.

### Dimensions

H 1.7–2.0 mm; W 2.8–3.3 mm.

## Distribution

This species lives in the Rumija Mts west of Lake Skadar. Type material came from two locations (Reischütz & Reischütz 2009; Subai 2009), but none of them were georeferenced. Our findings extend the known species range and reveal its co-occurrence with three congeneric species, namely *V. aspectulabeatidis*, *V. gittenbergeri* sp. nov. and *V. pesici* sp. nov. The georeferenced occurrences are from Upper–Middle Triassic and from Upper–Middle Jurassic limestone.

## Remarks

Referring to alleged morphological similarity, Welter-Schultes (2012) claimed that this species might be a junior synonym of *V. adrianae* Gittenberger, 1969. In our opinion, the shapes of the shells of these two species are totally different; *V. stojaspali* is the only known species of *Virpazaria* with an entirely flat topside. Furthermore, the apertures are also very different, with three knobs in *V. adrianae* and without any knobs in *V. stojaspali*.

### *Virpazaria (Aemiliella) ripkeni ripkeni* Gittenberger, 1969

Fig. 6A–L

*Virpazaria (Aemiliella) ripkeni* Gittenberger, 1969: 302–303, fig. 7.

*Virpazaria (Aemiliella) pastorpueri* A. Reischütz, N. Reischütz & P. L. Reischütz, 2011: 57, figs 1–2.

*Virpazaria (Aemiliella) ripkeni* – Welter-Schultes 2012: 212.

*Virpazaria (Aemiliella) ripkeni pastorpueri* – Gittenberger 2015: 24, figs 3–4.

## Diagnosis

Discoid shell, somewhat raised spire. Crescent-shaped aperture, thin but prominent parietal flap. Except for sinus area entire peristome thickened, with three protrusions: upper one is external end of palatal lamella, which reaches deeply inside (more than  $\frac{1}{10}$  of a whorl) and melts into a callous structure (which is often non-parallel with palatal lamella); middle one is external end of basal lamella, which reaches deeply inside (more than  $\frac{1}{10}$  of a whorl); lower one is a small denticle at half way between columellar insertion and basal lamella.

## Material examined

### Holotype (*V. ripkeni*)

MONTENEGRO • “Soko Höhle bei Dupilo westl. von Virpazar, Jugosl.” [W of Virpazar, Velika Jama, at Mt Soko near Dupilo = same type locality as for *V. adrianae*, but see Remarks]; NHMW 77153.

### Holotype (*V. pastorpueri*)


ALBANIA • Drisht, small cave beneath ruins of Drisht; Jul. 2010; A. Reischütz, N. Reischütz and P.L. Reischütz leg.; NHMW 107855.

### Paratypes (*V. ripkeni*)

MONTENEGRO • 6 spec.; same collection data as for holotype; “*Trissexodon kusceri* Montenegro Pečina Grbočica bei Virpazar” [original Edlauer label]; NHMW 77153a • 1 spec.; same collection data as for holotype; RMNH 53302.

### Paratypes (*V. pastorpueri*)

ALBANIA • 1 spec.; same collection data as for holotype; NMBE 539850 • further spec.; same collection data as for holotype; Reischütz collection [not seen].



**Fig. 6.** Shells of *Virpazaria (Aemiliella) ripkeni* Gittenberger, 1969. **A–E.** Holotype (NHMW 77153). **F.** Paratype (NHMW 77153a). **G.** Malkolaj (HNHM 103437). **H–I.** Gadoc (HNHM 103429). **J.** Drisht (HNHM 104392). **K.** Holotype of *V. pastorpueri* Reischütz *et al.*, 2011 (NHMW 107855). **L.** Kruni i Baksit (HNHM 103433). Scale bar = 1 mm.

### Other material

ALBANIA • Shkodër District, Gadoc, on Velipojë–Baks–Rjollë road, near Kruni i Gadocit; 41.8980° N, 19.4425° E; 3 m alt.; 27 May 2015; T. Deli, Z. Eröss and Z. Fehér leg.; HNHM 103429 • same collection data as for preceding; NHMW 110430/MN/0991 • 2 km W of Baks–Rjollë, on road to Velipojë; 41.8763° N, 19.4592° E; 10 m alt.; 28 May 2015; T. Deli, Z. Eröss and Z. Fehér leg.; HNHM 103431 • same collection data as for preceding; NHMW 110430/MN/0992 • 3 km W of Baks–Rjollë, Kruni i Baksit; 41.8831° N, 19.4533° E; 5 m alt.; 28 May 2015; T. Deli, Z. Eröss and Z. Fehér leg.; HNHM 103433 • same collection data as for preceding; NHMW 110430/MN/0993 • Qafë Shtiqë (Renc Mt), on Shkodër–Velipojë road; 41.9161° N, 19.4268° E; 50 m alt.; 28 May 2015; T. Deli, Z. Eröss and Z. Fehér leg.; HNHM 103435 • same collection data as for preceding; NHMW 110430/MN/0994 • Malkolaj, at Torovicë junction on Shkodër–Velipojë road; 41.9270° N, 19.4387° E; 2 m alt.; 28 May 2015; T. Deli, Z. Eröss and Z. Fehér leg.; HNHM 103437 • same collection data as for preceding; NHMW 110430/MN/0995 • Shkodër District, Drisht, E slope of Kalaja e Drishtit [type locality of *V. pastorpueri*]; 42.1248° N, 19.6150° E; 190 m alt.; 4 Jul. 2015; Z. Eröss, Z. Fehér and J. Grego leg.; HNHM 103439 • same collection data as for preceding; NHMW 110430/MN/0996 • Drisht, right bank of Kiri River at Shpellë Korbe; 42.1250° N, 19.6031° E; 90 m alt.; 14 Jul. 2018; T. Fehér, Z. Fehér and R. Lika leg.; HNHM 104392.

MONTENEGRO • 1 spec.; same collection data as for holotype; “Albanien Velika Jama bei Soko Dupilo Umgebung von Virpazar lg. Dabović *Trissexodon kusceri* n. sp. Smmlg. Edlauer 57441”, “Innenfallen der Außenwand der Mündung Virpazar, Smmlg. Edlauer” [original Edlauer labels]; on label indicated as paratype; NHMW 77154.

### Dimensions

H 1.5–2.25 mm; W 2.7–4.0 mm (see Remarks for *V. (A.) r. arbeni*).

### *Virpazaria (Aemiliella) ripkeni arbeni* Gittenberger, 2015

*Virpazaria (Aemiliella) ripkeni arbeni* Gittenberger, 2015: 27, fig. 8.

### Diagnosis

The upper palatal lamella which does not or hardly reaches the peristome and the non-thickened, knobless peristome are the features that distinguish this subspecies from the nominate subspecies.

### Material examined

#### Holotype

ALBANIA • Shkodër, Rozafa Cave [= Shpella e Rozafës]; 42.0487° N, 19.4938° E; 18 Oct. 2011; E. Gittenberger leg.; RMNH 5003974 [not seen].

#### Paratypes

ALBANIA • 2 spec.; same locality as for holotype; May 2013; M. Slooff leg.; RMNH 5003975 [not seen].

### Dimensions

H 1.7–1.8 mm; W 3.1–3.2 mm (based on three specimens according to Gittenberger 2015).

### Distribution

Recent records of this species are from the southern rim of the Shkodër Lake Basin, from the surroundings of Shkodër. The Shkodër site is located on Upper Triassic, the Drisht site on Middle Triassic and the

Renc Mts site on Upper Cretaceous limestone. If these were the only known localities, this would already make *V. ripkeni* the widest distributed species of its genus. Strangely, the type locality of the nominate subspecies is located at the northwestern side of the Shkodër Lake Basin, very far from these Albanian sites (but see Remarks).

### Remarks

This species was described on the basis of eight specimens. Their width and height are 3.5–4.0 mm and 2.05–2.25 mm, respectively, and the number of ribs on the last whorl is 73–86. In the original description (Gittenberger 1969) somewhat lower values were given, probably because he applied a different measuring method. The holotype of *V. pastorpueri* is 3.6 mm wide and 2.15 mm high with 60 ribs on its last whorl (although Reischütz *et al.* (2011) provided a smaller width (3.0–3.5 mm), a larger height (2.5 mm) and an average of 65 ribs on the last whorl). Smaller and more tapered shells were defined by Reischütz *et al.* (2011) as the most important features to distinguish *V. pastorpueri* from *V. ripkeni*. Later, Gittenberger (2015) studied a larger series of topotypical *V. pastorpueri* specimens and concluded that their shells are neither smaller nor relatively higher and thicker discoid than those of typical *V. ripkeni*. However, he found that the ribbing of *V. pastorpueri* is less dense (less than 70 ribs on the last whorl) and therefore he preserved it as a distinct subspecies, *V. ripkeni pastorpueri*. In the same paper, Gittenberger described another subspecies, *V. ripkeni arbeni*, which is smaller (H 1.7–1.8 mm; W 3.0–3.1 mm) and has a weakly developed peristome.

We collected several hundred specimens of this species at four different locations in the Renc Mts. Their shell width ranges between 2.7 and 3.6 mm, shell height between 1.5 and 2.1 mm and the number of ribs on the last whorl between 60 and 82. Compared to topotypical *V. pastorpueri*, their size, shape and rib density values largely overlap. Although one might say that they are generally somewhat smaller and more tapered than typical *V. ripkeni*, neither these characters nor the rib density are able to provide a clear-cut distinction. This large variability in rib density refutes Gittenberger's argument (Gittenberger 2015), and makes any distinction between *V. pastorpueri* and *V. ripkeni* meaningless. However, none of the many adult specimens from the Renc Mts have such a weakly developed peristome as in the three specimens that were described as *V. ripkeni arbeni*. Having no more material available from the type locality of *V. r. arbeni* (Shkodër), it is not possible to test whether the entire population consistently has an underdeveloped peristome there or the subspecies was described on the basis of three freak individuals. Until this question is settled, we must preserve *V. ripkeni arbeni*.

The type material of *V. ripkeni* was presumably collected by Petar Dabović in the 1930s, passed somehow to the Viennese private collector Aemilian Edlauer, and finally landed in the NHMW Mollusc Collection. It appears that Edlauer, or someone else who studied his collection, recognized, but never described, three different species of *Virpazaria*. The proposed names were '*Trissexodon kuščeri*', '*Trissexodon montenegrina*' and '*Trissexodon Käufeli*'. The latter name was obviously proposed for *V. pageti*, but the situation is more dubious with the other two. Among the lots of *V. adrianae* we could find both '*Trissexodon kuščeri*' and '*Trissexodon montenegrina*' labels, whereas the name '*Trissexodon kuščeri*' also appears in the material of *V. ripkeni*. Gittenberger (1969) mentioned eight type specimens of *V. ripkeni* from the same original lot (of which one was designated as a holotype, one made its way to the RMNH and six remained in the original vial and kept the original labels (NHMW 77153a)). Interestingly, another site, "Pečina Grbočica bei Virpazar", is indicated on the original label in that vial, whereas "Soko Höhle bei Dupilo", which is given in the description, can be found on a newer label written in someone else's handwriting. Furthermore, there is another lot (NHMW 77154), which is from Soko Cave according to the original label and contains one specimen of *V. ripkeni*, but not mentioned by Gittenberger (1969). Even stranger, this lot contains an original Edlauer label with the same locality, the same voucher number (E-57441) and the same proposed name, '*Trissexodon kusceri*', as the type lot of

*V. adrianae*. It seems very difficult, if not impossible, to trace back whether there was any mixup with the labels or the specimens in the past, and, if so, where exactly the type specimens of *V. ripkeni* came from (Soko?, Grbočica? or from somewhere else?). The fact that neither Gittenberger (1975) nor anyone else has been able to refind *V. ripkeni* at its alleged type locality or anywhere else in the Virpazar area further discredits the validity of the type locality.

## Discussion

Species of *Virpazaria* inhabit the karstic hills surrounding the basin of Lake Shkodër and a small isolated spot in the coastal foothills of the Orjen Mts at the northern part of Kotor Bay. This narrow-range genus comprises ten narrow-range species.

In this paper, we describe two of them as new and also propose the synonymization of *V. ripkeni pastorpuei* and *V. pageti alexanderi* with their nominate forms. At the time of their description, based on the knowledge at that time, they fulfilled the requirements to consider them subspecies, i.e., they seemed to occupy distinct geographic ranges and they seemed to have distinct morphological character(s). Now, however, available new distribution records and larger samples allow us to see them in a new light. Now we know that the intra-population variability of the features believed to be important for subspecific distinction in *V. ripkeni* and *V. pageti* (shell size, shell shape, rib density) is far larger than the inter-population differences. At the same time, we did not change the status of three other subspecies, namely *V. deelemanorum dhorai*, *V. pageti kleteckii* and *V. ripkeni arbeni*, more because we do not have enough data than because we are convinced of their distinct status. Since the genus *Virpazaria* is far from well explored, it can be assumed that after more intense sampling, new populations will be discovered and more material will be available in the future. This will hopefully provide a better foundation on which to test and reevaluate the taxonomic status of the three aforementioned subspecies.

*Virpazaria* is obviously bound to karstic areas, but we could not discover any preference in terms of the type or age of the limestone baserock. Usually, one species of *Virpazaria* is found at a given location, but there are cases when two or three congeners have been found to co-occur. The associated fauna frequently includes other MSS-specific species of *Platyla*, *Vitrea* and *Gyalina*, and occasionally *Pholeotheras euthrix* and species of *Spelaeodiscus*, *Klemmia* and *Agardhiella* as well.

Generally speaking, there were two different types of (not always easily distinguishable) micro-habitats where we sampled: (i) holes that are probably terminal caverns, with clearly visible signs of dissolution of the rock by water and (ii) fissures or cracks in the rock formed mechanically and filled with a deposit of particulate matter. At the moment we still have too few data to draw far-reaching conclusions, but we had the impression during our field work that *Virpazaria* prefers the former, whereas *Agardhiella* and *Spelaeodiscus* prefer the latter type of micro-habitat. This might be a potential explanation of why they cannot be found more often syntopically; however, this idea needs to be statistically tested on a larger dataset (as in Fehér *et al.* 2018).

## Acknowledgements

We are grateful to Anita Eschner (NHMW) and Sigrid Hof (SMF) for allowing access to the museum collections in their care, to Eike Neubert (NMBE) for loaning shells for examination, to Nesrine Akkari (NHMW) for her help in photographing shells, and to Alexander Reischütz, Edmund Gittenberger and an anonymous reviewer for their constructive and helpful review comments. Tamás Fehér, Jozef Grego, András Hunyadi, Barna Páll-Gergely and Miklós Szekeres helped during field trips and Péter Subai provided inspiration and important information about collecting methods. This study was supported by a grant from the Austrian Science Fund (FWF P- 26581-B25) to Z. Fehér.

## References

- Arndt E. & Subai P. 2013. Gastropoda from subterranean habitats in Greece. *Vernate* 31/2012: 213–224.
- Bole J. 1965. Die Vertreter der Gattung *Spelaeodiscus* Brusina 1886 (Gastropoda, Pulmonata) in Jugoslawien. *International Journal of Speleology* 1 (3): 349–356.  
<https://doi.org/10.5038/1827-806X.1.3.9>
- Camacho A.I. 1992. *The Natural History of Biospeleology*. Monografias del M.N.C.N. 7. CSIC, Madrid.
- Culver D.C. & Pipan T. 2014. *Shallow Subterranean Habitats*. Oxford University Press, Oxford.
- Fehér Z., Mason K., Szekeres M., Haring E., Bamberger S., Páll-Gergely B. & Sólmos P. 2018. Range-constrained co-occurrence simulation reveals little niche partitioning among rock-dwelling *Montenegrina* land snails (Gastropoda: Clausiliidae). *Journal of Biogeography* 45 (6): 1444–1457.  
<https://doi.org/10.1111/jbi.13220>
- Gittenberger E. 1969. Beiträge zur Kenntnis der Pupillacea. I. Die Spelaeodiscinae. *Zoologische Mededelingen* 43 (22): 287–306.
- Gittenberger E. 1975. Beiträge zur Kenntnis der Pupillacea. V. Die Spelaeodiscinae, erster Nachtrag. *Zoologische Mededelingen* 48 (23): 263–277.
- Gittenberger E. 1988. Über *Virpazaria (V.) pageti* Gittenberger 1969, *Gyalina (G.) pageti* n. sp. und deren Sekundärfundorte (Mollusca: Pulmonata). *Annalen des Naturhistorischen Museums in Wien* 90B: 423–426.
- Gittenberger E. 2015. Caving for snails in northern Albania. *Basteria* 79 (1–3): 23–29.
- Hudec V. 1965. Neue Erkenntnisse über den Geschlechtsapparat von *Argna bielzi* (Rossmässler) und Bemerkungen zur systematischen Stellung der Gattung *Argna*. *Archiv für Molluskenkunde* 94: 157–165.
- Juberthie C., Delay B. & Bouillon M. 1980. Sur l'existence du milieu souterrain superficiel en zone non calcaire. *Comptes Rendus de l'Académie des Sciences de la France D* 290: 49–52.
- Juberthie C.B., Delay B. & Bouillon M. 1981. Sur l'existence d'un milieu souterrain superficiel en zone calcaire. *Mémoires de Biospéologie* 8: 77–93.
- Kerney M.P. & Cameron R.A.D. 1979. *A Field Guide to the Land Snails of Britain and North-West Europe*. Collins, London.
- Mammola S., Giachino P.M., Piano E., Jones A., Barberis M., Badino G. & Isaia M. 2016. Ecology and sampling techniques of an understudied subterranean habitat: the *Milieu Souterrain Superficiel* (MSS). *The Science of Nature* 103: 88. <https://doi.org/10.1007/s00114-016-1413-9>
- Mikinčić V. 1953. *Geološka Karta FNR Jugoslavije i susednih Zemalja 1 : 500 000*. Naučna Knjiga, Belgrade.
- MMKS 1983. *Harta gjeologjike e RPS të Shqipërisë: Shkalla 1 : 200 000*. Mjeteve Mësimore, Kulturore e Sportive “Hamid Shijaku”, Tirana.
- Páll-Gergely B., Deli T., Eröss Z.P., Reischütz P.L., Reischütz A. & Fehér Z. 2018. Revision of the subterranean genus *Spelaeodiscus* Brusina, 1886 (Gastropoda, Pulmonata, Spelaeodiscidae). *ZooKeys* 769:13–48. <https://doi.org/10.3897/zookeys.769.25258>
- Reischütz A. & Reischütz P.L. 2009. Ein Beitrag zur Kenntnis der Molluskenfauna von Montenegro, nebst Beschreibung zweier neuer Arten der Gattung *Virpazaria* Gittenberger 1969. *Nachrichtenblatt der Ersten Vorarlberger Malakologischen Gesellschaft* 16: 51–60.

- Reischütz A. & Reischütz P.L. 2012. *Virpazaria nicoleae* nov. spec. (Spelaeodiscidae: Pulmonata: Gastropoda) aus Montenegro. *Nachrichtenblatt der Ersten Vorarlberger Malakologischen Gesellschaft* 19: 17–18.
- Reischütz A., Reischütz N. & Reischütz P.L. 2010. Beiträge zur Kenntnis der Molluskenfauna Albaniens. *Nachrichtenblatt der Ersten Vorarlberger Malakologischen Gesellschaft* 17: 29–31.
- Reischütz A., Reischütz N. & Reischütz P.L. 2011. Anmerkungen zur Verbreitung von *Virpazaria* Gittenberger 1969 (Spelaeodiscidae: Gastropoda) und bemerkenswerte Mollusken aus Albanien. *Nachrichtenblatt der Ersten Vorarlberger Malakologischen Gesellschaft* 18: 57–60.
- Reischütz P.L., Štamol V. & Subai P. 2012. Description of two new subspecies of *Virpazaria pageti* Gittenberger 1969 from Croatia (Gastropoda: Pulmonata: Spelaeodiscidae). *Archiv für Molluskenkunde* 141 (1): 31–34. <https://doi.org/10.1127/arch.moll/1869-0963/141/031-034>
- Rendoš M., Čejka T., Šteffek J. & Mock A. 2014. Land snails from subterranean traps exposed in a forested scree slope (Western Carpathians, Slovakia). *Folia Malacologica* 22 (4): 255–261. <https://doi.org/10.12657/folmal.022.022>
- Schileyko A.A. 1998. Treatise on Recent Terrestrial Pulmonate Molluscs, Part 1. Achatinellidae, Amastridae, Orculidae, Strobilopsidae, Spelaeodiscidae, Valloniidae, Cochlicopidae, Pupillidae, Chondrinidae, Pyramidulidae. *Ruthenica, Supplement* 2 (1): 1–127.
- Schneider C.A., Rasband W.S. & Eliceiri K.W. 2012. NIH Image to ImageJ: 25 years of image analysis. *Nature Methods* 9 (7): 671–675. <https://doi.org/10.1038/nmeth.2089>
- Subai P. 2009. New land snails from Montenegro and Albania. *Archiv für Molluskenkunde* 138 (1): 103–112. <https://doi.org/10.1127/arch.moll/0003-9284/138/103-112>
- Subai P. & Dedov I. 2008. A review of the Bulgarian species of *Aspasita* Westerlund, 1889 (Gastropoda; Pulmonata; Spelaeodiscidae), with description of *A. bulgarica* spec. nov. *Basteria* 72: 111–118.
- Welter-Schultes F.W. 2012. *European Non-marine Molluscs, a Guide for Species Identification*. Planet Poster Editions, Göttingen.

*Manuscript received: 16 May 2019*

*Manuscript accepted: 1 August 2019*

*Published on: 26 September 2019*

*Topic editor: Rudy Jocqué*

*Section editor: Thierry Backeljau*

*Desk editor: Danny Eibye-Jacobsen*

Printed versions of all papers are also deposited in the libraries of the institutes that are members of the *EJT* consortium: Muséum national d'Histoire naturelle, Paris, France; Meise Botanic Garden, Belgium; Royal Museum for Central Africa, Tervuren, Belgium; Royal Belgian Institute of Natural Sciences, Brussels, Belgium; Natural History Museum of Denmark, Copenhagen, Denmark; Naturalis Biodiversity Center, Leiden, the Netherlands; Museo Nacional de Ciencias Naturales-CSIC, Madrid, Spain; Real Jardín Botánico de Madrid CSIC, Spain; Zoological Research Museum Alexander Koenig, Bonn, Germany; National Museum, Prague, Czech Republic.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [European Journal of Taxonomy](#)

Jahr/Year: 2019

Band/Volume: [0558](#)

Autor(en)/Author(s): Feher Zoltan, Deli Tamas, Eröss Zoltan, Lika Romilda

Artikel/Article: [Taxonomic revision of the subterranean genus \*Virpazaria\* Gittenberger, 1969 \(Gastropoda, Spelaeodiscidae\) 1-25](#)