

Research article

urn:lsid:zoobank.org:pub:7A993025-6C28-4B38-98C7-683CD96E7885

Sisubiotus hakaiensis sp. nov. (Tardigrada, Macrobiotidae), a new tardigrade species from Calvert Island (British Columbia, Canada)

Matteo VECCHI^{1,*}, Henry CHOONG² & Sara CALHIM³

^{1,3}Department of Biological and Environmental Science, University of Jyväskylä,
PO Box 35, FI-40014 Jyväskylä, Finland.

²Invertebrate Zoology Department, Royal BC Museum, 675 Belleville Street,
Victoria, BC, V8W 9W2, Canada.

*Corresponding author: matteo.vecchi15@gmail.com; matteo.m.vecchi@jyu.fi

²Email: HChoong@royalbcmuseum.bc.ca

³Email: sara.calhim@jyu.fi

¹urn:lsid:zoobank.org:author:94FA8A73-9524-4FFE-93FC-DAADCE33C2A3

²urn:lsid:zoobank.org:author:FD15D310-EE1D-49F2-A46E-DE5D91463AFC

³urn:lsid:zoobank.org:author:B309F420-F7F9-4708-B59E-66880642F93E

Abstract. Tardigrades reports from British Columbia (Canada) trace back to 1908 and numerous species have been recorded from this region, despite the relatively few published sampling studies. We describe by integrative taxonomy (light microscopy morphology, morphometrics, and DNA sequencing) a new tardigrade species, *Sisubiotus hakaiensis* sp. nov. from the British Columbia central coast. The new species has been found in moss collected from a vertical rock outcrop near the Hakai Institute Calvert Island Field Station. *Sisubiotus hakaiensis* sp. nov. differs from all the other known species in the genus by the presence of a labyrinthine layer inside the egg process walls, whereas no consistent differences in the animals were found. This unique egg characteristic therefore required the amendment of the *Sisubiotus* generic diagnosis to account for the presence of the labyrinthine layer inside the egg process walls.

Keywords. Hakai, BC central coast, egg morphology, tardigrades, integrative taxonomy.

Vecchi M., Choong H. & Calhim S. 2022. *Sisubiotus hakaiensis* sp. nov. (Tardigrada, Macrobiotidae), a new tardigrade species from Calvert Island (British Columbia, Canada). *European Journal of Taxonomy* 823: 64–81. <https://doi.org/10.5852/ejt.2022.823.1815>

Introduction

Reports of tardigrades from Canada can be traced back to two publications in the early 20th century: Richters (1908), who identified three species collected from Vancouver, British Columbia during a Pacific expedition from 1896–1897; and Murray (1910), who collected 31 species during his limited Canadian survey. To date, approximately 121 species of tardigrades are known in Canada, and almost half of these (58 species from 32 genera) were recorded from British Columbia (Meyer 2013; Kaczmarek

et al. 2016). Of 58 species known from British Columbia, six have their type locality in the province: *Echiniscus canadensis* Murray, 1910; *Echiniscus reymondi* Marcus, 1928; *Insuetifurca arrowsmithi* (Kathman & Nelson, 1989); *Ursulinius woodsae* (Kathman, 1990); *Macrobiotus occidentalis* Murray, 1910; and *Platicrista cheuleusis* Kathman, 1990. The genus *Sisubiotus* was recently erected (Stec *et al.* 2021a) based on both morphological and molecular data to accommodate for three species previously assigned to *Macrobiotus* C.A.S. Schultze, 1834. This genus can be distinguished from the otherwise similar *Macrobiotus* by the absence of cuticular pores, the presence of elongated teeth in the second band of the oral cavity armature (OCA), and by a characteristic egg morphology (laid free with areolation and conical processes without the labyrinthine layer) (Stec *et al.* 2021a). The first two described species, *S. spectabilis* (Thulin, 1928) and *S. grandis* (Richters, 1911), were considered for many years as species inquirenda (Dastych 1973). Maucci & Pilato (1974) showed that these two species are valid, and they can be discriminated by their morphology. *Sisubiotus wuyishanensis* (Zhang & Sun, 2014), described from China (Zhang & Sun 2014) was deemed to be insufficiently characterized to be differentiated from *S. spectabilis* and *S. grandis* and thus considered as species inquirenda by Stec *et al.* (2021a). Integrative taxonomy integrates multiple lines of evidence from multiple technique (morphological, molecular) to solve taxonomic issues, and it has been providing to be extremely useful in tardigrades taxonomy (see for example Kiosya *et al.* 2021; Stec *et al.* 2021b; Stec & Morek 2022). Thus, using an integrative taxonomy approach we here describe *Sisubiotus hakaiensis* sp. nov. from Calvert Island (British Columbia, Canada), which can be easily differentiated from the other three species in the genus by the presence of a labyrinthine layer in the egg processes. Therefore, we also provide an amended diagnosis of the genus considering this new character state for *Sisubiotus*.

Material and methods

Samples and specimens

Specimens were recovered from moss samples collected by one of the authors (HC) from a vertical rock outcrop near the Hakai Institute Calvert Island Field Station during a biodiversity survey in 2018 and on a subsequent collecting trip by Hakai researchers in 2021. The Field Station is located within the Hakai Lúxvbálís Conservancy of the Province of British Columbia on the central coast of British Columbia. The central coast of British Columbia extends from approximately the north end of Aristazabal Island at ~52°49' N to the entrance to Queen Charlotte Strait at ~50°59' N, excluding the offshore Haida Gwaii Archipelago (Lindstrom *et al.* 2021).

Details on samples and specimens found are shown in Table 1. The samples were examined for tardigrades using the protocol by Stec *et al.* (2015). Animals and eggs were split into several groups for specific analyses, i.e., morphological analysis in PCM (Phase Contrast Microscopy) and DNA sequencing (for details see Table 1).

Microscopy and imaging

Specimens for light microscopy were mounted on microscope slides in a small drop of Hoyer's (~200 mg) medium, secured with a cover slip (22*22 mm) and dried at 60°C for a week. Freshly mounted specimens were checked for the presence of sperm in the gonad (Coughlan & Stec 2019; Coughlan *et al.* 2019). Slides were examined under an Olympus BX53 light microscope with PCM, associated with an Olympus DP74 digital camera. All figures were assembled in Figure J (Mutteter & Zinck 2013). For structures that could not be satisfactorily focused on a single light microscope photograph, a stack of 2–3 images were taken with an equidistance of ca 0.2 µm and assembled manually into a single deep-focus image in GIMP ver. 2-10 (GIMP Development Team 2019). Deep-focus images obtained by stacking are indicated in the figures caption with an asterisk (*).

Table 1. Samples details and number of specimens analyzed [animals + eggs].

Sample and field numbers	Locality	Coordinates for all samples listed	Elevation m a.s.l.	Coll. date	Sample collector	Specimens analyzed [A+E]
S418 (HC2018-14-1)			50	15/06/2018	Henry Choong	PCM Paratypes [2 + 2]
S1910 (GSB-Loc.4)	Steep wall, trail to Lookout, Calvert Island	51°38'54" N, 128°8'38" W	40	29/06/2021	Gillian Sadlier-Brown	PCM Paratypes [0 + 1]
S1911 (GSB-Loc.5)			40	29/06/2021	Gillian Sadlier-Brown	PCM Holotype [1 + 0] + PCM Paratypes [52 + 0] + DNA [3 + 0]

Morphometrics and morphological nomenclature

All measurements are given in micrometers (μm). Sample size was adjusted following the recommendations by Stec *et al.* (2016). Structures were measured only if their orientation was suitable. Body length was measured from the anterior extremity to the posterior end of the body, excluding the hind legs. Buccal tube length and the level of the stylet support insertion point were measured according to Pilato (1981). The *pt* index is the ratio of the length of a given structure to the length of the buccal tube (Pilato 1981). Measurements of buccal tube widths, heights of claws and eggs, as well as the terminology used to describe the OCA and eggshell morphology follow Kaczmarek & Michalczyk (2017). Morphometric data were handled using the “Parachela” ver. 1.7 template available from the Tardigrada Register (Michalczyk & Kaczmarek 2013). The raw morphometric data are provided as Supplementary file 1 (SM.01). Tardigrade taxonomy follows Bertolani *et al.* (2014) and Stec *et al.* (2021a).

Genotyping

DNA was extracted from individual animals following a Chelex® 100 resin (BioRad) extraction method by Casquet *et al.* (2012) with modifications described in detail in Stec *et al.* (2020a). We sequenced four DNA fragments, three nuclear (18S rRNA, 28S rRNA, ITS2) and one mitochondrial (COI). All fragments were amplified and sequenced according to the protocols described in Stec *et al.* (2020a); primers with original references are listed in Appendix 1. Sequencing products were read with an ABI 3130xl sequencer at the Department of Biological and Environmental Sciences (University of Jyväskylä, Finland).

Phylogenetic analysis

The phylogenetic analyses were conducted using concatenated 18S rRNA+28S rRNA+ITS-2+COI sequences. All *Sisubiotus* sequences available in GenBank were used. In addition, representative sequences of Macrobiotidae Thulin, 1928, Adorybiotidae Stec, Vecchi & Michalczyk, 2020, Richtersiusidae Guidetti, Schill, Giovannini, Massa, Goldoni, Ebel, Förschler, Rebecchi & Cesari, 2021 and Murrayidae Guidetti, Rebecchi & Bertolani, 2000 were included as outgroups (Appendix 2). Alignment of 18S and 28S markers was done on reference alignments as in Vecchi *et al.* (2022b). The ITS-2 sequences were aligned using MAFFT ver. 7 (Katoh *et al.* 2002; Katoh & Toh 2008) with the G-INS-i method (thread=4, threadtb=5, threadit=0, reorder, adjust direction, any symbol, max iterate=1000, retree=1, global pair input). The COI sequences were aligned according to their amino acid sequences (translated using the invertebrate mitochondrial code) with the MUSCLE algorithm (Edgar 2004) in MEGA7 with default settings (all gap penalties=0, max iterations=8, clustering method=UPGMB, lambda=24). Alignments were visually inspected and trimmed in MEGA7. Sequences were concatenated with the R package ‘concatipede’ ver. 1.0.0 (Vecchi & Bruneaux 2021). Model selection was performed for each alignment partition (6 in total: 18S rRNA, 28S rRNA, ITS-2 and three COI codons) with PartitionFinder2 (Lanfear *et al.* 2016), partitions and models selection process and results are present in Supplementary file 2 (SM.02). Bayesian Inference (BI) phylogenetic reconstruction was done with MrBayes ver. 3.2.6 (Ronquist *et al.* 2012). Two runs with one cold chain and three heated chains were run for 20 million generations with a burning of 2 million generations, sampling a tree every 1000 generations. Posterior distribution sanity was checked with the Tracer ver. 1.7 (Rambaut *et al.* 2018). MrBayes input file with the input alignment is available as Supplementary file 3 (SM.03). The phylogenetic tree was visualized with FigTree ver. 1.4.4 (Rambaud 2007) and the image was edited with Inkscape 0.92.3 (Bah 2011). The complete phylogenetic tree is available in Supplementary file 4 (SM.04).

Results

Phylogenetic analysis

The phylogenetic analysis (Fig. 1) of the sequences from three individuals confirmed the generic assignation to *Sisubiotus* of the new species. The new species forms a separate clade in sister group

relationship with (*S. spectabilis*+*S. grandis*). The genera *Sisubiotus*, *Macrobiotus*, *Mesobiotus* Vecchi, Cesari, Bertolani, Jönsson, Rebecchi & Guidetti, 2016, *Tenuibiotus* Pilato & Lisi, 2011 and *Paramacrobiotus* Guidetti, Schill, Bertolani, Dandekar & Wolf, 2009 are recovered as monophyletic, whereas the genus *Minibiotus* is paraphyletic (SM.04).

Tardigrade taxa found

Other than the new species of *Sisubiotus*, the analysed samples contained:

Sample S418: *Adorybiotus* cf. *granulatus*.

Sample S1910: *Macrobiotus occidentalis* Murray, 1910.

Sample S1911: *Acanthechiniscus goedeni* (Grigarick, Mihelčič & Schuster, 1964), *Diploechiniscus* sp., *Hypechiniscus gladiator* (Murray, 1905), *Calohypsibius ornatus* (Richters, 1900), *Platicrista* sp., *Adorybiotus* cf. *granulatus*.




Fig. 1. Bayesian phylogenetic placement of the new species of *Sisubiotus* Stec, Vecchi, Calhim & Michalczyk, 2021. Values above/below branches are Bayesian posterior probability values. Scale bar indicates mutations/site.

Systematic and taxonomic account

Phylum Tardigrada Doyère, 1840
Class Eutardigrada Richters, 1926
Superfamily Macrobiotidea Thulin, 1928
Family Macrobiotidae Thulin, 1928

Genus *Sisubiotus* Stec, Vecchi, Calhim & Michalczyk, 2021

Amended diagnosis

Large and whitish Macrobiotidae with: (i) a poreless cuticle, (ii) a wide rigid buccal tube with well-developed oral cavity armature (all three bands of teeth clearly visible in PCM, anterior teeth of the second band longitudinally elongated), (iii) two macroplacoids and a large microplacoid positioned close to them in the pharynx, (iv) Y-shaped claws of the Macrobiotus type with lunules on each leg, (v) ornamented eggs, laid freely, with areolation and conical processes with or without the labyrinthine layer.

Genus composition

Sisubiotus spectabilis (Thulin, 1928) (type species), *Sisubiotus grandis* (Richters, 1911), *Sisubiotus wuyishanensis* (Zhang & Sun, 2014) species inquirenda, *Sisubiotus hakaiensis* Vecchi, Choong & Calhim, 2022 sp. nov.

Sisubiotus hakaiensis sp. nov.

urn:lsid:zoobank.org:act:837BB60F-E17A-4023-959B-EA6E4169198B

Figs 2–5, Tables 2–3

Differential diagnosis

Reliable differences in differentiating the animal morphology between different species of *Sisubiotus* were not found. By the presence of a labyrinthine layer in the egg processes walls, *S. hakaiensis* sp. nov. can be easily differentiated from *S. spectabilis*, *S. grandis* and *S. wuyishanensis* (labyrinthine layer absent in these three species). In addition, *S. hakaiensis* differs from *S. wuyishanensis* by the presence of granulation on legs (present in *S. hakaiensis* vs absent in *S. wuyishanensis*) and by the shape of the egg processes walls (straight to slightly sigmoidal in *S. hakaiensis* vs concave in *S. wuyishanensis*).

Etymology

This species name refers to the Hakai Institute, which conducts and advances long-term scientific research at remote locations at the coastal margin of British Columbia, Canada, and which includes the Calvert Island Field Station from where the samples were collected.

Material examined

58 animals and 3 embryonated eggs. Specimens mounted on microscope slides in Hoyer's medium (55 animals + 3 embryonated eggs) and processed for DNA sequencing (3 animals).

Holotype

CANADA – **British Columbia** • Lookout, Calvert Island, British Columbia; 51°38'54" N, 128°8'38" W; 40 m a.s.l.; 29 Jun. 2021; Gillian Sadlier-Brown leg.; moss on rock; JYUt.S1911_SL5_B.

Paratypes

CANADA – **British Columbia** • 35 animals; Lookout, Calvert Island, British Columbia; 51°38'54" N, 128°8'38" W, 40–50 m a.s.l.; 15 Jun. 2018, 29 Jun. 2021; Henry Choong & Gillian Sadlier-Brown leg.; moss on rock; JYVt.S418_SL2, JYVt.S1911_SL1 to SL6 • 19 animals; same collection data as for

preceding; RBCM.S1911_SL9, SL_10 (RBCM 022-00001-002) • 2 embryonated eggs; same collection data as for preceding; JYVt.S418_SL1 • 1 embryonated egg; same collection data as for preceding; RBCM.S1910_SL2_A (RBCM 022-00001-001).

Voucher specimens are deposited in the Natural history collections of the Jyväskylä University Museum, Ihantolantie 5, Jyväskylä, Finland (JYV), Survontie 9, 40520 Jyväskylä, Finland (Slides JYVt.S1911_SL5, JYVt.S418_SL1-2, JYVt.S1911_SL1 to SL6) and in the Invertebrate Zoology Department, Royal BC Museum (RBCM), 675 Belleville Street, Victoria, BC, Canada (RBCM.S1910_SL2 (RBCM 022-00001-001), RBCM.S1911_SL9, SL10 (RBCM 022-00001-002)) .

Description

Animals (measurements and statistics in Table 2)

Body whitish; after fixation in Hoyer's medium body transparent (Fig. 2A). Eyes present in animals before and after fixation in Hoyer's medium. Cuticle poreless. Patches of fine granulation on the internal and external surfaces of legs I–III (Fig. 2B–C) as well as dorsal and dorso lateral of legs IV clearly visible in PCM (Fig. 2D). A pulvinus is present on the internal surface of legs I–III (Fig. 2B). Claws slender, of the *hufelandi* type. Primary branches with distinct accessory points, a long common tract, and with an evident stalk connecting the claw to the lunula (Fig. 3). All lunulae smooth (Fig. 3A, D). Single cuticular bar on legs I–III often visible in PCM (Fig. 3C), whereas the horseshoe-shaped structure under




Fig. 2. *Sisubiotus hakaiensis* sp. nov., habitus in PCM. **A.** Holotype (JYUt.S1911_SL5_B, Hoyer's medium), dorso-ventral projection. **B***. Paratype (JYUt.S1911_SL3_A). Pulvinus (indented arrowhead) and granulation (arrowhead) on the internal side of leg I. **C–D.** Paratype (JYUt.S1911_SL4_C). **C.** Granulation on external side of leg II. **D.** Granulation on claws IV. Deep-focus images obtained by stacking are indicated in the figures caption with an asterisk (*). Scale bars: A = 100 µm; B–D = 20 µm.

Table 2. Measurements (in μm) of selected morphological structures of the animals of *Sisubiotus hakaensis* sp. nov. mounted in Hoyer's medium. Abbreviations: N = number of eggs/structures measured; Range = the smallest and the largest structure among all measured specimens; SD = standard deviation.

Character	N	Range		Mean		SD		Holotype	
		μm	<i>pt</i>	μm	<i>pt</i>	μm	<i>pt</i>	μm	<i>pt</i>
Body length	25	387–685	707–1070	573	948	72	87	650	1034
Buccopharyngeal tube									
Buccal tube length	30	35.7–73.3	–	58.3	–	8.2	–	62.9	–
Stylet support insertion point	30	30.1–61.7	78.9–85.4	48.5	83.2	7.1	1.4	52.6	83.6
Buccal tube external width	25	5.7–14.4	15.2–21.2	10.6	18.4	2.1	1.4	11.9	19.0
Buccal tube internal width	25	3.7–11.5	10.3–17.7	8.3	14.4	1.7	1.5	10.0	15.8
Ventral lamina length	25	25.5–51.7	65.2–75.1	40.9	70.7	6.2	2.6	42.2	67.1
Placoid lengths									
Macroplacoid 1	30	7.1–21.0	17.7–36.4	16.0	27.3	3.5	4.0	17.7	28.1
Macroplacoid 2	30	4.5–15.5	12.7–21.2	10.2	17.3	2.2	1.9	11.6	18.4
Microplacoid	30	3.1–7.6	7.3–11.4	5.5	9.4	1.0	1.0	5.7	9.1
Macroplacoid row	30	13.5–37.0	34.6–57.2	29.0	49.3	6.0	5.4	32.1	51.0
Placoid row	30	17.9–46.1	46.0–72.5	36.9	62.7	7.2	6.1	41.8	66.4
Claw 1 heights									
External primary branch	19	7.4–15.5	19.4–23.2	12.8	21.5	1.7	1.0	13.7	21.8
External secondary branch	19	5.4–12.0	14.9–18.0	10.0	16.8	1.4	0.9	10.3	16.4
Internal primary branch	18	9.6–13.7	17.1–21.9	12.3	20.3	1.1	1.3	13.7	21.8
Internal secondary branch	17	8.4–10.9	12.6–17.6	9.8	16.1	0.7	1.3	10.9	17.4
Claw 2 heights									
External primary branch	22	6.9–14.5	19.2–24.2	12.9	21.5	1.6	1.6	14.2	22.5
External secondary branch	22	4.5–12.2	12.5–18.9	9.8	16.3	1.6	1.6	11.5	18.3
Internal primary branch	22	5.6–14.4	15.6–23.0	12.2	20.3	1.8	1.9	14.4	22.9
Internal secondary branch	20	8.8–11.3	14.5–17.9	9.9	16.4	0.7	1.0	11.0	17.4
Claw 3 heights									
External primary branch	18	8.1–15.2	18.9–23.9	13.0	22.0	1.6	1.2	14.2	22.6
External secondary branch	16	6.7–12.6	15.5–18.7	10.3	17.5	1.4	1.0	11.3	17.9
Internal primary branch	19	7.6–14.6	17.8–23.1	12.6	21.3	1.6	1.5	14.5	23.0
Internal secondary branch	18	5.3–12.0	13.5–18.8	9.9	16.9	1.4	1.4	11.6	18.4
Claw 4 heights									
Anterior primary branch	19	9.8–20.1	22.0–37.3	17.1	29.6	2.4	3.5	20.1	31.9
Anterior secondary branch	19	6.3–14.5	15.7–26.8	12.2	20.8	1.9	2.7	14.5	23.1
Posterior primary branch	20	8.8–23.6	21.5–43.1	16.1	27.7	2.8	4.7	19.4	30.8
Posterior secondary branch	18	6.1–13.8	14.6–23.0	11.3	19.3	1.8	2.4	13.4	21.3

claws IV poorly visible only in PCM (Fig. 3D). Mouth antero-ventral. Bucco-pharyngeal apparatus of the *Macrobiotus* type (Fig. 4A), with the ventral lamina and ten peribuccal lamellae. Pharyngeal bulb spherical, with trapezoidal apophyses with a median constriction, two rod-shaped macroplacoids and a large microplacoid positioned close to them; Fig. 4B–C). The macroplacoid length sequence is $2 < 1$. The first macroplacoid is anteriorly narrowed and constricted in the middle whereas the second has a sub-terminal constriction (Fig. 4B–C). The oral cavity armature well developed and composed of three bands of teeth, always clearly visible under PCM (Fig. 4D–E). The first band of teeth is composed of numerous small teeth visible in PCM as granules (Fig. 4D–E), arranged in several rows, situated anteriorly in the oral cavity, which start behind the bases of the peribuccal lamellae and extend on the lamellae bases. The second band of teeth is situated between the ring fold and the third band of teeth and comprised of 3–4 rows of teeth visible in PCM as granules (Fig. 4D–E) larger than those in the first band. The most anterior row of teeth within the second band comprises larger and longitudinally elongated teeth than the subsequent posterior rows (Fig. 4D–E). The teeth of the third band are located within the posterior portion of the oral cavity, between the second band of teeth and the buccal tube anterior ending (Fig. 4D–E). The third band of teeth is divided into the dorsal and the ventral portion. Under PCM, both bands are divided into three distinct transverse ridges, with the medio-dorsal larger than the medio-ventral one. In some specimens, additional mucrones can occur behind the medio-ventral ridge (Fig. 4E). Typically-shaped stylet furca, with spherical condyles supported by short branches provided with small apophyses (Fig. 4A).




Fig. 3. *Sisubiotus hakaensis* sp. nov., claws in PCM. **A***, **C.** Paratype (JYUt.S1911_SL3_A). Claws II–III. **B**, **D.** Holotype (JYUt.S1911_SL5_B). Claws IV. Arrowhead indicates horseshoe-shaped structure under claws IV. Deep-focus images obtained by stacking are indicated in the figures caption with an asterisk (*). Scale bars =10 μ m.




Fig. 4. *Sisubiotus hakaensis* sp. nov., buccal-pharyngeal apparatus in PCM. **A, D–E.** Paratype (JYUt. S1911_SL4_C). **B–C.** Paratype (JYUt.S1911_SL2_B). **A.** In toto buccal-pharyngeal apparatus. **B–C.** Placoids, arrowheads indicate constrictions in the macroplacoids. **D*.** Dorsal Oral Cavity Armature (OCA), empty arrowheads indicate the three bands of the OCA. **E*.** OCA, empty arrowheads indicate the three bands of the OCA. Deep-focus images obtained by stacking are indicated in the figures caption with an asterisk (*). Scale bars: A = 50 µm; B–E = 20 µm.

Eggs (measurements and statistics in Table 3)

Laid freely, white, spherical with large conical processes, areolated (Fig. 5A). About ten processes on the circumference. Each process is surrounded by usually eight to twelve deep areolae. Usually, two rows of areolae are present between the neighboring processes (Fig. 5A). The areolae rims are thin and high, and the areolae surface is reticulated (Fig. 5B). The labyrinthine layer between the process walls present and composed by a very fine mesh (Fig. 5C–D). Processes walls straight to slightly sigmoidal, and processes tips usually blunted or flat (Fig. 5C–D).




Fig. 5. *Sisubiotus hakaiensis* sp. nov. Paratype (JYUt.S418_SL1_A). Eggs in PCM. **A.** Egg surface. **B.** Reticulated chorion between areolations. **C.** Egg process with labyrinthine layer (arrowhead); seen as reticulation. **D.** Egg process in section showing labyrinthine layer (arrowhead). Scale bars: A = 50 μm ; B = 5 μm ; C = 20 μm ; D = 10 μm .

Table 3. Measurements (in μm) of selected morphological structures of the eggs of *Sisubiotus hakaiensis* sp. nov. mounted in Hoyer's medium. Abbreviations: N = number of eggs/structures measured; Range = the smallest and the largest structure among all measured specimens; SD = standard deviation.

Character	N	Range	Mean	SD
Egg bare diameter	2	126.0–134.0	130.0	5.7
Egg full diameter	1		171.0	
Process height	9	15.6–31.4	25.8	5.2
Process base width	9	18.6–29.2	23.9	3.0
Process base/height ratio	9	71%–170%	98%	32%
Inter-process distance	6	3.2–11.7	7.5	3.7
Number of processes on the egg circumference	1		10.0	

Reproduction

The examination of animals freshly mounted in Hoyer's medium revealed the presence of testis filled with sperm, so this species can be considered gonochoric.

DNA sequences

Sequences from 3 individuals from sample S1911 were obtained. 18S rRNA (3 sequences: OM523054-6); 28S rRNA (3 sequences: OM523059-61); COI (2 sequences: OM523181-2); ITS2 (2 sequences: OM523057-8).

Discussion

The analyzed animals and eggs belong to a new species in the genus *Sisubiotus*, morphologically and phylogenetically distinct from all other species in the genus. The finding of three embryonated eggs allowed us to link the animal and eggs morphologies. The presence of a labyrinthine layer in the egg processes walls is a new character state for the genus, requiring an amendment of its morphological diagnosis. This newly described species increases the number of tardigrade species recorded from British Columbia to 59, of which seven have their type locality specifically in this province. This new species finding highlights how important is to keep sampling in already explored areas as more biodiversity still awaits to be found (see for example Vuori *et al.* 2020).

The genus *Sisubiotus* has been reported mostly from Europe (Croatia, Finland, France, Italy, Norway, Poland, Romania, Switzerland), the Arctic (Franz Josef Land) and Asia (China, Japan, Siberia-Russia) (McInnes 1994; Stec *et al.* 2021a). Only a few records are known for the Americas: *S. grandis* was found in Greenland and Alaska (Grøngaard *et al.* 1990; Johansson *et al.* 2013) whereas *S. spectabilis* was recorded for Greenland (Maucci 1996), Alaska (USA) (Dastyh 1982; Johansson *et al.* 2013), West Virginia (USA) (Tarter & Nelson 1993) and Argentina (Claps & Rossi 1984). The current description of *S. hakaiensis* sp. nov. is a valuable addition to the scarce records of this genus in the Americas. It is important to note that reticulation in the egg process is a unique feature in an already uncommon and peculiar genus. Thus, it is unlikely to have been overlooked in previous records of *Sisubiotus* from the Americas. Nevertheless, doubts remain with respect to the records of *S. grandis* and *S. spectabilis*: the

considerable distance to their respective type localities and geographical distribution could be indicative of separate (and likely cryptic/pseudocryptic) species being present in the Americas.

Acknowledgments

We are grateful to the Hakai Institute, Tula Foundation, Eric Peterson and Christina Munck for making research on Calvert Island possible. We also wish to thank Gustav Paulay (University of Florida), Matt Lemay (Hakai), and Matt Whalen (UBC-Hakai) for facilitating collecting opportunities, Gillian Sadlier-Brown (Hakai) for collecting part of the samples used in this study, and Hugh MacIntosh (Royal BC Museum) for assistance with the processing of field samples. We are also grateful to Daniel Stec (Institute of Systematics and Evolution of Animals, Polish Academy of Sciences) for acquiring the new species microphotographs. The study was supported by an Academy of Finland Fellowships to S.C. (#314219 and #335759).

References

- Bah T. 2011. Inkscape: guide to a vector drawing program. Upper Saddle River, NJ, USA. Available from <http://tavmjong.free.fr/INKSCAPE/MANUAL/html/> [accessed 13 May 2022].
- Bertolani R., Guidetti R., Marchioro T., Altiero T., Rebecchi L. & Cesari M. 2014. Phylogeny of Eutardigrada: new molecular data and their morphological support lead to the identification of new evolutionary lineages. *Molecular Phylogenetics and Evolution* 76: 110–126. <https://doi.org/10.1016/j.ympev.2014.03.006>
- Casquet J.T., Thebaud C. & Gillespie R.G. 2012. Chelex without boiling, a rapid and easy technique to obtain stable amplifiable DNA from small amounts of ethanol-stored spiders. *Molecular Ecology Resources* 12: 136–141. <https://doi.org/10.1111/j.1755-0998.2011.03073.x>
- Claps M.C. & Rossi G.C. 1984. Contribución al conocimiento de los tardígrados de Argentina, IV. *Acta Zoologica Lilloana* 38: 45–50.
- Coughlan K. & Stec D. 2019. Two new species of the *Macrobotus hufelandi* complex (Tardigrada: Eutardigrada: Macrobiotidae) from Australia and India, with notes on their phylogenetic position. *European Journal of Taxonomy* 573: 1–38. <https://doi.org/10.5852/ejt.2019.573>
- Coughlan K., Michalczyk Ł. & Stec D. 2019. *Macrobotus caelestis* sp. nov., a new tardigrade species (Macrobiotidae: *hufelandi* group) from the Tien Shan Mountains (Kyrgyzstan). *Annales Zoologici* 69: 499–513. <https://doi.org/10.3161/00034541ANZ2019.69.3.002>
- Dastych H. 1973. Redescription of *Macrobotus spectabilis* Thulin 1928 (Tardigrada). *Bulletin of the Polish Academy of Sciences* 21: 823–825.
- Dastych H. 1982. An annotated list of Alaskan Tardigrada. *Polish Polar Research* 3: 95–102.
- Edgar R. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* 32: 1792–1797. <https://doi.org/10.1093/nar/gkh340>
- Folmer O., Black M., Hoeh W., Lutz R. & Vrijenhoek R. 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3: 294–299.
- Gąsiorek P., Stec D., Zawierucha K., Kristensen R.M. & Michalczyk Ł. 2018. Revision of *Testechiniscus* Kristensen, 1987 (Heterotardigrada: Echiniscidae) refutes the polar-temperate distribution of the genus. *Zootaxa* 4472: 261–297. <https://doi.org/10.11646/zootaxa.4472.2.3>
- GIMP Development Team. 2019. GIMP. Available from <https://www.gimp.org> [accessed 13 May 2022].

- Grøngaard A., Kristensen R.M. & Petersen M.K. 1990. Tardigradafaunaen på Disko. *Feltkurs i Arktisk biologi*, Godhavn: 155–179.
- Guil N., Jørgensen A. & Kristensen R. 2019. An upgraded comprehensive multilocus phylogeny of the Tardigrada tree of life. *Zoologica Scripta* 48: 120–137. <https://doi.org/10.1111/zsc.12321>
- Johansson C., Miller W.R., Linder E.T., Adams B.J. & Boreliz-Alvarado E. 2013. Tardigrades of Alaska: distribution patterns, diversity and species richness. *Polar Research* 32: 18793. <https://doi.org/10.3402/polar.v32i0.18793>
- Kaczmarek Ł. & Michalczyk Ł. 2017. The *Macrobiotus hufelandi* (Tardigrada) group revisited. *Zootaxa* 4363: 101–123. <https://doi.org/10.11646/zootaxa.4363.1.4>
- Kaczmarek Ł., Michalczyk Ł. & McInnes S.J. 2016. Annotated zoogeography of non-marine Tardigrada. Part III: North America and Greenland. *Zootaxa* 4203: 1–249. <https://doi.org/10.11646/zootaxa.4203.1.1>
- Kaczmarek Ł., Zawierucha K., Buda J., Stec D., Gawlak M., Michalczyk Ł. & Roszkowska M. 2018. An integrative redescription of the nominal taxon for the *Mesobiotus harmsworthi* group (Tardigrada: Macrobiotidae) leads to descriptions of two new *Mesobiotus* species from Arctic. *PLoS ONE* 13: e0204756. <https://doi.org/10.1371/journal.pone.0204756>
- Katoh K. & Toh H. 2008. Recent developments in the MAFFT multiple sequence alignment program. *Briefings in Bioinformatics* 9: 286–298. <https://doi.org/10.1093/bib/bbn013>
- Katoh K., Misawa K., Kuma K. & Miyata T. 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research* 30: 3059–3066. <https://doi.org/10.1093/nar/gkf436>
- Kiosya Y., Pogwizd J., Matsko Y., Vecchi M. & Stec D. 2021. Phylogenetic position of two *Macrobiotus* species with a revisional note on *Macrobiotus sottilei* Pilato, Kiosya, Lisi & Sabella, 2012 (Tardigrada: Eutardigrada: Macrobiotidae). *Zootaxa* 4933 (1): 113–135. <https://doi.org/10.11646/zootaxa.4933.1.5>
- Lanfear R., Frandsen P.B., Wright A.M., Senfeld T. & Calcott B. 2016. PartitionFinder 2: new methods for selecting partitioned models of evolution for molecular and morphological phylogenetic analyses. *Molecular Biology and Evolution* 34: 772–773. <https://doi.org/10.1093/molbev/msw260>
- Lindstrom S., Lemay M., Starko S., Hind K. & Martone P. 2021. New and interesting seaweed records from the Hakai area of the central coast of British Columbia, Canada: Chlorophyta. *Botanica Marina* 64: 343–361. <https://doi.org/10.1515/bot-2021-0038>
- Mapalo M.A., Stec D., Mirano-Bascos D. & Michalczyk Ł. 2016. *Mesobiotus philippinicus* sp. nov., the first limnoterrestrial tardigrade from the Philippines. *Zootaxa* 4126: 411–426. <https://doi.org/10.11646/zootaxa.4126.3.6>
- Mapalo M.A., Stec D., Mirano-Bascos D. & Michalczyk Ł. 2017. An integrative description of a limnoterrestrial tardigrade from the Philippines, *Mesobiotus insanis*, new species (Eutardigrada: Macrobiotidae: *harmsworthi* group). *The Raffles Bulletin of Zoology* 65: 440–454.
- Maucci W. 1996. Tardigrada of the Arctic tundra with descriptions of two new species. *Zoological Journal of the Linnean Society* 116: 185–204. <https://doi.org/10.1111/j.1096-3642.1996.tb02343.x>
- Maucci W. & Pilato G. 1974. *Macrobiotus spectabilis* Thulin, 1928 e *Macrobiotus grandis* Richters, 1911: due buone species di Eutardigradi. *Animalia* 1: 245–255.
- McInnes S.J. 1994. Zoogeographic distribution of terrestrial/freshwater tardigrades from current literature. *Journal of Natural History* 28: 257–352. <https://doi.org/10.1080/00222939400770131>
- Meyer H.A. 2013. Terrestrial and freshwater Tardigrada of the Americas. *Zootaxa* 3747: 1–71. <https://doi.org/10.11646/zootaxa.3747.1.1>

- Michalczyk Ł. & Kaczmarek Ł. 2013. The Tardigrada Register: a comprehensive online data repository for tardigrade taxonomy. *Journal of Limnology* 72: 175–181. <https://doi.org/10.4081/jlimnol.2013.s1.e22>
- Mironov S.V., Dabert J. & Dabert M. 2012. A new feather mite species of the genus *Proctophyllodes* Robin, 1877 (Astigmata: Proctophyllodidae) from the Long-tailed Tit *Aegithalos caudatus* (Passeriformes: Aegithalidae): morphological description with DNA barcode data. *Zootaxa* 3253: 54–61. <https://doi.org/10.11646/zootaxa.3253.1.2>
- Murray J. 1910. Tardigrada. *Report on Scientific Investigations of the British Antarctic Expedition 1907–1909* 1 (5): 83–187. <https://doi.org/10.5962/bhl.title.22427>
- Mutterer J. & Zinck E. 2013. Quick-and-clean article figures with FigureJ. *Journal of Microscopy* 252: 89–91. <https://doi.org/10.1111/jmi.12069>
- Pilato G. 1981. Analisi di nuovi caratteri nello studio degli Eutardigradi. *Animalia* 8: 51–57.
- Rambaut A. 2007. FigTree, a graphical viewer of phylogenetic trees. Available from <http://tree.bio.ed.ac.uk/software/figtree/> [accessed 13 May 2022].
- Rambaut A., Drummond A., Xie D., Baele G. & Suchard M. 2018. Posterior summarization in Bayesian phylogenetics using Tracer 1.7. *Systematic Biology* 67: 901–904. <https://doi.org/10.1093/sysbio/syy032>
- Richters F. 1908. Beitrag zur Kenntnis der Moosfauna Australiens und der Inseln des Pacifischen Ozeans. *Zoologischer Jahrbücher, Abteilung für Systematik, Geographie und Biologie der Tiere* 26: 196–213. Available from <https://www.biodiversitylibrary.org/page/27521375> [accessed 13 May 2022].
- Ronquist F., Teslenko M., van der Mark P., Ayres D., Darling A., Höhna S., Larget B., Liu L., Suchard M. & Huelsenbeck J. 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61: 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Roszkowska M., Ostrowska M., Stec D., Janko K. & Kaczmarek Ł. 2017. *Macrobotus polypiformis* sp. nov., a new tardigrade (Macrobotidae; *hufelandi* group) from the Ecuadorian Pacific coast, with remarks on the claw abnormalities in eutardigrades. *European Journal of Taxonomy* 327: 1–19. <https://doi.org/10.5852/ejt.2017.327>
- Stec D. & Morek W. 2022. Reaching the monophyly: re-evaluation of the enigmatic species *Tenuibiotus hyperonyx* (Maucci, 1983) and the genus *Tenuibiotus* (Eutardigrada). *Animals* 12: 404. <https://doi.org/10.3390/ani12030404>
- Stec D., Smolak R., Kaczmarek Ł. & Michalczyk Ł. 2015. An integrative description of *Macrobotus paulinae* sp. nov. (Tardigrada: Eutardigrada: Macrobotidae: *hufelandi* group) from Kenya. *Zootaxa* 4052: 501–526. <https://doi.org/10.11646/zootaxa.4052.5.1>
- Stec D., Gąsiorek P., Morek W., Koszyła P., Zawierucha K., Michno K., Kaczmarek Ł., Prokop Z.M. & Michalczyk Ł. 2016. Estimating optimal sample size for tardigrade morphometry. *Zoological Journal of the Linnean Society* 178: 776–784. <https://doi.org/10.1111/zoj.12404>
- Stec D., Zawierucha K. & Michalczyk Ł. 2017. An integrative description of *Ramazzottius subanomalous* (Biserov, 1985) (Tardigrada) from Poland. *Zootaxa* 4300: 403–420. <https://doi.org/10.11646/zootaxa.4300.3.4>
- Stec D., Krzywański Ł. & Michalczyk Ł. 2018a. Integrative description of *Macrobotus canaricus* sp. nov. with notes on *M. recens* (Eutardigrada: Macrobotidae). *European Journal of Taxonomy* 452: 1–36. <https://doi.org/10.5852/ejt.2018.452>
- Stec D., Roszkowska M., Kaczmarek Ł. & Michalczyk Ł. 2018b. *Paramacrobotus lachowskiae*, a new species of Tardigrada from Colombia (Eutardigrada: Parachela: Macrobotidae). *New Zealand Journal of Zoology* 45: 43–60. <https://doi.org/10.1080/03014223.2017.1354896>

- Stec D., Kristensen R.M. & Michalczyk Ł. 2020a. An integrative description of *Minibiotus ioculator* sp. nov. from the Republic of South Africa with notes on *Minibiotus pentannulatus* Londoño *et al.*, 2017 (Tardigrada: Macrobiotidae). *Zoologischer Anzeiger* 286: 117–134. <https://doi.org/10.1016/j.jcz.2020.03.007>
- Stec D., Krzywański Ł., Arakawa K. & Michalczyk Ł. 2020b. A new redescription of *Richtersius coronifer*, supported by transcriptome, provides resources for describing concealed species diversity within the monotypic genus *Richtersius* (Eutardigrada). *Zoological Letters* 6: 1–25. <https://doi.org/10.1186/s40851-020-0154-y>
- Stec D., Krzywański Ł., Zawierucha K. & Michalczyk Ł. 2020c. Untangling systematics of the *Paramacrobiotus areolatus* species complex by an integrative redescription of the nominal species for the group, with multilocus phylogeny and species delineation in the genus *Paramacrobiotus*. *Zoological Journal of the Linnean Society* 188: 694–716. <https://doi.org/10.1093/zoolinnean/zlzl63>
- Stec D., Vecchi M., Maciejowski W. & Michalczyk Ł. 2020d. Resolving the systematics of Richtersiidae by multilocus phylogeny and an integrative redescription of the nominal species for the genus *Crenubiotus* (Tardigrada). *Scientific Reports* 10: 1–20. <https://doi.org/10.1038/s41598-020-75962-1>
- Stec D., Vecchi M., Calhim S. & Michalczyk Ł. 2021a. New multilocus phylogeny reorganises the family Macrobiotidae (Eutardigrada) and unveils complex morphological evolution of the *Macrobiotus hufelandi* group. *Molecular Phylogenetics and Evolution* 160: 106987. <https://doi.org/10.1016/j.ympev.2020.106987>
- Stec D., Vecchi M., Dudziak M., Bartels P. J., Calhim S. & Michalczyk Ł. 2021b. Integrative taxonomy resolves species identities within the *Macrobiotus pallarii* complex (Eutardigrada: Macrobiotidae). *Zoological Letters* 7: 1–45. <https://doi.org/10.1186/s40851-021-00176-w>
- Stec D., Vončina K., Kristensen R.M. & Michalczyk Ł. 2022. The *Macrobiotus ariekammensis* species complex provides evidence for parallel evolution of claw elongation in macrobiotid tardigrades. *Zoological Journal of the Linnean Society* zlab101. <https://doi.org/10.1093/zoolinnean/zlab101>
- Tarter D. & Nelson D. 1993. First records of tardigrades (Phylum: Tardigrada) from mosses in the Cranberry Glades Botanical Area in the Monongahela National Forest, Pocahontas County, West Virginia. *Proceedings of the West Virginia Academy of Science* 65: 40–42.
- Vecchi M. & Bruneaux M. 2021. concatipede: an R package to concatenate fasta sequences easily. <https://doi.org/10.5281/zenodo.5130603>
- Vecchi M., Ferrari C., Stec D. & Calhim S. 2022a. Desiccation risk favours prevalence and diversity of tardigrade communities and influences their trophic structure in alpine ephemeral rock pools. *Hydrobiologia* 849: 1995–2007. <https://doi.org/10.1007/s10750-022-04820-0>
- Vecchi M., Stec D., Vuori T., Ryndov S., Chartrain J. & Calhim S. 2022b. *Macrobiotus naginae* sp. nov., a new xerophilous tardigrade species from Rokua sand dunes (Finland). *Zoological Studies* 61: 21.
- Vuori T., Massa E., Calhim S. & Vecchi M. 2020. Tardigrades of Finland: new records and an annotated checklist. *Zootaxa* 4851 (3): 477–521. <https://doi.org/10.11646/zootaxa.4851.3.3>
- Zawierucha K., Kolicka M. & Kaczmarek Ł. 2016. Re-description of the Arctic tardigrade *Tenuibiotus voronkovi* (Tumanov, 2007) (Eutardigrada; Macrobiotidea), with the first molecular data for the genus. *Zootaxa* 4196 (4): 498–510. <https://doi.org/10.11646/zootaxa.4196.4.2>
- Zhang P. & Sun X.Z. 2014. A new species of the genus *Macrobiotus* (Tardigrada: Macrobiotidae) from Southeastern China. *Zoological Systematics* 39: 224–228.

Manuscript received: 28 February 2022

Manuscript accepted: 12 April 2022

Published on: 10 June 2022

Topic editor: Tony Robillard

Section editor: Daniel Stec

Desk editor: Pepe Fernández

Printed versions of all papers are also deposited in the libraries of the institutes that are members of the *EJT* consortium: Muséum national d’histoire naturelle, Paris, France; Meise Botanic Garden, Belgium; Royal Museum for Central Africa, Tervuren, Belgium; Royal Belgian Institute of Natural Sciences, Brussels, Belgium; Natural History Museum of Denmark, Copenhagen, Denmark; Naturalis Biodiversity Center, Leiden, the Netherlands; Museo Nacional de Ciencias Naturales-CSIC, Madrid, Spain; Real Jardín Botánico de Madrid CSIC, Spain; Zoological Research Museum Alexander Koenig, Bonn, Germany; National Museum, Prague, Czech Republic.

Supplementary files

Supp. file 1. SM.01. Morphometric data for *Sisubiotus hakaiensis* sp. nov.

<https://doi.org/10.5852/ejt.2022.823.1815.6991>

Supp. file 2. SM.02. Model selection results on the concatenated alignment.

<https://doi.org/10.5852/ejt.2022.823.1815.6993>

Supp. file 3. SM.03. MrBayes input file for phylogenetic analysis including the used alignment.

<https://doi.org/10.5852/ejt.2022.823.1815.6995>

Supp. file 4. SM.04. Output phylogenetic tree from MrBayes analysis.

<https://doi.org/10.5852/ejt.2022.823.1815.6997>

Appendix 1. PCR primers for amplification of the four DNA fragments sequenced in the study.

DNA fragment	Primer name	Primer direction	Primer sequence (5'–3')	Primer source
18S rRNA	18S_Tar_1Ff	Forward	AGGCGAAACCGCGAATGGCTC	Stec <i>et al.</i> (2017)
	18S_Tar_1Rr	Reverse	GCCGCAGGCTCCACTCCTGG	
28S rRNA	28S_Eutar_F	Forward	ACCCGCTGAACTTAAGCATAT	Gąsiorek <i>et al.</i> (2018)
	28SR0990	Reverse	CCTTGGTCCGTGTTTCAAGAC	Mironov <i>et al.</i> (2012)
ITS-2	Eutar_Ff	Forward	CGTAACGTGAATTGCAGGAC	Stec <i>et al.</i> (2018a)
	Eutar_Rr	Reverse	TCCTCCGCTTATTGATATGC	
COI	LCO1490	Forward	GGTCAACAAATCATAAAGATATTGG	Folmer <i>et al.</i> (1994)
	HCO2198	Reverse	TAAACTTCAGGGTGACCAAAAAATCA	

Appendix 2. GenBank accession numbers of DNA sequences used in the phylogenetic analysis.

	18S	28S	COI	ITS2	References
<i>Sisubiotus hakaensis</i> sp. nov. 1	OM523054	OM523059	OM523181	OM523057	This study
<i>Sisubiotus hakaensis</i> sp. nov. 2	OM523055	OM523060	OM523182	OM523058	This study
<i>Sisubiotus hakaensis</i> sp. nov. 3	OM523056	OM523061			This study
<i>Sisubiotus spectabilis</i> 1	MN888371	MN888357	MN888323	MN888331	Stec <i>et al.</i> 2021a
<i>Sisubiotus spectabilis</i> 2	MN888371	MN888357	MN888322	MN888331	Stec <i>et al.</i> 2021a
<i>Sisubiotus spectabilis</i> 3	MN888372	MN888364	MN888324	MN888344	Stec <i>et al.</i> 2021a
<i>Sisubiotus grandis</i>		MH079490			Guil <i>et al.</i> 2019
<i>Crenubiotus crenulatus</i>	MT812474	MT812468	MT808079	MT812606	Stec <i>et al.</i> 2020b
<i>Dactylobiotus selenicus</i>	MT812476	MT812466	MT808076	MT812602	Stec <i>et al.</i> 2020b
<i>Macrobiotus ariekammensis</i>	MZ463668	MZ463674	MZ460999	MZ463656	Stec <i>et al.</i> 2022
<i>Macrobiotus pallarii</i>	MT809069	MT809081	MT807924	MT809094	Stec <i>et al.</i> 2021b
<i>Macrobiotus polypiformis</i>	KX810008	KX810009	KX810011	KX810010	Roszkowska <i>et al.</i> 2017
<i>Mesobiotus harmsworthi</i>	MH197146	MH197264	MH195150	MH197154	Kaczmarek <i>et al.</i> 2018
<i>Mesobiotus insanis</i>	MF441488	MF441489	MF441491	MF441490	Mapalo <i>et al.</i> 2017
<i>Mesobiotus philippinicus</i>	KX129793	KX129794	KX129796	KX129795	Mapalo <i>et al.</i> 2016
<i>Minibiotus cf. diversus</i>	OK663227	OK663238	MW306859	OK663216	Vecchi <i>et al.</i> 2022a, 2022b
<i>Minibiotus ioculator</i>	MT023998	MT024041	MT023412	MT024000	Stec <i>et al.</i> 2020a
<i>Minibiotus pentannulatus</i>	MT023999	MT024042	MT023413	MT024001	Stec <i>et al.</i> 2020a
<i>Paramacrobiotus areolatus</i>	MH664931	MH664948	MH675998	MH666080	Stec <i>et al.</i> 2020c
<i>Paramacrobiotus fairbanksi</i>	MH664941	MH664950	MH676011	MH666090	Stec <i>et al.</i> 2020c
<i>Paramacrobiotus lachowskiae</i>	MF568532	MF568533	MF568534	MF568535	Stec <i>et al.</i> 2018b
<i>Paramacrobiotus richtersi</i>	OK663224	OK663236	OK662995	OK663213	Vecchi <i>et al.</i> 2022b
<i>Paramacrobiotus tonollii</i>	MH664946	MH664963	MH676018	MH666096	Stec <i>et al.</i> 2020c
<i>Richtersius coronifer</i>	MH681760	MH681757	MH676053	MH681763	Stec <i>et al.</i> 2020d
<i>Tenuibiotus cf. ciprianoi</i>	MN888376	MN888361	MN888328	MN888348	Stec <i>et al.</i> 2021a
<i>Tenuibiotus danilovi</i>	MN888377	MN888362	MN888329	MN888349	Stec <i>et al.</i> 2021a
<i>Tenuibiotus voronkovi</i>	KX810045	KX810049	KX810042	KX810046	Zawierucha <i>et al.</i> 2016

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [European Journal of Taxonomy](#)

Jahr/Year: 2022

Band/Volume: [0823](#)

Autor(en)/Author(s): Vecchi Matteo, Choong Henry, Calhim Sara

Artikel/Article: [Sisubiotus hakaiensis sp. nov. \(Tardigrada, Macrobiotidae\), a new tardigrade species from Calvert Island \(British Columbia, Canada\) 64-81](#)