

Research article

urn:lsid:zoobank.org:pub:BE1232A7-064A-4D0C-B9BE-05559274B29D

Iron-isopods: new records and new species of terrestrial isopods (Isopoda, Oniscidea) from Brazilian Amazon iron ore caves

Carlos Mario LÓPEZ-OROZCO^{1,*}, Ivanklin Soares CAMPOS-FILHO²,
Jéssica S. GALLO³, Jonas E. GALLÃO⁴, Yesenia M. CARPIO-DÍAZ⁵,
Ricardo BORJA-ARRIETA⁶ & Maria Elina BICHUETTE⁷

^{1,3,4,5,6,7}Laboratório de Estudos Subterrâneos, Universidade Federal de São Carlos,
São Carlos, São Paulo, Brazil.

^{1,5,6}Grupo de Investigación en Biología Descriptiva y Aplicada, Universidad de Cartagena,
Programa de Biología, Campus San Pablo, Cartagena de Indias, Colombia.

²Department of Biological Sciences, University of Cyprus, Lefkosia (Nicosia), Cyprus.

^{3,4,7}Instituto Brasileiro de Estudos Subterrâneos, São Carlos, São Paulo, Brazil.

⁷Grupo Bambuí de Pesquisas Espeleológicas, Belo Horizonte, Minas Gerais, Brazil.

* Corresponding autor: clopezo1610@gmail.com

² Email: ivanklin.filho@gmail.com

³ Email: jessicasgallo@gmail.com

⁴ Email: jonasgallao@gmail.com

⁵ Email: ycarpiod@unicartagena.edu.co

⁶ Email: rborjaa@unicartagena.edu.co

⁷ Email: lina.cave@gmail.com

¹ urn:lsid:zoobank.org:author:8403E42C-C752-425C-934F-D1FA02C36C15

² urn:lsid:zoobank.org:author:31D6AB25-DB38-4E79-AA3F-EEB0D14A4F3F

³ urn:lsid:zoobank.org:author:6EE31621-298E-4C0A-A824-AB926F276C6E

⁴ urn:lsid:zoobank.org:author:4A4E46E9-A722-49FE-91CF-BC9F6B5A07E3

⁵ urn:lsid:zoobank.org:author:1A9C855D-7795-4986-A27E-209AC10843BF

⁶ urn:lsid:zoobank.org:author:1B3ADA69-81B4-49C9-9023-80048E989B17

⁷ urn:lsid:zoobank.org:author:97C7185B-33E2-46F3-AE0E-4FF7CC2B024C

Abstract. Two new species of terrestrial isopods are described from iron ore caves in Brazil, within the Amazon biome, *Circoniscus mendesi* López-Orozco, Campos-Filho & Bichuette sp. nov. and *C. xikrin* López-Orozco, Campos-Filho & Carpio-Díaz sp. nov. (Scleropactidae). In addition, the knowledge of the distribution of *Ctenorillo ferrarai* Campos-Filho, Araujo & Taiti, 2014 (Armadillidae) is extended to Parauapebas, and *Benthanooides tarzan* Cardoso & Ferreira, 2023 to south area of the Campos Ferruginosos National Park, both in the State of Pará. Moreover, a distribution map and photographs of the species are given.

Keywords. Neotropical, iron caves, Parauapebas, Canaã dos Carajás, woodlice.

López-Orozco C.M., Campos-Filho I.S., Gallo J.S., Gallão J.E., Carpio-Díaz Y.M., Borja-Arrieta R. & Bichuette M.E. 2024. Iron-isopods: new records and new species of terrestrial isopods (Isopoda, Oniscidea) from Brazilian Amazon iron ore caves. *European Journal of Taxonomy* 921: 116–135.
<https://doi.org/10.5852/ejt.2024.921.2421>

Introduction

Terrestrial isopods (Oniscidea) represent one of the most diverse groups among crustaceans, including more than 4000 species in more than 500 genera in 38 or 39 families (Schmalfuss 2003; Taiti 2004; Javidkar *et al.* 2015; Sfenthourakis & Taiti 2015; Dimitriou *et al.* 2019). Of these species, more than 300 have been recorded from subterranean environments around the world (e.g., Vandel 1973; Taiti & Gruber 2008; Bedek *et al.* 2011; Taiti & Xue 2012; Tabacaru & Giurginca 2013; Reboleira *et al.* 2015).

To date, more than 230 species of terrestrial isopods are known from Brazil, being the country with the higher diversity of Oniscidea within the Neotropical region (see Campos-Filho *et al.* 2018, 2019, 2020; Cardoso *et al.* 2020a, 2020b, 2021). In the last decade, the knowledge of the diversity of oniscideans associated with subterranean environments considerably increased in the Brazilian territory, from six species in 2011 to 78 in 2023 (Campos-Filho *et al.* 2018, 2019, 2020, 2022a, 2022b, 2023; Cardoso *et al.* 2020a, 2020b, 2021, 2023). Moreover, among them, more than 40 are considered obligatory cave-dwellers, representing more than 50% of the current diversity (Campos-Filho *et al.* 2018, 2019, 2020, 2022a, 2022b, 2023; Fernandes *et al.* 2019; Cardoso *et al.* 2020a, 2020b, 2021, 2023).

The Amazon biome has a great diversity of terrestrial isopods, which makes an important link between the evolutionary aspects of the group in the Neotropics (Lemos de Castro 1984; Leistikow 1999, 2001; Schmidt & Leistikow 2005; Schmidt 2007; Grangeiro *et al.* 2010, 2017, 2021). Regarding the cave fauna of the State of Pará, 18 species in the families Trichoniscidae Sars, 1899, Philosciidae Kinahan, 1857, Scleropactidae Verhoeff, 1938, Dubioniscidae Schultz, 1995, Platyarthridae Verhoeff, 1949, and Armadillidae Brandt, 1833 are currently known, of which eight are considered troglobitic, viz. *Benthana alba* Cardoso & Ferreira, 2023, *Benthanoides amazonicus* Cardoso & Ferreira, 2023, *B. tarzan* Cardoso & Ferreira, 2023, *Amazoniscus eleonora* Souza, Bezerra & Araujo, 2006, *A. leistikowi* Campos-Filho, Araujo & Taiti, 2014, *A. spica* Campos-Filho, Aguiar & Taiti, 2020, *Circoniscus buckupi* Campos-Filho & Araujo, 2011, and *C. carajasensis* Campos-Filho & Araujo, 2011 (Souza *et al.* 2006; Campos-Filho & Araujo 2011; Campos-Filho *et al.* 2014, 2020; Cardoso & Ferreira 2023).

The present study aims to describe two new species of terrestrial isopods from Brazilian iron ore caves within the Amazon biome, State of Pará: *Circoniscus mendesi* López-Orozco, Campos-Filho & Bichuette sp. nov. and *C. xikrin* López-Orozco, Campos-Filho & Carpio-Díaz sp. nov. (Scleropactidae). In addition, the knowledge of the distribution of *Ctenorillo ferrarai* Campos-Filho, Araujo & Taiti, 2014 (Armadillidae) and *B. tarzan* (Philosciidae) is extended.

Material and methods

Study area

The specimens were collected in iron caves of Mineral Province Carajás, in the municipalities of Canaã dos Carajás and Parauapebas, State of Pará, corresponding to the Equatorial Amazonian Domain, at elevations of 600–800 m a.s.l. (Fig. 1A). The Carajás National Forest (FLONA de Carajás, from Portuguese *Floresta Nacional de Carajás*) and Campos Ferruginosos National Park (PARNA dos Campos Ferruginosos from Portuguese *Parque Nacional dos Campos Ferruginosos*) have a high rate of endemism and more than 2000 caves (Campos & Castilho 2012; Piló *et al.* 2015). The vegetation is composed of dense tropical rainforest and canga (savanna systems established on ferrous soils) (Mota




Fig. 1. Study area. **A.** Distribution of the species. **B.** Vegetation of FLONA de Carajás. **C.** Mineral exploitation mine.

et al. 2015). The region has a humid tropical climate, with a rainy period from November to April and a dry season from June to September, an average annual precipitation of over 1500 mm, and an average annual temperature of about 25°C (Köppen 1948; Kottek *et al.* 2006) (Fig. 1B). Although the cavities are inserted in conservation units, mineral exploitation is allowed on FLONA de Carajás, and the largest iron ore mine in the world is there, placing these cave systems under a high level of threat (Fig. 1C) (Carmo *et al.* 2012; ICMBIO 2020).

Collections and taxonomy

Specimens were collected by active search with the aid of tweezers and brushes, and stored in 75% ethanol. Information about the microhabitat of the caves was also recorded. The identifications are based on morphological characters with the use of micropreparations in Hoyer's medium (Anderson 1954). For each new species, the diagnosis, etymology, type material, description, and remarks are given. The illustrations were made with the aid of a camera lucida mounted on Zeiss Stemi SV6 stereo microscope and Leica DMLS microscope. The final illustrations were prepared using the software GIMP (ver. 2.8) with the method proposed by Montesanto (2015, 2016).

The material examined is deposited in the Collection of the Laboratório de Estudos Subterrâneos (LES), Universidade Federal de São Carlos, São Carlos, Brazil (curator: Maria E. Bichuette).

Results

Systematic account

Class Malacostraca Latreille, 1802
Order Isopoda Latreille, 1817
Suborder Oniscidea Latreille, 1802
Family Philosciidae Kinahan, 1857

Genus *Benthanoides* Lemos de Castro, 1958

Type species

Benthanoides pauper (Jackson, 1926), by original designation (see Schmidt & Leistikow 2004).

Benthanoides tarzan Cardoso & Ferreira, 2023
Fig 1

Benthanoides tarzan Cardoso & Ferreira, 2023: 557, figs 2g, 3d, 8–9.

Material examined

BRAZIL – Pará, **Canaã dos Carajás** • 1 ♂; SB_0049 cave (also registered as GEM 1460 and known as Samuel II); 6°18'58.4" S, 49°53'41.1" W; 629 m a.s.l.; 16 Feb. 2022; J.E. Gallão, J.S. Gallo, D.F. Torres and V.F. Sperandei leg.; part in micropreparations; LES 0028764.

Distribution

Presently known from several caves on central part of the PARNA dos Campos Ferruginosos (Cardoso & Ferreira 2023). This work expands its distribution to the southern zone of the park, in SB_0049 cave (also registered as GEM 1460 and known as Samuel II by local people) (Fig. 1).

Family Scleropactidae Verhoeff, 1938

Genus *Circoniscus* Pearse, 1917

Type species

Circoniscus gaigei Pearse, 1917, by monotypy (see Schmidt & Leistikow 2004).

Circoniscus mendesi López-Orozco, Campos-Filho & Bichuette sp. nov.
urn:lsid:zoobank.org:act:5AEA1F23-397B-4C4B-A423-3659311F11C8
Figs 1–4, 8A–B

Diagnosis

Pigment and eyes absent; schisma on pereonite 1 and ventral triangular lobe in pereonite 2; antennula bearing eight lateral aesthetascs arranged in three sets; pleopod 1 and 2 exopods with respiratory areas; pereopod 7 merus bearing triangular lobe on rostral portion; pleopod 5 exopod apex elongated and acute.

Etymology

The new species is named after Francisco Alves Mendes Filho, popularly known as Chico Mendes, a rubber tapper and activist who gained an international reputation for the defense of the Amazonian biodiversity. Furthermore, the present name honors all workers of the ICMBio (in Portuguese, Instituto Chico Mendes de Conservação da Biodiversidade), who continue to keep alive Mendes' ideals.

Material examined

Holotype

BRAZIL – **Pará, Parauapebas** • ♂; CAV-N1-0174; 6°01'27.20" S, 50°17'54.73" W; 649 m a.s.l.; 22 Aug. 2021; M.E. Bichuette, J.E. Gallão and D.F. Torres leg.; LES 0028767.

Paratypes

BRAZIL – **Pará, Parauapebas** • 1 ♂, 1 ♀; same collection data as for holotype; part in micropreparations; LES 0028768 • 3 ♂♂, 4 ♀♀; same collection data as for holotype; LES 0028769 • 1 ♂, 1 ♀; same locality as for holotype; 12 Feb. 2022; J.E. Gallão, J.S. Gallo, D.F. Torres and V.F. Sperandei leg.; LES 0028770 • 3 ♂♂; same collection data as for preceding; LES 0028771.

Description

MEASUREMENTS. Maximum body length: male 5 mm, female 6 mm.

BODY. Color pale yellowish, with typical muscle spots insertions. Exoantennal conglobation (Fig. 2A); body strongly convex (Fig. 2A); dorsal surface smooth, bearing small triangular scale-setae (Fig. 2B). Noduli laterales not discernible at 400× magnification.

CEPHALON. Wider than long, frontal shield obtuse on medial upper margin and straight lateral margins, lateral lobes and eyes absent (Fig. 2C–E).

PEREON. Pereonite 1 epimera with schisma, anterior corners directed frontwards (Fig. 2A, E–F); pereonite 2 with ventral triangular lobes, epimera with outer margin rounded (Fig. 2A, E–F); pereonites 3–4 epimera with outer margin rounded (Fig. 2A); pereonites 5–7 epimera subquadrangular (Fig. 2A).

PLEON. Pleonites 3–5 epimera with outline continuous with that of pereonite 7, apex acute and directed backwards (Fig. 2G); telson triangular, slightly broader than long, with slightly concave sides, rounded apex (Fig. 2G).




Fig. 2. *Circoniscus mendesi* López-Orozco, Campos-Filho & Bichuette sp. nov., ♀, paratype (LES 0028768). **A.** Habitus, lateral view. **B.** Dorsal scale-seta. **C.** Cephalon, dorsal view. **D.** Cephalon, back view. **E.** Cephalon and pereonites 1 and 2, frontal view. **F.** Pereonite 1 and 2 epimera, ventral view. **G.** Pleonites 3–5, telson and uropods. **H.** Antennula. **I.** Antenna, with flagellum detail.

ANTENNULA. Composed of three articles, distal article longest and conical, bearing eight lateral aesthetascs arranged in three sets and apical tip (Fig. 2H).

ANTENNA. Short and stout, not surpassing pereonite 1 when extended backwards; flagellum consisting of two subequal articles, distal article bearing seven lateral aesthetascs in two groups; apical organ longer than distal article of flagellum (Fig. 2I).

MOUTH. Mandibles with molar penicil of many branches; left mandible (Fig. 3A) with 2+1 free penicils, right mandible (Fig. 3B) with 1+1 penicils. Maxillula (Fig. 3C) inner endite with two apical penicils, distal margin bearing outer tip; outer endite with 5+5 teeth, inner set with four teeth apically cleft.




Fig. 3. *Circoniscus mendesi* López-Orozco, Campos-Filho & Bichuette sp. nov., ♀, paratype (LES 0028768). **A.** Left mandible. **B.** Right mandible. **C.** Maxillula. **D.** Maxilla. **E.** Maxilliped, arrow illustrating the endite in caudal view. **F.** Uropod. Figure not to scale.

Maxilla (Fig. 3D) inner lobe rounded and covered with thick setae; outer lobe about twice as wide as inner lobe, covered with thin setae. Maxilliped (Fig. 3E) basis rectangular bearing sparse scale-setae; palp with one setae on proximal article, medial and distal articles with two tufts of setae; endite subrectangular, medial seta long, surpassing distal margin, distal margin covered with thin setae and bearing one seta on outer portion, rostral surface with setose sulcus ending with one short penicil.

UROPOD. Protopod flattened and enlarged, filling gap between pleonite 5 and telson, surpassing distal margin of telson; exopod inserted on median margin, endopod twice as long as exopod and inserted proximally (Fig. 3F).

PEREOPOD 1. Carpus short, transverse antennal grooming brush; dactylus with inner claw surpassing median portion of outer claw, unguis and dactylar organ simple, surpassing distal margin of outer claw (Fig. 4A).

PLEOPOD EXOPODS. Pleopod 1 and 2 exopods with respiratory areas.

Male

PEREOPOD 1. Without particular modifications (Fig. 4A).

PEREOPOD 7. Ischium elongated, sternal margin straight; merus bearing triangular lobe on rostral portion (Fig. 4B).

GENITAL PAPILLA. Triangular ventral shield and subapical orifices (Fig. 4C).

PLEOPODS. Pleopod 1 (Fig. 4C) exopod ovoid; endopod about twice as long as exopod and bent outwards, apex acute and bearing small setae along inner margin. Pleopod 2 (Fig. 4D) exopod triangular, outer margin concave with one seta; endopod flagelliform, longer than exopod. Pleopod 3–4 exopods as in Fig. 4E–F. Pleopod 5 (Fig. 4G) exopod triangular, outer margin almost straight with one tiny setae, inner margin covered with small setae, apex elongated and acute.

Taxonomic remarks

The genus *Circoniscus* comprises 11 species with a distribution restricted to South America (Schmidt 2007; Campos-Filho *et al.* 2017, 2018, 2023; Ocampo-Maceda *et al.* 2022). To date, except for *Circoniscus pallidus* Arcangeli, 1936, in the State of São Paulo, and *Circoniscus caeruleus* Campos-Filho, Sfenthourakis & Bichuette, 2023, in the state of Mato Grosso, all species are recorded from the State of Pará (Campos-Filho *et al.* 2018, 2023). The genus was morphologically defined by Schmidt (2007). Among the species of the genus, *Circoniscus buckupi* Campos-Filho & Araujo, 2011 and *C. carajasensis* Campos-Filho & Araujo, 2011 are considered troglotic, with records from several caves in the State of Pará (Campos-Filho & Araujo 2011; Campos-Filho *et al.* 2022c).

In the absence of body pigments and eyes, *C. mendesi* sp. nov. is similar to *C. buckupi* and *C. carajasensis*; however, it can be distinguished by the triangular sternal lobe of the male pereopod 7 merus, and male pleopod 5 exopod acute. The triangular sternal lobe on the merus 7 is also present in the species of *Circoniscus* in the paper of Campos-Filho *et al.* (2023) from Gruta Lagoa Azul, Bonito, State of Mato Grosso; however, this species is considered to be a troglitic with characters easily distinguishable from *Circoniscus mendesi* sp. nov.

Ecological remarks

The specimens were collected in the twilight zone, living under rocks, in leaf litter, between roots, and in moist soil (Fig. 11A–B). This species is considered to be troglitic due to the absence of body pigments and eyes and to the presence of elongated pereopods.



Fig. 4. *Circoniscus mendesi* López-Orozco, Campos-Filho & Bichuette sp. nov., ♂, paratype (LES 0028768). **A.** Pereopod 1, arrow illustrating the dactylus in rostral view. **B.** Pereopod 7. **C.** Genital papilla and pleopod 1. **D.** Pleopod 2. **E.** Pleopod 3 exopod. **F.** Pleopod 4 exopod. **G.** Pleopod 5 exopod. Figure not to scale.

Distribution

Presently known only from N1_0174 cave, FLONA de Carajás, Parauapebas, Pará, Brazil (Fig. 1).

Circoniscus xikrin López-Orozco, Campos-Filho & Carpio-Díaz sp. nov.

urn:lsid:zoobank.org:act:A22A4823-7BC1-428A-AE59-39C00021300B

Figs 1, 5–7

Diagnosis

Pigment and eyes absent; schisma on pereonite 1, pereonite 2 without ventral lobe; telson with acute apex; antennula bearing eleven lateral aesthetascs arranged in six sets; pleopod 1 and 2 exopods with respiratory areas; pereopod 7 ischium elongated, bearing semicircular lobe on rostral portion, merus bearing semicircular lobe on rostral portion; pleopod 5 exopod apex short and acute.

Etymology

The new species is named for the Xikrin native people, inhabitants of Terras Indígenas Cateté and Trincheira Bacajá, State of Pará.

Material examined

Holotype

BRAZIL – Pará, Canaã dos Carajás • ♂; SB_0049 cave (also registered as GEM 1460 and known as Samuel II by local people); 6°18'58.4" S, 49°53'41.1" W; 629 m a.s.l.; 16 Feb, 2022; J.E. Gallão, J.S. Gallo, D.F. Torres and V.F. Sperandei leg.; part in micropreparations; LES 0028772.

Paratype

BRAZIL – Pará, Parauapebas • 1 ♀; N3_0026 cave; 6°02'40.3" S, 50°13'10.6" W; 18 Feb. 2022; same collectors as for holotype; LES 0028773.

Description

MEASUREMENTS. Maximum body length: male 5.8 mm, female 6 mm.

BODY. Color pale yellowish, with typical muscle spots insertions. Exoantennal conglobation (Fig. 5A); body strongly convex (Fig. 5A); dorsal surface smooth, bearing small fan-shaped scale-setae (Fig. 5B). Noduli laterales not discernible at ×400 magnification.

CEPHALON. Wider than long, frontal shield obtuse on medial upper margin and straight lateral margins, lateral lobes and eyes absent (Fig. 5C–E).

PEREON. Pereonite 1 epimera with schisma, anterior corners directed frontwards (Fig. 5A, E–F); pereonite 2 without ventral lobes, epimera with outer margin rounded (Fig. 5A, E–F); pereonites 3–4 epimera with outer margin rounded (Fig. 5A); pereonites 5–7 epimera subquadrangular (Fig. 5A).

PLEON. Pleonites 3–5 epimera with outline continuous with that of pereonite 7, apex acute and directed backwards (Fig. 5G); telson triangular, broader than long, with slightly concave sides, acute apex (Fig. 5G).

ANTENNULA. Composed of three articles, distal article longest, conical, proximal portion protruding, bearing 11 lateral aesthetascs arranged in six sets plus apical tip (Fig. 5H).




Fig. 5. *Circoniscus xikrin* López-Orozco, Campos-Filho & Carpio-Díaz sp. nov., ♂, holotype (LES 0028772). A. Habitus, lateral view. B. Dorsal scale-seta. C. Cephalon, dorsal view. D. Cephalon, back view. E. Cephalon and pereonites 1 and 2, frontal view. F. Pereonite 1 and 2 epimera, ventral view. G. Pleonites 3–5, telson and uropods. H. Antennula. I. Antenna, with flagellum detail.

ANTENNA. Short and stout, not surpassing pereonite 2 when extended backwards; flagellum consisting of two subequal articles, distal article bearing 17 lateral aesthetascs in two groups; apical organ short than distal article of flagellum (Fig. 5I).

MOUTH. Mandibles with molar penicil of seven branches; left mandible (Fig. 6A) with 2+1 free penicils, right mandible (Fig. 6B) with 1+1 penicils. Maxillula (Fig. 6C) inner endite with two apical penicils, distal margin bearing outer tip; outer endite with 4+6 teeth, inner set with four teeth apically cleft. Maxilla (Fig. 6D) inner and outer lobe rounded and covered with thick setae. Maxilliped (Fig. 6E) as in *C. mendesi* sp. nov.

UROPOD. Protopod flattened and enlarged, filling gap between pleonite 5 and telson, surpassing distal margin of telson; exopod inserted on median margin; endopod short, twice as long as exopod and inserted proximally (Fig. 6F).




Fig. 6. *Circoniscus xikrin* López-Orozco, Campos-Filho & Carpio-Díaz sp. nov., ♂, holotype (LES 0028772). **A.** Left mandible. **B.** Right mandible. **C.** Maxillula. **D.** Maxilla. **E.** Maxilliped, arrow illustrating the endite in caudal view. **F.** Uropod. Figure not to scale.




Fig. 7. *Circoniscus xikrin* López-Orozco, Campos-Filho & Carpio-Díaz sp. nov., ♂, holotype (LES 0028772). **A.** Pereopod 1, arrow illustrating the dactylus in rostral view. **B.** Pereopod 7. **C.** Genital papilla and pleopod 1. **D.** Pleopod 2. **E.** Pleopod 3 exopod. **F.** Pleopod 4 exopod. **G.** Pleopod 5 exopod. Figure not to scale.

PEREPOD 1. Elongated, carpus short, transverse antennal grooming brush; dactylus with inner claw not surpassing median portion of outer claw, unguis seta, and dactylar organ simple not surpassing distal margin of outer claw (Fig. 7A).

PLEOPOD EXOPODS. Pleopod 1 and 2 exopods with respiratory areas.

Male

PEREPOD 1. Without particular modifications (Fig. 7A).

PEREPOD 7. Basis and ischium elongated, sternal margin straight, with one semicircular lobe on distal portion in rostral view; merus bearing semicircular lobe in proximal portion in rostral view (Fig. 7B).

GENITAL PAPILLA. Triangular ventral shield and subapical orifices (Fig. 7C).

PLEOPODS. Pleopod 1 (Fig. 7C) exopod ovoid, with semicircular apex; endopod about twice as long as exopod and bent outwards, apex acute and bearing small setae along inner margin. Pleopod 2 (Fig. 7D) exopod triangular, outer margin concave with one seta; endopod flagelliform, longer than exopod. Pleopod 3–4 exopods as in Fig. 7E–F, respectively. Pleopod 5 (Fig. 7G) exopod triangular, outer margin almost straight with one tiny seta, inner margin covered with small setae, apex short and acute.

Taxonomic remarks

Circoniscus xikrin sp. nov. is clearly distinguishable from *C. buckupi*, *C. carajasensis*, and *C. mendesi* sp. nov. in the semicircular lobe on the rostral surface of the male pereopod 7 ischium and merus, and telson with acute apex.

Ecological remarks

The specimens were collected in the aphotic zone, inhabiting very moist soil. This species is considered to be a troglobitic due to the absence of body pigments and eyes, and elongated pereopods.

Distribution

Presently known only from SB_0049 cave in PARNA dos Campos Ferruginosos, Canaã dos Carajás, and N3_0026 cave system, Parauapebas, Pará, Brazil (Fig. 1).

Family Armadillidae Brandt, 1831

Genus *Ctenorillo* Verhoeff, 1942

Type species

Ctenorillo buddelundi Verhoeff, 1942, by monotypy [= *Cubaris regulus* 893 Van Name, 1920], by monotypy (see Schmidt & Leistikow 2004).

Ctenorillo ferrarai Campos-Filho, Araujo & Taiti, 2014
Figs 8B–D

Ctenorillo ferrarai Campos-Filho, Araujo & Taiti in Campos-Filho *et al.*, 2014: 412, figs 37–40, table 1.

Ctenorillo ferrarai – Fernandes *et al.* 2019: 1119. — Campos-Filho *et al.* 2023: 560, figs 22–23.

Material examined

BRAZIL – Pará, Canaã dos Carajás • 1 ♂, 1 ♀; SB_0049 caves (also registered as GEM 1460 and known as Samuel II by local people); 6°18'58.4" S, 49°53'41.1" W; 629 m a.s.l.; 16 Feb. 2022; J.E. Gallão,

J.S. Gallo, D.F. Torres and V.F. Sperandei leg.; LES 0028774. – **Parauapebas** • 1 ♂; CAV-N1-0174; 6°01'27.20" S, 50°17'54.73" W; 649 m a.s.l.; 12 Feb. 2022; J.E. Gallão, J.S. Gallo, D.F. Torres and V.F. Sperandei leg.; LES 0028775 • 3 ♂♂, 1 ♀; same collection data as for preceding; LES 0028776 • 2 ♂♂, 1 ♀; same collection data as for preceding; LES 0028777.

Ecological remarks

The specimens were collected in the aphotic and twilight zones, living under rocks and among roots in moist soil (Fig. 8B–D).

Distribution

Ctenorillo ferrarai has been recorded in caves of the FLONA de Carajás, in the municipalities of Canaã dos Carajás and Curionópolis, Pará (Campos-Filho *et al.* 2014, 2023). This is the first record in Parauapebas (Fig. 1).

Discussion

As mentioned, the genus *Circoniscus* has a distribution restricted to South America, with the majority of records in the Brazilian territory. Campos-Filho *et al.* (2011) described *Circoniscus buckupi* and *C. carajasensis*, and Ferreira *et al.* (2015) recorded *Circoniscus* sp., all from iron ore caves in FLONA de Carajás. Campos-Filho *et al.* (2014) recorded the epigeal species *C. bezzi* Arcangeli, 1931, *C. intermedius*




Fig. 8. **A.** *Circoniscus mendesi* López-Orozco, Campos-Filho & Bichuette sp. nov., specimen from CAV-N1-0174. **B.** Parauapebas, Pará, in moist soil. **C.** *Ctenorillo ferrarai* Campos-Filho, Araujo & Taiti, 2014, specimen from CAV-SB-0049 (also known as Samuel II by local people). **D.** Canaã dos Carajás, Pará, in moist soil.

Souza & Lemos de Castro, 1991, and *C. incisus* Souza & Lemos de Castro, 1991 from the states Pará, Mato Grosso do Sul, and Minas Gerais, all in Brazilian caves. Moreover, Campos-Filho *et al.* (2019, 2023) described several epigeal species from several karstic areas in Brazil. Taking into account the above, it is clear that the subterranean habitat plays an important role in the conservation of the Oniscidea.

According to the CECAV/ICMbio database (CECAV 2020), Brazil has more than 22 000 caves, which hold a high taxonomic diversity (Gnaspini & Trajano 1994; Pinto-da-Rocha 1995; Gallão & Bichuette 2015, 2018; Trajano *et al.* 2016; Chagas-Jr & Bichuette 2018; Bichuette & Gallão 2021). Thus, our knowledge of the diversity of the terrestrial isopods from Brazilian caves is still far from complete. As mentioned by Campos-Filho *et al.* (2014), the taxonomic impediment is the primary reason for the delay in the assessment of the biodiversity of Oniscidea, both outside and inside caves. Moreover, it delays the progress in the knowledge of various other groups of invertebrates in subterranean ecosystems. Considering the national heritage of scientific collections, allied with works in development, it is important to emphasize the need to encourage the formation of more taxonomists to process the amount of material redeposited in these collections. In addition, it is worth mentioning that the iron ore caves from the Parauapebas region and surroundings suffer from iron mining for exportation. The FLONA de Carajás holds the largest mining project in Brazil (Fig. 1C). The impact of mining may be the worst to subterranean habitats due to the irreversible destruction, obliterating all the subterranean fauna. Knowing and describing our subterranean diversity are the first steps toward the conservation of our biological subterranean heritage.

The present work describes two new species of terrestrial isopods, *C. mendesi* sp. nov. and *C. xikrin* sp. nov., which belong to the troglobitic inventory of the iron ore cave diversity within the Amazonian domain. Moreover, it expands the knowledge of the distribution of *B. tarzan* and *C. ferrarai*. The diversity of terrestrial isopods associated with subterranean habitats from Brazil is increased to 80 species. Lastly, this information might be used in management and conservation plans both for this group as well as for Brazilian subterranean ecosystems.

Acknowledgements

The research was developed under the TCCE (Termo de Compromisso de Compensação Espeleológica) ICMBio (Instituto Chico Mendes de Conservação da Biodiversidade)/Vale S.A. Nº. TCCE 01/2018, with operational management carried out by the IABS (Instituto Brasileiro de Desenvolvimento e Sustentabilidade). We also thank the staff of IABS for all support and logistics management. We are also in debt to the ICMBio team José S. Moraes (Pezão), Airton d.S. Almeida, Deyvid R. Ramos, Hélio, Silvana Araujo, Paulo J.B. Faiad, and Wendelo S. Costa for helping us with logistics and fieldwork; to LES team (UFSCar) for help in the organization of scientific collection, especially to M.F.C. Zancheta and E.L. da Silva; to Odete Rocha for the loan of her laboratory facilities for the development of illustrations; this study was financed in part by the Brazilian funding agencies CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil) Finance Code 001, for a scholarship to CML-O and RB-A and CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico) for the scholarship to YMC-D; to Research Project titled “Biodiversity of terrestrial isopods (Crustacea, Isopoda, Oniscidea) from Cyprus in the light of integrative taxonomy”, “ONISILOS Research Program – 2018”, funded by the University of Cyprus, for the postdoctoral fellowship granted to ISC-F; to Vicerrectoría de Investigaciones de the University of Cartagena for the financial support to the Grupo de Investigación en Biología Descriptiva y Aplicada (Project Semilleros, Acta de Compromiso Nº 053–2022).

References

Anderson L.E. 1954. Hoyer's solution as a rapid permanent mounting medium for Bryophytes. *The Bryologist* 57: e242. <https://doi.org/10.2307/3240091>

- Bedek J., Taiti S. & Gottstein S. 2011. Catalogue and atlas of cave-dwelling terrestrial isopods (Crustacea: Oniscidea) from Croatia. *Natura Croatica* 20 (2): 237–354. <https://hrcak.srce.hr/75047>
- Bichuette M.E. & Gallão J.E. 2021. Under the surface: what we know about the threats to subterranean fishes in Brazil. *Neotropical Ichthyology* 19 (3): e210089. <https://doi.org/10.1590/1982-0224-2021-0089>
- Campos J. & Castilho A.F. 2012. Uma visão geográfica da Região de Carajás. In: Martins F.D., Castilho A.F., Campos J., Hatano F.M. & Rolim S.G. (orgs) *Fauna da Floresta de Carajás: Estudos sobre Vertebrados Terrestres*: 32–63. Nitro Imagens, São Paulo.
- Campos-Filho I.S. & Araujo P.B. 2011. Two new troglobitic species of Scleropactidae (Crustacea: Isopoda: Oniscidea) from Pará, Brazil. *Nauplius* 19: 27–39. <https://doi.org/10.1590/S0104-64972011000100004>
- Campos-Filho I.S., Araujo P.B., Bichuette M.E., Trajano E. & Taiti S. 2014. Terrestrial isopods (Crustacea: Isopoda: Oniscidea) from Brazilian caves. *Zoological Journal of the Linnean Society* 172 (2): 360–425. <https://doi.org/10.1111/zoj.12172>
- Campos-Filho I.S., Montesanto G., Araujo P.B. & Taiti S. 2017. New species and new records of terrestrial isopods (Crustacea, Isopoda, Oniscidea) from Brazil. *Iheringia, Série Zoologia* 107: e2017034. <https://doi.org/10.1590/1678-4766e2017034>
- Campos-Filho I.S., Cardoso G.M. & Aguiar J.O. 2018. Catalogue of terrestrial isopods (Crustacea, Isopoda, Oniscidea) from Brazil: an update with some considerations. *Nauplius* 26: e2018038. <https://doi.org/10.1590/2358-2936e2018038>
- Campos-Filho I.S., Fernandes C.S., Cardoso G.M., Bichuette M.E., Aguiar J.O. & Taiti S. 2019. Two new species and new records of terrestrial isopods (Crustacea, Isopoda, Oniscidea) from Brazilian caves. *Zootaxa* 4564 (2): 422–448. <https://doi.org/10.11646/zootaxa.4564.2.6>
- Campos-Filho I.S., Fernandes C.S., Cardoso G.M., Bichuette M.E., Aguiar J.O. & Taiti S. 2020. New species and new records of terrestrial isopods (Crustacea, Isopoda, Oniscidea) of the families Philosciidae and Scleropactidae from Brazilian caves. *European Journal of Taxonomy* 606: 1–38. <https://doi.org/10.5852/ejt.2020.606>
- Campos-Filho I.S., Gallo J.S., Gallão J.E., Torres D.F., Carpio-Díaz Y.M., López-Orozco C.M., Borja-Arrieta R., Taiti S. & Bichuette M.E. 2022a. Expanding the knowledge on the diversity of the cavernicolous Styloniscidae Vandel, 1952 (Oniscidea, Synocheta) from Brazil, with descriptions of two new species from the semiarid karst regions. *ZooKeys* 1101: 35–55. <https://doi.org/10.3897/zookeys.1101.79043>
- Campos-Filho I.S., Gallo J.S., Gallão J.E., Torres D.F., Horta L., Carpio-Díaz Y.M., López-Orozco C.M., Borja-Arrieta R., Aguiar J.O. & Bichuette M.E. 2022b. Unique and fragile diversity emerges from Brazilian caves – two new amphibious species of *Xangoniscus* Campos-Filho, Araujo & Taiti, 2014 (Oniscidea, Styloniscidae) from Serra do Ramalho karst area, state of Bahia, Brazil. *Subterranean Biology* 42: 1–22. <https://doi.org/10.3897/subtbiol.42.75725>
- Campos-Filho I.S., Cardoso G.M. & Bichuette M.E. 2022c. Isopoda, Oniscidea. In: Zampaulo R.A. & Prous X. (eds) *Fauna Cavernícola do Brasil*: 363–387. Editora Rupestre, Belo Horizonte, Brazil.
- Campos-Filho I.S., Sfenthourakis S., Gallo J.S., Gallão J.E., Torres D.F., Chagas-Jr A., Horta L., Carpio-Díaz Y.M., López-Orozco C.M., Borja-Arrieta R., Araujo P.B. & Bichuette M.E. 2023. Shedding light into Brazilian subterranean isopods (Isopoda, Oniscidea): expanding distribution data and describing new taxa. *Zoosystema* 45 (19): 531–599. <https://doi.org/10.5252/zoosystema2023v45a19>
- Cardoso G.M. & Ferreira R.L. 2023. New troglobitic species of *Benthana* Budde-Lund, 1908 and *Benthanoides* Lemos de Castro, 1958 from iron-ore caves and their important record in the Amazon biome (Crustacea: Isopoda: Philosciidae). *Zootaxa* 4 (27): 548–562, <https://doi.org/10.11646/zootaxa.5319.4.5>

- Cardoso G.M., Bastos-Pereira R., Souza L.A. & Ferreira R.L. 2020a. New troglotic species of *Xangoniscus* (Isopoda: Styloniscidae) from Brazil, with notes on their habitats and threats. *Zootaxa* 4819 (1): 84–108. <https://doi.org/10.11646/zootaxa.4819.1.4>
- Cardoso G.M., Bastos-Pereira R., Souza L.A. & Ferreira R.L. 2020b. New cave species of *Pectenoniscus* Andersson, 1960 (Isopoda: Oniscidea: Styloniscidae) and an identification key for the genus. *Nauplius* 28: e2020039. <https://doi.org/10.1590/2358-2936e2020039>
- Cardoso G.M., Bastos-Pereira R., Souza L.A. & Ferreira R.L. 2021. *Chaimowiczia*: a new Iuiuniscinae genus from Brazil (Oniscidea, Synocheta, Styloniscidae) with the description of two new troglotic species. *Subterranean Biology* 39: 45–62. <https://doi.org/10.3897/subtbiol.39.65305>
- Cardoso G.M. & Ferreira R.L. 2023. New troglotic species of *Benthana* Budde-Lund, 1908 and *Benthanoides* Lemos de Castro, 1958 from iron-ore caves and their important record in the Amazon biome (Crustacea: Isopoda: Philosciidae). *Zootaxa* 4 (27): 548–562. <https://doi.org/10.11646/ZOOTAXA.5319.4.5>
- Carmo F.F., Carmo F.F., Campos I.C. & Jacobi C.M. 2012. Cangas: ilhas de ferro estratégicas para a conservação. *Ciência Hoje* 295 (50): 48–53.
- CECAV – Centro Nacional de Pesquisas e Conservação de Cavernas. 2020. Cadastro Nacional de Informações Espeleológicas – CANIE. Available from <https://www.gov.br/icmbio/pt-br/assuntos/centros-de-pesquisa/cecav/cadastro-nacional-de-informacoes-espeleologicas> [accessed 6 Jan. 2021].
- Chagas-Jr A. & Bichuette M.E. 2018. A synopsis of centipedes in Brazilian caves: hidden species diversity that needs conservation (Myriapoda, Chilopoda). *ZooKeys* 737: 13–56. <https://doi.org/10.3897/zookeys.737.20307>
- Dimitriou A.C., Taiti S. & Sfenthourakis S. 2019. Genetic evidence against monophyly of Oniscidea implies a need to revise scenarios for the origin of terrestrial isopods. *Scientific Reports* 9: e18508. <https://doi.org/10.1038/s41598-019-55071-4>
- Fernandes C.S., Campos-Filho I.S., Araujo P.B. & Bichuette M.E. 2019. Synopsis of terrestrial isopods (Crustacea: Isopoda: Oniscidea) from Brazilian caves, with emphasis on new records from north, midwest, northeast and southeast regions. *Journal of Natural History* 53 (17–18): 1095–1129. <https://doi.org/10.1080/00222933.2019.1634225>
- Ferreira R.L., Oliveira M.P.A. & Silva M.S. 2015. Biodiversidade subterrânea em geossistemas ferruginosos. In: Carmo F.F. & Kamino L.H.Y. (eds) *Geossistemas Ferruginosos no Brasil*: 195–231. Instituto Prístino, Belo Horizonte, Brazil.
- Gallão J.E. & Bichuette M.E. 2015. Taxonomic distinctness and conservation of a new high biodiversity subterranean area in Brazil. *Anais da Academia Brasileira de Ciências* 87 (1): 209–217. <https://doi.org/10.1590/0001-3765201520140312>
- Gallão J.E. & Bichuette M.E. 2018. Brazilian obligatory subterranean fauna and threats to the hypogean environment. *ZooKeys* 746: 1–23. <https://doi.org/10.3897/zookeys.746.15140>
- Gnaspini P. & Trajano E. 1994. Brazilian cave invertebrates, with a checklist of troglomorphic taxa. *Revista Brasileira de Entomologia* 38 (4): 549–584.
- Grangeiro D.C. & Christoffersen M.L. 2010. A new species of *Androdeloscia* (Isopoda: Oniscidea) from Brazilian Amazon. *Revista Nordestina de Biologia* 10: 77–93.
- Grangeiro D.C., Souza L.A. & Christoffersen M.L. 2017. New species of *Xiphoniscus* and new record of *Androdeloscia escalonai* (Isopoda, Scutocoxifera, Oniscidea, Philosciidae) from Brazilian Amazon. *Zootaxa* 4350 (2): 374–384. <https://doi.org/10.11646/zootaxa.4350.2.11>

- Grangeiro D.C., Souza L.A. & Christoffersen M.L. 2021. Three new species of *Androdeloscia* Leistikow, 1999 (Oniscidea: Philosciidae) from the Brazilian Amazonia. *Zootaxa* 5047 (3): 321–341. <https://doi.org/10.11646/zootaxa.5047.3.5>
- ICMBio – Instituto Chico Mendes de Conservação da Biodiversidade. 2020. Unidades de Conservação – Amazônia. Floresta Nacional de Carajás. Available from <https://www.icmbio.gov.br/portal/unidadesdeconservacao/biomasbrasileiros/amazonia/unidades-de-conservacao-amazonia/1927-flona-de-carajas> [accessed 7 Feb. 2020].
- Javidkar M., Cooper S.J.B., King R.A., Humphreys W.F. & Austin A. 2015. Molecular phylogenetic analyses reveal a new southern hemisphere oniscidean family (Crustacea: Isopoda) with a unique water transport system. *Invertebrate Systematics* 29: 554–577. <https://doi.org/10.1071/IS15010>
- Köppen W. 1948. *Climatología: con un Estudio de los Climas de la Tierra*. Fondo de 1992 Cultura Economica, Pánuco, Veracruz, Mexico.
- Kottek M., Grieser J., Beck C., Rudolf B. & Rubel F. 2006. World Map of the Köppen-Geiger climate classification updated. *Meteorologische Zeitschrift* 15: 259–263. <https://doi.org/10.1127/0941-2948/2006/0130>
- Leistikow A. 1999. *Androdeloscia* gen. n., a new genus of South American terrestrial isopods with description of 13 new species (Crustacea: Oniscidea: “Philosciidae”). *Revue suisse de Zoologie* 106: 813–904. Available from <https://biodiversitylibrary.org/page/41279890> [accessed 7 Feb. 2020].
- Leistikow A. 2001. Phylogeny and biogeography of South American Crinocheta, traditionally placed in the family “Philosciidae” (Crustacea: Isopoda: Oniscidea). *Organisms, Diversity & Evolution* Electronic Supplement 4: 1–85. Available from <http://www.senckenberg.uni-frankfurt.de/odes/01-04.htm> [accessed 6 Feb. 2020].
- Lemos de Castro, A. 1984. Uma nova espécie de *Prosekia* (Philosciidae, Isopoda) de uma floresta inundável (Igapó) na Amazonia Central. *Amazoniana* 8 (4): 441–445.
- Montesanto G. 2015. A fast GNU method to draw accurate scientific illustrations for taxonomy. *ZooKeys* 515: 191–206. <https://doi.org/10.3897/zookeys.515.9459>
- Montesanto G. 2016. Drawing setae: a GNU way for digital scientific illustrations. *Nauplius* 24: e2016017. <https://doi.org/10.1590/2358-2936e2016017>
- Mota N.F.O., Silva L.V.C., Martins F.D. & Viana P.L. 2015. Vegetação sobre sistemas ferruginosos da Serra dos Carajás. In: Carmo F.F. & Kamino L.H.Y. (eds) *Geossistemas Ferruginosos do Brasil*: 289–315. Instituto Prístino, Belo Horizonte, Brazil.
- Ocampo-Maceda A.T., Ruelas-Cabana C.M., López-Orozco C.M. & López-Tejeda E.L. 2022. Catalogue of terrestrial isopods (Isopoda, Oniscidea) from Peru, with new records of *Circoniscus ornatus* (Scleropactidae) and *Ethelum americanum* (Eubelidae). *Nauplius* 30: e2022003. <https://doi.org/10.1590/2358-2936e2022003>
- Piló L.B., Coelho A. & Reino J.C.R. 2015. Geoespeleologia em rochas ferríferas: cenário atual e conservação. In: Carmo F.F. & Kamino L.H.Y. (eds) *Geossistemas Ferruginosos do Brasil*: 125–148. Instituto Prístino, Belo Horizonte, Brazil.
- Pinto-da-Rocha, R. 1995. Sinopse da fauna cavernícola do Brasil (1907–1994). *Papéis Avulsos de Zoologia* 39 (6): 61–173.
- Reboleira A.S.P.S., Goncalves F., Oromí P. & Taiti S. 2015. The cavernicolous Oniscidea (Crustacea: Isopoda) of Portugal. *European Journal of Taxonomy* 161: 1–61. <https://doi.org/10.5852/ejt.2015.161>

- Schmalzfuss H. 2003. World catalog of terrestrial isopods (Isopoda: Oniscidea). *Stuttgarter Beiträge zur Naturkunde, Serie A* 654: 1–341.
- Schmidt C. 2007. Revision of the Neotropical Scleropactidae (Crustacea: Oniscidea). *Zoological Journal of the Linnean Society* 151: 1–339. <https://doi.org/10.1111/j.1096-3642.2007.00286.x>
- Schmidt C. & Leistikow A. 2004. Catalogue of genera of the terrestrial Isopoda (Crustacea: Isopoda: Oniscidea). *Steenstrupia* 28 (1): 1–118.
- Schmidt C. & Leistikow A. 2005. Review of the genus *Androdeloscia* Leistikow, with description of four new species (Crustacea, Isopoda, Oniscidea). *Entomologische Abhandlungen* 62: 117–163.
- Sfenthourakis S. & Taiti S. 2015. Patterns of taxonomic diversity among terrestrial isopods. *ZooKeys* 515: 13–25. <https://doi.org/10.3897/zookeys.515.9332>
- Souza L.A., Bezerra A.V. & Araujo J.P. 2006. The first troglobitic species of Scleropactidae from Brazil (Crustacea, Isopoda, Oniscidea). *Subterranean Biology* 4: 37–43.
- Tabacaru I. & Giurginca, A. 2013. Cavernicolous Oniscidea of Romania. *Travaux de l'Institut de Spéologie "Emile Racovitza"* 42: 3–26.
- Taiti S. 2004. Crustacea: Isopoda: Oniscidea (woodlice). In: Gunn J. (ed.) *Encyclopedia of Caves and Karst Science*: 547–551. Fitzroy Dearborn, Taylor and Francis Group, New York.
- Taiti S. & Gruber G.A. 2008. Cave-dwelling terrestrial isopods from Southern China (Crustacea, Isopoda, Oniscidea), with descriptions of four new species. In: Latella L. & Zorzin R. (eds) *Research in South China Karsts*: 101–123. Memorie del Museo Civico di Storia Naturale di Verona, Monografie Naturalistiche.
- Taiti S. & Xue Z. 2012. The cavernicolous genus *Trogloniscus* nomen novum, with descriptions of four new species from southern China (Crustacea, Oniscidea, Styroniscidae). *Tropical Zoology* 25 (4): 183–209. <https://doi.org/10.1080/03946975.2012.751240>
- Trajano E., Gallão J.E. & Bichuette M.E. 2016. Spots of high diversity of troglobites in Brazil: the challenge of measuring subterranean diversity. *Biodiversity and Conservation* 25: 1805–1828. <https://doi.org/10.1007/s10531-016-1151-5>
- Vandel A. 1973. Les isopodes terrestres et cavernicoles de l'île de Cuba. In: Orghidan T., Núñez A., Botosaneanu L., Decou V., Negrea Ş. & Viña N. (eds) *Résultats des Expéditions biospéologiques cubano-roumaines à Cuba, Vol. I*: 153–188. Academiei Republicii Socialiste România, Bucharest.

Manuscript received: 7 June 2023

Manuscript accepted: 25 September 2023

Published on: 6 February 2024

Topic editor: Magalie Castelin

Desk editor: Pepe Fernández

Printed versions of all papers are also deposited in the libraries of the institutes that are members of the *EJT* consortium: Muséum national d'histoire naturelle, Paris, France; Meise Botanic Garden, Belgium; Royal Museum for Central Africa, Tervuren, Belgium; Royal Belgian Institute of Natural Sciences, Brussels, Belgium; Natural History Museum of Denmark, Copenhagen, Denmark; Naturalis Biodiversity Center, Leiden, the Netherlands; Museo Nacional de Ciencias Naturales-CSIC, Madrid, Spain; Leibniz Institute for the Analysis of Biodiversity Change, Bonn – Hamburg, Germany; National Museum of the Czech Republic, Prague, Czech Republic.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [European Journal of Taxonomy](#)

Jahr/Year: 2024

Band/Volume: [0921](#)

Autor(en)/Author(s): Lopez-Orozco Carlos Mario, Campos-Filho Ivanklin Soares, Gallo Jessica S., Gallao Jonas E., Carpio-Diaz Yesenia M., Borja-Arrieta Ricardo, Bichuette Maria Elina

Artikel/Article: [Iron-isopods: new records and new species of terrestrial isopods \(Isopoda, Oniscidea\) from Brazilian Amazon iron ore caves 116-135](#)