

European Journal of Taxonomy 930: 157–181 https://doi.org/10.5852/ejt.2024.930.2491

This work is licensed under a Creative Commons Attribution License (CC BY 4.0).

Research article

urn:lsid:zoobank.org:pub:8FAF9EED-F707-4407-9885-2E571DB89F32

Taxonomic account of the ant genus *Syscia* Roger, 1816 (Hymenoptera, Formicidae) from Asia, with descriptions of two new species from China

Congcong DU⁰, Zhenghui XU⁰ & Zhilin CHEN⁰,*

 ^{1,3}Key Laboratory of Ecology of Rare and Endangered Species and Environmental Protection (Guangxi Normal University), Ministry of Education, Guilin 541004, China.
²Key Laboratory of Forest Resources Conservation and Utilization in the Southwest Mountains of China Ministry of Education, Southwest Forestry University, Kunming, Yunnan Province 650224, China.
^{1,3}Guangxi Key Laboratory of Rare and Endangered Animal Ecology, Guangxi Normal University, Guilin 541004, China.
²Key Laboratory of Forest Disaster Warning and Control in Yunnan Province, College of Biodiversity Conservation, Southwest Forestry University, Kunming, Yunnan Province 650224, China.

> *Corresponding author: chenzhilin35@163.com ¹Email: cleverduwang@gmail.com ²Email: xuzhenghui1962@163.com

¹urn:lsid:zoobank.org:author:630068B0-6A0D-42CA-B084-3D638B72D83E ²urn:lsid:zoobank.org:author:1A507854-7B05-4BB4-BDB9-CC62BC97AE66 ³urn:lsid:zoobank.org:author:F148C223-F388-4397-8E8C-2FAAE42157A6

Abstract. The ant genus *Syscia* Roger, 1861 is a cryptobiotic group of the subfamily Dorylinae. In this study, eight species of this genus from Asia are recognized, including two new species, *Syscia arcodorsa* sp. nov. and *S. zhoui* sp. nov., both described from Guangxi in China. In addition, *Cerapachys guizhouensis* Zhou, 2006 is redescribed and transferred to *Syscia* as *S. guizhouensis* (Zhou, 2006) comb. nov. The species *Syscia typhla* Roger, 1861 is removed from the ant fauna of China. Lastly, a key to species of *Syscia* from Asia based on the worker caste is presented.

Keywords. Dorylinae, redescription, *Syscia arcodorsa* sp. nov., *Syscia zhoui* sp. nov., *Cerapachys guizhouensis* Zhou, 2006.

Du C., Xu Z. & Chen Z. 2024. Taxonomic account of the ant genus *Syscia* Roger, 1816 (Hymenoptera, Formicidae) from Asia, with descriptions of two new species from China. *European Journal of Taxonomy* 930: 157–181. https://doi.org/10.5852/ejt.2024.930.2491

Introduction

The ant genus Syscia Roger, 1861 (Hymenoptera, Formicidae, Dorylinae) was first proposed by Roger (1861) in a paper on 'Ponera-like ants' based on the type species Syscia typhla Roger, 1861, and was regarded as a member of the tribe "Cerapachysii". Since then, the taxonomic status of this genus has changed many times by different myrmecologists, e.g., considered as a valid genus (Forel 1900; Dalla Torre 1903; Bingham 1903), as a subgenus of Cerapachys Smith, 1857 (Wheeler 1902, 1910, 1922; Emery 1902, 1911; Forel 1917; Donisthorpe 1943) or a junior synonym of *Cerapachys* (Kempf 1972; Brown 1975). More recently, Brady et al. (2014) found Cerapachys to be non-monophyletic. Additionally, according to phylogenetic inference based on molecular data for the major dorylomorph lineages, they found that Neotropical and Indomalayan species of Syscia together formed a clade distinct from other genera. Later, Borowiec (2016) systematically revised the generic classification of the subfamily Dorylinae Leach, 1815 based on molecular phylogenetic evidence and a critical reappraisal of doryline morphology. The result of their analysis divided the former Cerapachys into 9 different monophyletic genera and recognized Syscia as a valid genus again. Meanwhile, they comprehensively described the specific characteristics of different types of Syscia, and diagnosed their unique autapomorphic characteristics: basal segment of hind tarsus widening distally with a light patch of cuticle on the inner (flexor) side, and abdominal tergite IV anteriorly folding over sternite. That outstanding work provided a solid foundation for the subsequent classification of the genus Syscia.

Syscia is the only doryline genus with a disjunct distribution between the Old and New World. Members of this genus are usually found in leaf litter, rotting wood and soil cores, and are thus considered part of the cryptobiotic fauna (Borowiec 2016; Jaitrong et al. 2020; Longino & Branstetter 2021). Previously, only five species had been described worldwide (Borowiec 2016). Therefore, as Borowiec (2016) pointed out at the time, there might be more morphospecies presented in collections from the Old World and additional undescribed species in the New World. Since then, Jaitrong et al. (2020) described two new species from Thailand and Aswaj et al. (2021) described and illustrated one new species from India. Longino and Branstetter (2021) integrated ultra-conserved element (UCE) phylogenomics with traditional taxonomy to reveal and name 31 new species of Syscia in Central and South America, resulting in an increase in the number of known species of *Syscia* in the New World from 3 to 34. By contrast, the known species of Syscia in the Old World are still few, with only 5 recorded (Antweb 2022). Except for Syscia typhla, which is widespread, the remaining species of Syscia are generally native to their type-localities or surrounding country, e.g., S. chaladthanyakiji Jaitrong, Wiwatwitaya & Yamane, 2020 and S. reticularis Jaitrong Wiwatwitaya & Yamane, 2020 in Thailand, S. humicola Ogata, 1983 in Korea and Japan, and S. indica Aswaj, Sahanashree, Udayakantha, Aniruddha & Priyadarsanan, 2021 in India. The type-locality of Syscia typhla is Sri Lanka, and this species also was recorded in northern India (Ghosh et al. 2007), southern China (Xu 1998; Xu et al. 1999; Gu et al. 2019), and Japan (Sonobe 1973; Abe 1974) However, a detailed examination of relevant specimens of Xu (1998) and Xu et al. (1999), revealed that the specimens of Ooceraea biroi (Forel 1907) were misidentified as S. typhla. And the new species S. zhoui sp. nov. was misidentified as S. typhla by Gu et al. (2019). Therefore, there is currently no distribution of the Syscia in the ant fauna of China. Given the complexity and heterogeneity of the geography and climate in China, there are likely to be additional cryptic and undescribed species of Syscia to be discovered there.

In this paper, two new species are described, i.e., *Syscia arcodorsa* sp. nov. and *S. zhoui* sp. nov. *Cerapachys guizhouensis* Zhou, 2006 was combined in *Ooceraea* Roger, 1862 by Bharti *et al.* (2021). Here, we redescribe this species and revise it as a member of *Syscia*, *S. guizhouensis* (Zhou, 2006) comb. nov. A key to Asian species of *Syscia* based on the worker caste is provided.

Material and methods

The specimens of *Syscia arcodorsa* sp. nov. and *S. zhoui* sp. nov. were collected during field trips to Huaping village in the Huaping National Nature Reserve in Guilin City, and Pairu village in the Chongzuo White-Headed Langur National Reserve in Guangxi, China, by hand alive (Fig. 1), and later preserved in a vial containing absolute ethyl alcohol. Then the specimens were pin-mounted and examined with a Leica M205A stereo microscope. High-quality multifocused montage images were produced with the KEYENCE (VHX–6000) digital imaging system.

The examined specimens are deposited in the repositories with acronyms as follows: (1) GXNU (Insect Collection, Guangxi Normal University, Guilin, Guangxi, China); (2) SWFU (Insect Collection, Southwest Forestry University, Kunming, Yunnan Province, China); (3) IZCAS (Institute of Zoology, Chinese Academy of Sciences, Beijing, China). The type specimen images of *Syscia chaladthanyakiji*, *S. reticularis*, and *S. typhla* available on the AntWeb (http://www.antweb.org) were examined and compared. The general terminology of the *Syscia* worker ants follows Borowiec (2016) and Jaitrong *et al.* (2020). All measurements are given in millimeters.



Fig. 1. The sampled sites of *Syscia guizhouensis* comb. nov., *Syscia arcodorsa* sp. nov. and *Syscia zhoui* sp. nov. in China (Source: Esri, Maxar, GeoEye, Earthstar geographics, CNES/Airbus DS, USDA, USGS, AeroGRID, IGN, and the GIS User Community).

Abbreviations for	me	asurements and indices									
BL	=	body length, the total body outstretched length of the individual in lateral view, from the mandibular apex to the gastral apex, calculated by summing lengths of head, mesosoma, petiole, and gaster									
CI	=	cephalic index, $HW \times 100/HL$									
HL	=	head length, the maximum length of the head in full-face view, measured from a transverse line spanning the anteriormost points of the clypeus to that of posteriormost points of the cranium									
HW	=	head width, the maximum width of the cranium in full-face view (excluding eyes)									
ML	=	mesosomal length, maximum diagonal length of mesosoma in lateral view, measured from posterodorsal border of the pronotal flange to posterior basal angle of metapleuron									
MW	=	mesosomal width, maximum width of promesonotum in dorsal view									
SI	=	scape index, $SL \times 100/HW$									
SL	=	scape length, the maximum length of antennal scape excluding basal condular bulb									
РН	=	petiolar height, the maximum height of petiole in lateral view (including subpetiolar process)									
PI1	=	petiolar index 1, PL/PH×100									
PI2	=	petiolar index 2, PW/PL×100									
PL	=	petiolar length, the maximum length of petiole in lateral view (excluding helium)									
PPH	=	postpetiole height, maximum height in lateral view									
PPI1	=	postpetiole index 1, PPL/PPH × 100									
PPI2	=	postpetiole index 2, PPW/PPL×100									
PPL	=	= postpetiole length, maximum length in dorsal view									
PPW	=	postpetiole width, maximum width in dorsal view									
PW	=	petiolar width, maximum width of petiole in dorsal view									
WI	=	waist index, $PPW/PW \times 100$									

Institutional abbreviations

GXNU = Insect Collection, Guangxi Normal University, Guilin, Guangxi, China
IZCAS = Institute of Zoology, Chinese Academy of Sciences, Beijing, China
SWFU = Insect Collection, Southwest Forestry University, Kunming, Yunnan Province, China

Results

Taxonomy

Class Insecta Linnaeus, 1758 Order Hymenoptera Linnaeus, 1758 Family Formicidae Latreille, 1809 Subfamily Dorylinae Leach, 1815

Genus Syscia Roger, 1861

Key to Asian species of Syscia Roger, 1861 based on the worker caste

1.	Antennae with 9 segments (Figs 9A, 10D)	. 2
_	Antennae with 11 segments (Figs 2C, 3A, 4D, 5A, 7A)	. 3

DU C. et al., Taxonomic account of Syscia Roger, 1816 (Hymenoptera, Formicidae)

2.	Mesosoma rather elongate, long-ellipsoidal, with lateral sides and anterior margin obviously convex in dorsal view (Fig. 9C); both anterior margin and posterior margin of petiole convex in lateral view (Fig. 9D); posteroventral corner of subpetiolar process blunt rounded in lateral view (Fig. 9D); postpetiole in dorsal view longer than wide (Fig. 9C)
3. _	Head in full-face view oval (Figs 2A, 8B)
4.	Body entire reticulate, with deep bottoms (Fig. 8)
5.	Petiole in dorsal view broader than long (Fig. 3C); in lateral view the front and the back margins of the petiole vertical (Fig. 3A)
6. –	Subpetiolar process in lateral view subtriangular (Fig. 4C); anterior margin of petiole slightly convex (Fig. 4D)
7.	Propodeal declivity straight or slightly convex in lateral view (Figs 6–7)
-	Propodeal declivity slightly concave in lateral view (Fig. 5)

Syscia arcodorsa sp. nov.

urn: lsid: zoobank.org: act: 838 DEBA2-DA7B-473D-A7D3-475A3D3C1F09

Fig. 2, Table 1

Diagnosis

Body color brown with antennae and legs yellowish brown. Head in full-face view oval, distinctly longer than broad; antennae 11-segmented. Mesosoma in dorsal view subrectangular; in lateral view, dorsal outline distinctly convex. Petiole in dorsal view subrectangular, slightly longer than broad; in lateral view, petiolar node (excluding sternite) trapezoidal, slightly longer than high; subpetiolar process in lateral view subtriangle. Postpetiole in lateral view slightly higher than long, sternite anteroventrally produced as a blunt angle directed downward and forward.

Etymology

The specific epithet refers to the dorsal outline of the arched mesosoma in lateral view.



Fig. 2. *Syscia arcodorsa* sp. nov., holotype, worker (GXNU210848). **A**. Head in full-face view. **B**. Body in dorsal view. **C**. Body in lateral view (imaged by Zhilin Chen).

Type material

Holotype (worker)

CHINA•Guangxi, Guilin City, Longshen Town, Huaping Village in the Huaping National Nature Reserve; 25.626944° N, 109.183333° E; 503.50 m a.s.l.; 22 Jul. 2021; Zhilin Chen leg.; No. GXNU210848; GXNU: GXNU210848.

Paratypes (workers)

CHINA • 1 worker; same collection data as for holotype; GXNU: GXNU210848-1 • 1 worker; same collection data as for holotype; SWFU: GXNU210848-2 • 1 worker; same collection data as for holotype; IZCAS: GXNU210848-3.

Description

MEASUREMENTS AND INDICES. Holotype worker: BL 3.70, HL 0.63, HW 0.44, CI 70, SL 0.40, SI 91, MW 0.30, ML 0.70, PL 0.32, PH 0.30, PW 0.30, PI1 107, PI2 94, PPL 0.48, PPH 0.55, PPW 0.46, PPI1 86, PPI2 96, WI 152. Paratype workers: BL 3.61–3.73, HL 0.62–0.69, HW 0.43–0.49, CI 69–71, SL 0.36–0.43, SI 80–100, MW 0.33–0.35, ML 0.75–0.83, PL 0.27–0.33, PH 0.25–0.33, PW 0.25–0.33, PI1 100–108, PI2 96–100, PPL 0.44–0.45, PPH 0.42–0.45, PPW 0.40–0.50, PPI1 97–108, PPI2 89–100.00, WI 133–175 (n = 3) (individual measurements see Table 1).

Worker

HEAD. Antennae with 11 segments; apical antennal segment (XI) conspicuously enlarged, much broader than and longer than two preceding segments (IX–X) combined; scape short, not reaching mid-length of head when folded back (Fig. 2A). Head in full-face view oval, distinctly longer than broad, lateral margins convex, middle part of posterior margin slightly concave, with posterolateral corners rounded obviously (Fig. 2A). Frontal carinae short and narrow, not reaching to middle of head, with anterior half sharply elevated and posterior half low and vestigial in profile. Clypeus short, anterior margin almost straight. Eyes and ocelli completely absent (Fig. 2A). Mandibles triangular; masticatory margin serrated, without distinct teeth.

MESOSOMA. In dorsal view, rather robust, subrectangular, with lateral margins slightly convex, anterior margin convex, and posterior margin slightly concave (Fig. 2B); in lateral view, dorsal outline strongly convex; promesonotal suture and metanotal groove absent; mesopleuron demarcated from pronotum by distinct suture, but not demarcated from mesonotum and metapleuron; mesopleuron also not differentiated from metapleuron (Fig. 2C). Propodeal declivity feebly convex, encircled with distinct thin rim (Fig. 2C).

METASOMA. In dorsal view, petiole subrectangular, slightly longer than broad, lateral margins slightly convex (Fig. 2B); in lateral view, petiolar node (excluding sternite) trapezoidal with slightly convex dorsal outline, slightly longer than high, with bilateral edge sloped slightly (Fig. 2C). Subpetiolar process in lateral view subtriangle, with ventral outline concave (Fig. 2C). Postpetiole in dorsal view clearly larger than petiole, subtrapezoid, anterior margin slightly concave and shorter than posterior margin, lateral margins apparently convex (Fig. 2B); in lateral view, postpetiole slightly higher than long; dorsal outline slightly convex; postpetiolar sternite in lateral view low, with ventral margin slightly convex, anteroventrally produced as blunt angle directed downward and forward (Fig. 2C). Abdominal tergite IV (second gastral tergite) in dorsal view elongate, subrectangular with lateral margins convex, anterior margin obviously concave, and lateral margins slightly convex (Fig. 2B).

SCULPTURE. Entire body with numerous, relatively small, closely-spaced foveolae (Fig. 2). Antennal scape, outer surface of mandibles, and legs with fine dense micropunctures (Fig. 2A–C).

	Syscia arcodorsa sp. nov.				Syscie	<i>Syscia zhoui</i> sp. nov.				Syscia guizhouensis comb. nov.			
ABBV	holotype	paratypes		holotype	paratypes		holotype	paratypes					
BL	3.70	3.61	3.73	3.75	3.55	3.98	3.75	3.80	3.20	3.10	3.25	3.20	
HL	0.63	0.65	0.62	0.69	0.65	0.69	0.65	0.62	0.60	0.61	0.625	0.60	
HW	0.44	0.45	0.43	0.49	0.53	0.56	0.55	0.50	0.45	0.43	0.45	0.43	
CI	70	69	68	71	82	81	85	77	75	70	72	71	
SL	0.40	0.36	0.43	0.43	0.42	0.43	0.43	0.43	0.31	0.30	0.33	0.32	
SI	91	80	100	89	79	76	78	85	69	69	72	75	
MW	0.30	0.33	0.35	0.33	0.38	0.31	0.39	0.38	0.33	0.35	0.33	0.34	
ML	0.7	0.75	0.83	0.79	0.60	0.65	0.72	0.80	0.85	0.81	0.84	0.83	
PL	0.32	0.27	0.33	0.30	0.30	0.30	0.30	0.33	0.35	0.33	0.325	0.34	
РН	0.30	0.25	0.33	0.29	0.30	0.30	0.25	0.30	0.32	0.325	0.32	0.33	
PW	0.30	0.25	0.33	0.30	0.30	0.35	0.25	0.33	0.27	0.26	0.28	0.26	
PI1	107	108	100	105	100	100	120	110	107	102	102	105	
PI2	94	96	100	100	100	117	83	98	77	79	85	75	
PPL	0.48	0.44	0.45	0.45	0.44	0.48	0.45	0.49	0.41	0.40	0.42	0.43	
РРН	0.55	0.45	0.42	0.45	0.45	0.48	0.43	0.44	0.45	0.42	0.43	0.43	
PPW	0.46	0.44	0.50	0.40	0.40	0.43	0.42	0.45	0.42	0.4	0.43	0.43	
PPI1	86	97	108	100	97	99	106	111	91	95	98	99	
PPI2	96	100	92	89	91	89	93	93	102	100	102	101	
WI	152	175	154	133	133	121	168	138	156	154	155	169	

Table 1. The morphometric data of *Syscia arcodorsa* sp. nov., *S. zhoui* sp. nov. and *S. guizhouensis* comb. nov.

PILOSITY. Body entirely covered with long and short decumbent, sub-erect, and erect hairs.

COLORATION. Body brown. Antennae and legs yellowish brown (Fig. 2).

Distribution

China (Guangxi).

Habitat

The nest of *Syscia arcodorsa* sp. nov. was found in the Huaping National Nature Reserve, Guangxi, China. The sampled site is located in the evergreen broad-leaf forest, in the soil surface under the falling leaves near a flowing mountain stream.

Recognition

Syscia arcodorsa sp. nov. is similar to *S. reticularis*, because both of them have oval heads in full-face view. But it can be distinguished from the latter by the body covered with punctures or shallow fovea; however, the latter body is entirely reticulate, with deep bottoms.

This new species is also similar to an Indian species, *S. indica*, in general appearance of the workers with 11-segmented antennae. However, *S. arcodorsa* sp. nov. can be distinguished from the latter by the following characteristics: 1) head in full-face view oval, with posterolateral corners of head rounded obviously in *S. arcodorsa* (but head in full-face view rectangular, with posterolateral corners

of head angular obtusely in *S. indica*); 2) dorsal outline of mesosoma in lateral view curved or arched in *S. arcodorsa* (but almost straight in *S. indica*); 3) declivity of propodeum in lateral view almost convex in *S. arcodorsa* (but slightly concave in *S. indica*).

Syscia chaladthanyakiji Jaitrong, Wiwatwitaya & Yamane, 2020 Fig. 3

Syscia chaladthanyakiji Jaitrong et al., 2020: 3, figs 1-6, 11 (w.q.) Thailand.

Diagnosis

Workers have 11-segmented antennae. Head in full-face view rectangular, longer than broad, with posterior margin straight. Mesosoma in dorsal view subrectangular with almost parallel lateral sides; in lateral view, dorsal outline is almost straight. Petiole in dorsal view rectangular, broader than long; in lateral view (excluding sternite) rectangular with convex dorsal outline, bilateral edge vertical; subpetiolar process in lateral view subtriangular with large acute tooth on anteroventral margin, with posterior outline strongly concave. Postpetiole in lateral view slightly higher than long, sternite anteroventrally produced as blunt angle directed downward and forward.

Material examined

Unexamined. But the holotype of *S. chaladthanyakiji* was examined from images available from AntWeb (https://www.antweb.org; THNHM-I-05483, imaged by Yudthana Samsung).

Syscia humicola (Ogata, 1983) Fig. 4

Cerapachys humicola Ogata, 1983: 132, figs 1-13 (w. ergatoid q. m.). Japan.

Syscia humicola – Borowiec 2016: 224.

Diagnosis

Workers have 11-segmented antennae. Head in full-face view subrectangular, longer than broad, with posterior margin slightly convex and posterolateral corners angular. Mesosoma in dorsal view subrectangular, with almost parallel lateral sides; in lateral view, dorsal outline slightly convex. Petiole in dorsal view nearly subquadrate, as broad as long, with lateral sides convex; in lateral view (excluding sternite) trapezoidal, with slightly convex dorsal outline, bilateral edge sloped slightly; subpetiolar process in lateral view subtriangular, with posterior outline slightly concave. Postpetiole in lateral view as broad as long, sternite anteroventrally produced as nearly orthogonal directed downward and forward.

Type material

Paratype (worker)

JAPAN • 1 worker; Mt. Shiratake, Tsushima Nagasaki Pref.; 14 Jul. 1981; K.Ogata leg.; Colony No.CE-0003; currently stored at the IZCAS.



Fig. 3. *Syscia chaladthanyakiji* Jaitrong, Wiwatwitaya & Yamane, 2020, holotype, worker (THNHM-I-05483). A. Head in full-face view. **B**. Label. **C**. Body in dorsal view. **D**. Body in lateral view (images cited from https://www.antweb.org; imaged by Yudthana Samung).



Fig. 4. *Syscia humicola* (Ogata, 1983), paratype, worker (IZCAS). **A**. Head in full-face view. **B**. Body in dorsal view. **C**. Body in lateral view (imaged by Zhilin Chen).

Syscia indica Aswaj, Sahanashree, Udayakantha, Aniruddha & Priyadarsanan, 2021 Fig. 5

Syscia indica Aswaj et al., 2021: 67, figs 1, 3a-f (w.). India (Arunachal Pradesh).

Diagnosis

Workers have 11-segmented antennae. Head in full-face view subrectangular, distinctly longer than broad; lateral margins slightly convex. Mesosoma in dorsal view elongates with almost parallel lateral sides; in lateral view, dorsal outline almost straight. Petiole in dorsal view subrectangular, distinctly longer than broad; lateral margins almost parallel-sided; in lateral view (excluding sternite) trapezoidal, slightly shorter than high, dorsal outline slightly convex, bilateral edge slope slightly; subpetiolar process in lateral view subrectangular, ventral outline strongly concave; ventrolateral surface of petiole with two distinct almost parallel carinae just above subpetiolar process. Postpetiole in lateral view slightly shorter than high, sternite anteroventrally produced as blunt angle directed downward and forward.

Type material

Unexamined. But holotype of S. indica was examined from images available from Aswaj et al. (2021).

Syscia guizhouensis (Zhou, 2006) comb. nov. Figs 6–7, Table 1

Cerapachys guizhouensis Zhou, 2006: 581, figs 1-2 (w.). China.

Ooceraea guizhouensis - Bharti et al. 2021: 181.

Diagnosis

Workers have 11-segmented antennae, body color brown with antennae and legs yellowish brown. Head in full-face view subrectangular, distinctly longer than broad. Mesosoma in dorsal view subrectangular with almost parallel lateral sides; in lateral view, dorsal outline slightly convex. Petiole in dorsal view elliptical, longer than broad, with lateral margins convex; in lateral view, petiolar node (excluding sternite) subtrapezoid with slightly convex dorsal outline, bilateral edge slop, and slightly concave; subpetiolar process in lateral view subtrapezoid, with posterior outline strongly concave, and ventral margin very short and almost straight. Postpetiole in lateral view slightly higher than long, sternite anteroventrally produced as nearly orthogonal directed downward and forward.

Type material

Holotype (worker)

CHINA • Guizhou, Tongren City, in the Fanjingshan National Nature Reserve; 27.911945° N, 108.696634° E; 2195.00 m a.s.l.; 3 Jun. 2002; Shanyi Zhou leg; No. GXNU 060127; GXNU: GXNU060127.

Paratypes (workers)

CHINA • 1 worker; same collection data as for holotype; GXNU: SWFU 060127-1 • 1 worker; same collection data as for holotype; SWFU: GXNU 060127-2 • 1 worker; same collection data as for holotype; IZCAS: GXNU 060127-3.

Other material examined

CHINA • 1 spec; Xizang, Linzhi City, Motuo Town, Motuo Village; 29.32° N, 95.34° E; 1230 m a.s.l. ; 27 Aug. 2016; Zhilin Chen leg.; No. G160363; GXNU: G160363.



Fig. 5. *Syscia indica* Aswaj, Sahanashree, Udayakantha, Aniruddha & Priyadarsanan, 2021, holotype, worker. **A**. Head in full-face view. **B**. Petiole and postpetiole in dorsal view. **C**. Body in dorsal view **D**. Body in lateral view (images cited from Aswaj *et al.* 2021).



Fig. 6. *Syscia guizhouensis* (Zhou, 2006) comb. nov., holotype, worker (No. G160363). A. Head in full-face view. **B**. Body in dorsal view. **C**. Body in lateral view (imaged by Zhilin Chen).

Redescription

MEASUREMENTS AND INDICES. Holotype worker: BL 3.20, HL 0.60, HW 0.45, CI 75, SL 0.31, SI 69, MW 0.33, ML 0.85, PL 0.35, PH 0.32, PW 0.27, PI1 107, PI2 77, PPL 0.41, PPH 0.45, PPW 0.42, PPI1 91, PPI2 102, WI 156. Paratype worker: BL 3.10–3.25, HL 0.60–0.63, HW 0.43–0.45, CI 71–72.00, SL 0.30–0.33, SI 69–75, MW 0.33–0.35, ML 0.81–0.84, PL 0.33–0.34, PH 0.32–0.33, PW 0.26–0.28, PI1 102–105, PI2 75–85, PPL 0.40–0.43, PPH 0.42–0.43, PPW 0.40–0.43, PPI1 95–99, PPI2 100–102, WI 154–169 (n = 3) (individual measurements see Table 1).

Worker

HEAD. Antennae with 11 segments; apical antennal segment (XI) conspicuously enlarged, much broader than and longer than two preceding segments (IX–X) combined. In full-face view, head subrectangular, distinctly longer than broad, lateral margins slightly convex and convergent from middle to posterior, and posterior margin slightly concave, with posterolateral corner acute (Fig. 6A). Frontal carinae short and narrow, not reaching to middle of head, with anterior half sharply elevated and posterior half low and vestigial in profile. Clypeus short, anterior margin almost straight. Eyes and ocelli completely absent (Fig. 6A). Mandibles triangular; masticatory margin serrated, without distinct teeth (Fig. 6A).

MESOSOMA. In dorsal view, robust and subrectangular, with almost parallel lateral sides, anterior margin slightly convex, and posterior margin slightly concave (Fig. 6B); in lateral view, dorsal outline slightly convex; promesonotal suture and metanotal groove absent; mesopleuron demarcated from pronotum by distinct suture, but not demarcated from mesonotum and metapleuron; mesopleuron also not differentiated from metapleuron (Fig. 6C). Propodeal declivity straight, encircled with distinct thin rim (Fig. 6C).

METASOMA. In dorsal view, petiole elliptical, longer than broad (Fig. 6B); in lateral view, petiolar node (excluding sternite) subtrapezoid with slightly convex dorsal outline, anterior edge slightly concave and poster edge sloped (Fig. 6C). Subpetiolar process in lateral view subtrapezoid, with posterior outline strongly concave, and ventral margin very short and almost straight (Fig. 6C). Postpetiole in dorsal view clearly larger than petiole, subtrapezoid, anterior margin slightly concave and shorter than posterior margin, lateral margins slightly convex (Fig. 6B); in lateral view, postpetiole higher than long; dorsal outline obviously convex; postpetiolar sternite in lateral view really low, with ventral margin convex, anteroventrally produced as nearly orthogonal directed downward and forward (Fig. 6C). Abdominal tergite IV (second gastral tergite) in dorsal view elongate, subrectangular with lateral margins slightly convex, anterior margin strongly concave, and lateral margins slightly convex (Fig. 6B).

SCULPTURE. Entire body with numerous, relatively small, closely-spaced foveolae (Fig. 6). Antennal scape, outer surface of mandible, and legs with fine dense micropunctures (Fig. 6).

PILOSITY. Body entirely covered with densely long and short decumbent, sub-erect and erect hairs.

COLORATION. Body brown. Antennae and legs yellowish brown (Fig. 6).

Distribution

China (Guizhou, Xizang).

Habitat

The nest of *Syscia guizhouensis* was found in the Fanjingshan National Nature Reserve, Guizhou, China. The sampled site is located in an evergreen broad-leaf forest, on the soil surface under the falling leaves.



Fig. 7. *Syscia guizhouensis* (Zhou, 2006) comb. nov., worker (No. G160363). **A**. Head in full-face view. **B**. Petiole and postpetiole in dorsal view. **C**. Body in dorsal view. **D**. Body in lateral view (imaged by Zhilin Chen).

Recognition

Syscia guizhouensis is similar to *S. humicola*, but can be distinguished from *S. humicola* by the following characteristics: 1) subpetiolar process in lateral view subrectangular in *S. guizhouensis* (subpetiolar process in lateral view subtriangular in *S. humicola*); 2) in full-face view posterior half of the head trapezoidal, narrower posteriorly in *S. guizhouensi* (in full-face view posterior half of the head rectangular in *S. humicola*); 3) in lateral view anterior margin of petiole concave in *S. guizhouensis* (in lateral view anterior margin of petiole sloped and slightly convex in *S. humicola*).

Syscia guizhouensis is also similar to a species from Thailand, *S. chaladthanyakiji*, in the general appearance of workers with 11-segmented antennae. However, *S. guizhouensis* can be distinguished from *S. chaladthanyakiji* by the following characteristics: 1) dorsal outline of mesosoma in lateral view slightly convex in *S. guizhouensis* (nearly straight in dorsal view in *S. chaladthanyakiji*); 2) petiole in dorsal view obviously elliptical, longer than broad; in lateral view trapezoidal with slightly convex dorsal outline, bilateral edge slop and anterior edge slightly concave in *S. guizhouensi* (obviously rectangular in dorsal view, broader than long; in lateral view rectangular with convex dorsal outline, bilateral edge vertical in *S. chaladthanyakiji*); 3) subpetiolar process in lateral view subtrapezoid, with posterior outline strongly concave, and ventral margin very short and almost straight in *S. guizhouensis* (subrectangular with ventral outline concave in *S. chaladthanyakiji*); 4) postpetiolar sternite in lateral view anteroventrally produced as nearly orthogonal directed downward and forward in *S. guizhouensis* (reduced into blunt angle directed downward and forward in *S. chaladthanyakiji*); 5) body color darker, reddish brown in *S. guizhouensis* (brown in *S. chaladthanyakiji*).

Syscia reticularis Jaitrong, Wiwatwitaya & Yamane, 2020 Fig. 8

Syscia reticularis Jaitrong et al, 2020: 6, figs 7-10, 11 (w.). Thailand.

Diagnosis

Workers have 11-segmented antennae. Body color dark reddish brown. Head in full-face view elliptical, distinctly longer than broad, lateral margins strongly convex, posterior margin almost straight, with posterolateral corner as blunt. Mesosoma in dorsal view rather robust, subrectangular, with almost parallel-sided lateral sides; in lateral view, dorsal outline slightly convex; propodeal declivity flat, encircled with indistinct rim. Petiole in dorsal view subquadrate, almost as long as broad, lateral margins nearly parallel-sided; in lateral view (excluding sternite) subrectangular, slightly shorter than high, dorsal outline convex, bilateral edge vertical; subpetiolar process in lateral view subrectangular with its ventral outline concave; ventrolateral surface of petiole with two distinct almost parallel carinae just above subpetiolar process. Postpetiole in lateral view rounded, shorter than high, with ventral outline concave; sternite anteroventrally produced as blunt angle directed downward and forward. Entire body finely reticulated with deep bottoms; antennal scape, outer face of mandible and legs punctate.

Type material

Unexamined. But the holotype of *S. reticularis* was examined from images available from AntWeb (https://www.antweb.org; THNHM-I-02624, imaged by Yudthana Samsung).

Distribution

Thailand.



Fig. 8. *Syscia reticularis* Jaitrong, Wiwatwitaya & Yamane, 2020, holotype, worker (THNHM-I-02624). A. Head in full-face view. **B**. Label. **C**. body in dorsal view. **D**. Body in lateral view (images cited from https://www.antweb.org; imaged by Yudthana Samung).

Syscia typhla Roger, 1861 Fig. 9

Syscia typhla Roger, 1861: 20 (w.). Sri Lanka.

Cerapachys typhla – Emery 1902: 24. *Cerapachys typhlus* – Xu 1998: 119. — Xu *et al.* 1999: 382. — Gu *et al.* 2019: 147, misidentification. *Syscia typhla* – Borowiec 2016: 224.

Diagnosis

Workers have 9-segmented antennae. Head in full-face view subrectangular, distinctly longer than broad, lateral margins slightly convex, posterior margin also slightly convex. Mesosoma in dorsal view rather elongate, long-ellipsoidal, with lateral sides and anterior margin obviously convex in dorsal view; in lateral view, dorsal outline convex; propodeal declivity feebly concave, encircled with, distinct thin rim. Petiole in dorsal view subtrapezoid in dorsal view, anterior margin shorter than posterior margin; in lateral view (excluding sternite) subquadrate, almost as long as high, lateral margins convex; subpetiolar process in lateral view subtriangle, with ventral corner clypeiform, ventral outline strongly concave. Postpetiole in lateral view subrectangular, shorter than high, with ventral outline slightly concave; sternite anteroventrally produced as blunt angle directed downward and forward.

Type material

Unexamined. But the type worker of *S. typhla* was examined in the online database of the AntWeb (https://www.antweb.org/; FOCOL0804, imaged by Christiana Klingenberg).

Distribution

Japan (doubtful), West Bengal, Sri Lanka.

Remarks

Based on a reexamination the specimens which were reported from Yunnan of China by Xu (1998) and Xu *et al.* (1999), we found that those specimens were misidentified as *Cerapachys typhlus*, but *Ooceraea biroi* (Forel, 1907). In addition, Gu *et al.* (2019) reported that the species *Cerapachys typhlus* was also distributed in Guangxi of China, which was a misidentification of *S. zhoui* sp. nov. Therefore, the species *S. typhla* is excluded from the ant fauna of China.

Syscia zhoui sp. nov. urn:lsid:zoobank.org:act:02E6945A-8EE0-4244-85F8-6CC638006484 Fig. 10, Table 1

Cerapachys typhus Gu et al., 2019: 147, misidentification.

Diagnosis

Workers have 9-segmented antennae, body color reddish brown with antennae and legs yellowish brown. Head in full-face view subrectangular, longer than broad. Mesosoma in dorsal view subrectangular with almost parallel lateral sides; in lateral view, dorsal outline slightly convex. Petiole in dorsal view subquadrate, with lateral margins convex; in lateral view, petiolar node (excluding sternite) subquadrate with slightly convex dorsal outline, bilateral edge sloped slightly; subpetiolar process in lateral view subtriangle. Postpetiole in lateral view slightly higher than long, sternite anteroventrally produced as nearly orthogonal directed downward and forward.



Fig. 9. *Syscia typhla* Roger, 1861, holotype, worker (FOCOL0804) **A**. Head in full-face view. **B**. Label. **C**. Body in dorsal view. **D**. Body in lateral view (images cited from https://www.antweb.org; imaged by Christiana Klingenberg).

Etymology

The new species is named in honor of Shanyi Zhou (Guangxi Normal University) for his outstanding contributions to the ant fauna of China.

Type material

Holotype (worker)

CHINA • Guangxi, Chongzuo City, Zuozhou Town, PaiRu Village in the Chongzuo White-Headed Langur National Reserve; 22.555816° N, 107.416772° E; 242.00 m a.s..l.; 13 Jun. 2016; Zhilin Chen leg.; No. GXNU160017.

Paratypes (workers)

CHINA • 1 worker; same collection data as for holotype; GXNU: GXNU160017-1 • 1 worker; same collection data as for holotype; SWFU: GXNU160017-2 • 1 worker; same collection data as for holotype; IZCAS: GXNU160017-3.

Description

MEASUREMENTS AND INDICES. Holotype worker: BL 3.55, HL 0.65, HW 0.53, CI 82, SL 0.42, SI 79, MW 0.38, ML 0.60, PL 0.30, PH 0.30, PW 0.30, PI1 100, PI2 100, PPL 0.44, PPH 0.45, PPW 0.40, PPI1 97, PPI2 91, WI 133. Paratype workers: BL 3.75–3.98, HL 0.62–0.69, HW 0.50–0.56, CI 77–85, SL 0.42–0.43, SI 76–85, MW 0.31–0.38, ML 0.65–0.80, PL 0.30–0.33, PH 0.25–0.30, PW 0.25–0.35, PI1 100–120, PI2 83–117, PPL 0.45–0.48, PPH 0.42–0.48, PPW 0.42–0.45, PPI1 99–111, PPI2 89–93, WI 121–168 (n = 3) (individual measurements see Table 1).

Worker

HEAD. Antennae with 9 segments; scape short, not reaching mid-length of head when folded back (Fig. 10A). In full-face view, head subrectangular, longer than broad, lateral margins slightly convex and convergent posteriorly, posterior margin slightly concave, with posterolateral corner acute (Fig. 10A). Frontal carinae short and narrow, not reaching to middle of head, with anterior half sharply elevated and posterior half low and vestigial in profile. Clypeus short, anterior margin almost straight. Eyes and ocelli completely absent (Fig. 10A). Mandibles triangular; masticatory margin serrated, without distinct teeth (Fig. 10A).

MESOSOMA. In dorsal view, robust and subrectangular, with almost parallel lateral sides, anterior margin slightly convex, and posterior margin slightly concave (Fig. 10B); in lateral view, dorsal outline slightly convex; promesonotal suture and metanotal groove absent; mesopleuron demarcated from pronotum by distinct suture, but not demarcated from mesonotum and metapleuron; mesopleuron also not differentiated from metapleuron (Fig. 10D). Propodeal declivity straight, encircled with distinct thin rim (Fig. 10D).

METASOMA. In dorsal view, petiole subquadrate, with lateral margins convex (Fig. 10C); in lateral view, petiolar node (excluding sternite) subquadrate with slightly convex dorsal outline, bilateral edge sloped slightly (Fig. 10D). Subpetiolar process in lateral view subtriangle, with ventralopasterior corner acute and posterior outline strongly concave (Fig. 10D). Postpetiole in dorsal view clearly larger than petiole, subtrapezoid, anterior margin slightly concave and shorter than posterior margin, lateral margins apparently convex (Fig. 10C); in lateral view, postpetiole apparently shorter than high; dorsal outline slightly convex; postpetiolar sternite in lateral view really low, with ventral margin convex, anteroventrally produced as nearly orthogonal directed downward and forward (Fig. 10D). Abdominal tergite IV (second gastral tergite) in dorsal view elongate, subrectangular with lateral margins slightly convex, anterior margin strongly concave, and lateral margins obviously convex (Fig. 10C).



Fig. 10. *Syscia zhoui* sp. nov., holotype, worker (GXNU160017). **A**. Head in full-face view. **B**. Petiole and postpetiole in dorsal view. **C**. Mesosoma in dorsal view. **D**. Body in lateral view (imaged by Zhilin Chen).

Sculpture. Entire body with numerous, relatively small, closely-spaced foveolae (Fig. 10). Antennal scape, outer surface of mandible, and legs with fine dense micropunctures (Fig. 10A, D).

PILOSITY. Body entirely covered with densely long and short decumbent, sub-erect and erect hairs.

COLORATION. Body reddish brown. Antennae and legs yellowish brown (Fig. 10).

Distribution

China (Guangxi).

Habitat

The nest of *Syscia zhoui* sp. nov. was found in the Chongzuo White-Headed Langur National Reserve, Guangxi, China. The sampled site is located in an evergreen broad-leaf forest, on the soil surface under the falling leaves.

Recognition

Syscia zhoui sp. nov. is a special species with antennae 9 segments, similar to *S. typhla* Roger, 1861, distinctly distinguished from all other species in *Syscia* with antennae 11 segments. However, *S. zhoui* can be distinguished from *S. typhla* by the following characteristics: 1) head in lateral view trapezoidal in *S. zhoui* (head in lateral view oval in *S. typhla*); 2) mesosoma rather robust, subrectangular with almost parallel lateral sides, and anterior margin slightly convex in dorsal view in *S. zhoui* (mesosoma rather elongate, long-ellipsoidal, with lateral sides and anterior margin obviously convex in dorsal view in *S. typhla*); 3) both anterior margin and posterior margin of petiole straight in *S. zhoui* (both anterior margin of petiole convex in *S. typhla*); 4) ventroposterior corner of subpetiolar process blunt rounded in *S. typhla*); 5) postpetiole in dorsal view wider than long in *S. zhoui* (postpetiole in dorsal view longer than its width in *S. typhla*).

Discussion

Syscia is a representative group of rare ants belonging to the subfamily Dorylinae with a disjunct distribution between the Old and New World. Previously, *Syscia* was included in the genus *Cerapachys*, but recent phylogenetic studies have revived it as a valid genus. Before this study, only 5 species of *Syscia* were recorded in the Old World. The low number of species of *Syscia* in the Old World compared to the recorded species richness (34 species) found in the New World indicates that there should be more speces of *Syscia* yet to be described from collections in the Old World. Eastern Asia and China represent one of the most important regions of the Old World, due to the vast land area, heterogeneous landscape, and diverse climate all contributing to a high biodiversity. In this paper, *S. typhla* is removed from the ant fauna of China, whereas two new species, *Syscia arcodorsa* sp. nov. and *S. zhoui* sp. nov. are added. *Cerapachys guizhouensis* Zhou, 2006 is re-described and revised as a member of *Syscia*, *S. guizhouensis* (Zhou, 2006). These results lay an important foundation for a new understanding of the myrmecofauna of China and Eastern Asia.

Acknowledgments

This study was supported by the National Natural Science Foundation of China (No. 32360127), Natural Science Foundation of Guangxi (2022GXNSFAA035524), National Animal Collection Resource Center of China, Key Laboratory of Ecology of Rare and Endangered Species and Environmental Protection (Guangxi Normal University), Ministry of Education and Guangxi Key Laboratory of Rare and Endangered Animal Ecology (Guangxi Normal University). This research also was supported by the National Animal Collection Resource Center, China.

References

Abe T. 1974. Notes on the fauna of ants in Iriomote Island. [In Japanese.]. *In*: Ikehara S. (ed.) *Ecological Studies of Nature Conservation of the Ryukyu Islands* (1): 105–111. Report for the Fiscal Year of 1973. [In Japanese.]. University of the Ryukyus, Naha, Okinawa.

AntWeb. Version 8.75.3. California Academy of Science, Available from https://www.antweb.org [accessed 27 Apr. 2022].

Aswaj P., Sahanashree R., Udayakantha W.S., Aniruddha M. & Priyadarsanan D.R. 2021. Two new species of doryline ants (Hymenoptera, Formicidae) with 11-segmented antennae from India. *ZooKeys* 1056: 59–72. https://doi.org/10.3897/zookeys.1056.68722

Bingham C.T. 1903. *The Fauna of British India, Including Ceylon and Burma. Hymenoptera, Vol. II. Ants and Cuckoo-wasps.* Taylor and Francis, London. https://doi.org/10.5962/bhl.title.100740

Bharti H., Rilta J.S. & Dhadwal T. 2021 Two new species of *Ooceraea* (Hymenoptera, Formicidae, Dorylinae) from India with ten-segmented antennae. *ZooKeys* 1010: 165–183. https://doi.org/10.3897/zookeys.1010.58436

Borowiec M.L. 2016. Generic revision of the ant subfamily Dorylinae (Hymenoptera, Formicidae). *ZooKeys* 608: 1–280. https://doi.org/10.3897/zookeys.608.9427

Brady S.G., Fisher B.L., Schultz T.R., Ward P.S. 2014. The rise of army ants and their relatives: diversification of specialized predatory doryline ants. *BMC Evolutionary Biology* 14 (93): 1–14. https://doi.org/10.1186/1471-2148-14-93

Brown W.L. 1975. Contributions toward a reclassification of the Formicidae. V. Ponerinae, tribes Platythyreini, Cerapachyini, Cylindromyrmecini, Acanthostichini, and Aenictogitini. Search. *Agriculture (Ithaca, New York)* 5 (1): 1–115.

Dalla Torre K.W. 1893. *Catalogus Hymenopterorum hucusque descriptorum systematicus et synonymicus*. Vol. 7. Formicidae (Heterogyna). W. Engelmann, Leipzig. https://doi.org/10.5962/bhl.title.10348

Donisthorpe H. 1943. A list of the type-species of the genera and subgenera of the Formicidae. [concl.]. *Annals and Magazine of Natural History* (11) 10: 721–737. https://doi.org/10.1080/00222934308527388

Emery C. 1902. Note mirmecologiche. *Rendiconti delle Sessioni della Reale Accademia delle Scienze dell'Istituto di Bologna (n.s.)* 6: 22–34.

Emery C. 1911. Hymenoptera. Fam. Formicidae. Subfam. Ponerinae. Genera Insectorum 118: 1-125.

Forel A. 1900. Les formicides de l'empire des Indes et de Ceylan. Part VII. *Journal of the Bombay Natural History Society* 13: 303–332.

Forel A. 1917. Cadre synoptique actuel de la faune universelle des fourmis. *Bulletin de la Société Vaudoise des Sciences Naturelles* 51: 229–253.

Ghosh S.N., Chattopadhyay A. & Bhattacharyya D.K. 2007. Studies on roadside soil inhabiting ants (Hymenoptera: Formicidae) of Kolkata with reference to the effects of lead emitted through automobile exhaust. *Zoological Survey of India, Occ. Paper* 257: 1–149.

Gu B., Chen Z.L., Wei X., Qin R.Y. & Zhou S.Y. 2019. Faunal analysis of ant of Guangxi Chongzuo White-headed Langur Nature Reserve, China. *Journal of Guangxi Normal University (Nature Science Edition)* 37 (4): 144–152. https://doi.org/10.16088/j.issn.1001-6600.2019.04.018

Holt B.G., Lessard J.P., Borregaard M.K., Fritz S.A., Araújo M.B., Dimitrov D., Fabre P.-H., Graham C.H, Graves G.R., Jønsson K.A., Nogués-Bravo D., Wang Zhiheng, Witthaker R.J., Fjelsdå J. & Rahbek C. 2013. An update of Wallace's zoogeographic regions of the world. *Science* 339 (6115): 74–78. https://doi.org/10.1126/science.1228282

Jaitrong W., Wiwatwitaya D. & Yamane S. 2020. First record of the ant genus *Syscia* Roger, 1861 (Hymenoptera: Formicidae) from Thailand, with descriptions of two new species. *Far Eastern Entomologist* 411: 1–9. https://doi.org/10.25221/fee.411.1

Kempf W.W. 1972. Catálogo abreviado das formigas da região Neotropical. *Studia Entomologica* 15: 3–344.

Longino J.T. & Branstetter M.G. 2021. Integrating UCE phylogenomics with traditional taxonomy reveals a trove of New World *Syscia* species (Formicidae: Dorylinae). *Insect Systematics and Diversity* 5 (2): 1–51. https://doi.org/10.1093/isd/ixab001

Roger J. 1861. Die Ponera-artigen Ameisen (Schluss). Berliner Entomologische Zeitschrift 5: 1-54.

Sonobe R. 1973. Ant fauna of the Sesoko Island, Okinawa. *Sesoko Marine Science Laboratory Technical Report* 2: 15–16.

Wheeler W.M. 1902. An American *Cerapachys*, with remarks on the affinities of the Cerapachyinae. *Biological Bulletin (Woods Hole)* 3: 181–191. https://doi.org/10.2307/1535872

Wheeler W.M. 1910. Ants: Their Structure, Development and Behavior. Columbia University Press, New York. https://doi.org/10.5962/bhl.title.1937

Wheeler W.M. 1922. Ants of the American Museum Congo expedition : a contribution to the myrmecology of Africa. *Bulletin of the American Museum of Natural History*: 45: 1–1139.

Xu Z.H. 1998. A report of fourty-one ant species newly recorded in China from Xishuangbanna District of Yunnan Province (Hymenoptera: Formicidae). *Zhongguo Xue Shu Qi Kan Wen Zhai* 4: 1119–1121.

Xu Z.H., Yang B.L. & Hu G. 1999. Formicidae ant communities in fragments of montane rain forest in Xishuangbanna, China. *Zoological Research* 20 (4): 288–293.

Zhou S.Y. 2006. Formicidae. In: Li Z.Z. & Jin D.C. (eds) Insects from Fanjingshan Landscape: 579–589. Guizhou Science and Technology Publishing House, Guiyang.

Manuscript received: 22 October 2022 Manuscript accepted: 23 November 2023 Published on: 2 April 2024 Topic editor: Tony Robillard Section editor: Fabio Cianferoni Desk editor: Eva-Maria Levermann

Printed versions of all papers are also deposited in the libraries of the institutes that are members of the *EJT* consortium: Muséum national d'histoire naturelle, Paris, France; Meise Botanic Garden, Belgium; Royal Museum for Central Africa, Tervuren, Belgium; Royal Belgian Institute of Natural Sciences, Brussels, Belgium; Natural History Museum of Denmark, Copenhagen, Denmark; Naturalis Biodiversity Center, Leiden, the Netherlands; Museo Nacional de Ciencias Naturales-CSIC, Madrid, Spain; Leibniz Institute for the Analysis of Biodiversity Change, Bonn – Hamburg, Germany; National Museum of the Czech Republic, Prague, Czech Republic.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: European Journal of Taxonomy

Jahr/Year: 2024

Band/Volume: 0930

Autor(en)/Author(s): Xu Zhenghui, Du Congcong, Chen Zhilin

Artikel/Article: <u>Taxonomic account of the ant genus Syscia Roger, 1816</u> (<u>Hymenoptera, Formicidae</u>) from Asia, with descriptions of two new species from <u>China 157-181</u>