



Received: 7 October 2024 • Accepted: 27 March 2025 • Published: 10 July 2025

Topic editor: Tony Robillard • Section editor: Jurate De Prins • Desk editor: Radka Rosenbaumová

Research article

urn:lsid:zoobank.org:pub:398EFB77-4619-4AEE-B9BA-65282C2C9F13

New species of *Notalina* Mosely, 1939 (Trichoptera: Leptoceridae) from Chapada Diamantina National Park, Northeast Brazil, with new records of other leptocerids from the Caatinga biome

Ana L. HENRIQUES-OLIVEIRA^{1,*}  , André A. ALVES²   &
Felipe F.F. MOREIRA³  

^{1,2}Laboratório de Entomologia – LABENT, Departamento de Zoologia, Instituto de Biologia, Universidade Federal do Rio de Janeiro – UFRJ, Av. Carlos Chagas Filho, 373, sala A1-107, Cidade Universitária, CEP 21941-971, Rio de Janeiro, RJ, Brazil.

^{1,3}Laboratório de Entomologia – LABE, Instituto Oswaldo Cruz, Fundação Oswaldo Cruz – FIOCRUZ, Avenida Brasil, 4365, Pavilhão Mourisco, sala 214, Manguinhos, CEP 21040-360, Rio de Janeiro, RJ, Brazil.

*Corresponding author: alholiveira@gmail.com

²Email: andrjrje@gmail.com

³Email: ppmeiameiameia@gmail.com

Abstract. The genus *Notalina* Mosely, 1939 contains 29 species divided into two subgenera: *Notalina* (*Notalina*) Mosely, 1939, which includes 15 endemic species from Australia and Tasmania, and *Notalina* (*Neonotalina*) Holzenthal, 1986, which comprises 14 species exclusively from South America. In Brazil, eleven species have been recorded, with the main distribution areas in the Cerrado and Atlantic Forest biomes of the Southeast and Central-West regions, while only three species have been recorded in the Northeast region. In this paper, we describe and illustrate two new species of *Notalina* (*Neonotalina*), *Notalina* (*Neonotalina*) *diamantina* sp. nov. and *Notalina* (*Neonotalina*) *daniae* sp. nov., from Chapada Diamantina National Park, Bahia, Northeast Brazil. We also provide new distributional records for other leptocerids from the Caatinga biome.

Keywords. Hudsonemini, taxonomy, South America, long-horned caddisflies.

Henriques-Oliveira A.L., Alves A.A & Moreira F.F.F. 2025. New species of *Notalina* Mosely, 1939 (Trichoptera: Leptoceridae) from Chapada Diamantina National Park, Northeast Brazil, with new records of other leptocerids from the Caatinga biome. *European Journal of Taxonomy* 1004: 1–18. <https://doi.org/10.5852/ejt.2025.1004.2957>

Introduction

The long-horned caddisfly Leptoceridae Leach, 1815 is one of the largest three families of Trichoptera Kirby, 1813, with more than 2235 described species and 49 genera worldwide (Holzenthal & Calor 2017; Morse *et al.* 2019). Currently, the family is divided into four subfamilies: Grumichellinae

Morse, 1981 and Triplectidinae Ulmer, 1906, both occurring mainly in the Australian and Neotropical regions; Leptocerinae Ulmer, 1903 with a cosmopolitan distribution; and Leptorussinae Morse, 1981, which includes the single Australian-endemic genus *Leptorussa* Mosely, 1953 (in Mosely & Kimmins 1953) (Malm & Johanson 2011). In Brazil, Leptoceridae is represented by 103 species in ten genera: *Achoropsyche* Holzenthal, 1984, *Amazonatolica* Holzenthal & Pes, 2004, *Atanatolica* Mosely, 1936, *Grumichella* Müller, 1879, *Ibyacerina* Henriques-Oliveira, Silva, Nessimian & Takiya, 2021, *Nectopsyche* Müller, 1879, *Neoathripsodes* Holzenthal, 1989, *Notalina* Mosely, 1936, *Oecetis* McLachlan, 1877, and *Triplectides* Kolenati, 1859. Among these, *Achoropsyche*, *Amazonatolica*, and *Ibyacerina* are monotypic and endemic (Calor *et al.* 2024a).

The genus *Notalina* belongs to the tribe Hudsonemini of the subfamily Triplectidinae and was established by Mosely (1936) for three Australian species: *Notalina parkeri* Mosely, 1936, *N. delicatula* (Ulmer, 1908), and *N. flava* (Ulmer, 1908). Since then, many species have been described. Currently, *Notalina* is divided into two subgenera: *Notalina* (*Notalina*) Mosely, 1936, with 15 endemic species from Australia and Tasmania, and *Notalina* (*Neonotalina*) Holzenthal, 1986 with 14 species exclusively from South America (Holzenthal & Calor 2017; Henriques-Oliveira *et al.* 2018; Silva-Pereira *et al.* 2022; Calor *et al.* 2024b). Holzenthal (1986) informally recognized two species groups in the Neotropical subgenus *Neonotalina*, based mainly on characters of the male genitalia: *N. roraima* species group, occurring in the upper Amazon basin, the Northern Andes, and the Guiana Highlands, represented by *Notalina* (*Neonotalina*) *mathiasi* Holzenthal, 1986, *N. (Neonotalina) nanay* Holzenthal, 1986, and *N. (Neonotalina) roraima* Holzenthal, 1986, and *N. brasiliensis* species group mainly occurring in the highlands of the Cerrado and Atlantic Forest biomes of the Southeast and Central areas of the country, which includes *Notalina* (*Neonotalina*) *brasiliensis* Holzenthal, 1986, *N. (Neonotalina) cipo* Holzenthal, 1986, *N. (Neonotalina) claudiofroehlichii* Desidério & Calor, 2024 (in Calor *et al.* 2024b), *N. (Neonotalina) franciscana* Henriques-Oliveira, Rocha & Nessimian, 2018, *N. (Neonotalina) froehlichii* Calor & Holzenthal, 2006 (in Calor *et al.* 2006), *N. (Neonotalina) goianensis* Calor, 2008, *N. (Neonotalina) hamiltoni* Holzenthal, 1986, *N. (Neonotalina) jordanensis* Henriques-Oliveira, Spies & Dumas, 2012, *N. (Neonotalina) morsei* Holzenthal, 1986, *N. (Neonotalina) paulista* Calor & Holzenthal, 2006 (in Calor *et al.* 2006), and *N. (Neonotalina) ralphi* Silva-Pereira, Oliveira, Desidério, Calor & Hamada, 2022 (Calor *et al.* 2024a).

The Caatinga biome extends over an area of approximately 800 000 km² in Northeastern Brazil, encompassing parts of nine Brazilian states: Alagoas, Bahia, Ceará, Paraíba, Pernambuco, Piauí, Rio Grande do Norte, and Sergipe in the Northeast region, plus Minas Gerais in the Southeast region. According to Prado (2003), the Caatinga is a mosaic of different forest types, characterized by short trees and shrubs with xerophytic traits, influenced by climatic extremes when compared to other Brazilian formations: highest solar radiation, highest annual temperature means, lowest rates of relative humidity, and lowest precipitation, which is limited to a very short period of the year. Currently, three species of *Notalina* (*Neonotalina*) are recorded from the Northeast region: *N. brasiliensis*, *N. cipo*, and *N. claudiofroehlichii* (Calor *et al.* 2024b). Here, we describe and illustrate two new species of *Notalina* (*Neonotalina*) from Chapada Diamantina National Park (PNCD) and its surroundings within the Caatinga biome. We also provide new records of other Leptoceridae from Northeast Brazil.

Material and methods

Study area

Specimens of caddisflies were collected from various streams in Chapada Diamantina National Park and its surroundings, Bahia, Brazil (Fig. 1). Chapada Diamantina National Park (PNCD), covering an area of 152 000 hectares, is located in the Serra do Sincorá, in the central region of Chapada Diamantina (plateau). It spans the municipalities of Lençóis, Mucugê, Ibicoara, Andaraí, and Palmeiras, in the state of Bahia (MMA 2007). Chapada Diamantina forms a large set of plateaus in the northern part of the Serra do Espinhaço, a mountain range that stretches from the states of Minas Gerais to Bahia. The Chapada Diamantina rises like an imposing wall, with average altitudes over 1000 meters, acting as a watershed divide between the rivers flowing into the São Francisco River basin and those heading directly to the




Fig. 1. Distribution map of the *Notalina* Mosely, 1936 and *Nectopsyche* Müller, 1879 species treated in this study.

Atlantic Ocean (IBGE 1977). The park boasts a significant ecological and environmental diversity, being part of both the Caatinga (RBCAAT) and Atlantic Forest (RBMA) Biosphere Reserves (MMA 2007).

Sampling and illustrations

Adult specimens were collected using a collapsible light trap (Nessimian *et al.* 2024), a Malaise trap (Gressitt & Gressitt 1962), suspended over bodies of water, a white sheet, or by hand net. All specimens were preserved in 80–96% ethanol. To observe the genital structures, the abdomen was removed and cleared using 10% KOH or 85% lactic acid, as described by Blahnik *et al.* (2007). Abdomens were temporarily mounted on a cavity slide with glycerin for viewing and drawing and then transferred to a microvial with 80% ethanol for storage alongside the remainder of the specimen. The terminology for adults follows that of Holzenthal (1986) and Calor *et al.* (2006) for males, and Ito *et al.* (2013) for females.

Pencil sketches of wings and genital structures were made under a stereo microscope (Carl Zeiss, model Stemi SV6) and a compound microscope (Carl Zeiss, model AxioLab), both equipped with a camera lucida. The sketches were scanned and imported into an Adobe Illustrator (ver. 22.0.1, Adobe Systems, Inc.) document to create graphic vectors. Photographs were taken with a digital Leica Camera (DFC450) coupled to a Leica stereo microscope (M205C). A series of photographs of each structure at different focal planes were stacked using the Leica Application Suite (ver. 4.6.2). The map was created using QGIS ver. 3.34.4 (QGIS Development Team 2020).

The types and all other examined specimens are deposited at the Coleção Entomológica Professor José Alfredo Pinheiro Dutra, Departamento de Zoologia, Universidade Federal do Rio de Janeiro, Rio de Janeiro (DZRJ), and some paratypes of the new species housed in Museu Nacional, Universidade Federal do Rio de Janeiro, Rio de Janeiro (MNRJ).

Abbreviations of taxonomic structures

alp	=	apicolateral process
apm. gon.	=	apicomesal process of gonopod plate
bdp	=	basodorsal process
bvp	=	basoventral process
d	=	discoidal cell
dmp	=	dorsomesal process
inf. app.	=	inferior appendage
mdp	=	mesodorsal process
mvp	=	mesoventral process
pr. app.	=	preanal appendages
sp. scl.	=	spermathecal sclerite
th	=	thyridial cell
I, III, V	=	forks I, III or V
VIII	=	segment VIII
IX	=	segment IX
X	=	segment X

Institutional abbreviations

DZRJ	=	Coleção Entomológica Prof. José Alfredo Pinheiro Dutra, Departamento de Zoologia, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil
MACN	=	Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”, Buenos Aires, Argentina
MCZ	=	Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts, USA
MNRJ	=	Museu Nacional, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil
MZBS	=	Museo de Zoologia, Barcelona, Spain

Results

Taxonomy

Class Insecta Linnaeus, 1758
Order Trichoptera Kirby, 1813
Suborder Integripalpia Martynov, 1924
Superfamily Leptoceroidea Leach, 1815
Family Leptoceridae Leach, 1815
Subfamily Triplectidinae Ulmer, 1906
Genus *Notalina* Mosely, 1936
Subgenus *Neonotalina* Holzenthal, 1986

Notalina (*Neonotalina*) *diamantina* sp. nov.

urn:lsid:zoobank.org:act:23271900-3C70-46AE-981D-0FD18FDF6087

Figs 2–6

Diagnosis

This new species can be identified by the following characteristics: segment X with a very long dorsomesal process, with a rounded apex and a tuft of very long setae, in lateral view, facing upwards at an angle $>45^\circ$; the apicolateral processes short, truncate, and equipped with stout setae; and the mesodorsal process of the inferior appendages digitate, long, and flat (like a blade), bearing a single seta on the apical third.

The new species is assigned to the *N. brasiliانا* species group as defined by Holzenthal (1986) based on the presence of acuminate flanges on the phallobase and the well-developed phallosclerite. *Notalina* (*Neonotalina*) *diamantina* sp. nov. resembles *N. brasiliانا*, *N. froehlichii*, and *N. goianensis*, due to the presence of long, setose apicodorsal processes of segment X. However, it differs from its relatives in the appearance of segment X in the dorsal view: in the new species, the apicodorsal processes are well developed, rounded, and enlarged apically with long setae. Furthermore, they are very long and well upturned dorsally with an inclination angle of almost 90° . In contrast, *N. brasiliانا* and *N. goianensis* have apicomesal processes that are long, thin, and have an inclination angle of 45° or less. Although *N. froehlichii*, in lateral view, shows apicomesal processes with the same angle of inclination as the new species, the apicolateral processes are digitate and rounded, while in *N. diamantina*, they are short and truncate. Another difference is that segment X in *N. froehlichii* is elevated in the middle length in lateral view, whereas in the new species, it is only slightly elevated near the base. The mesoventral process of the inferior appendages in the new species is subtriangular, similar to *N. brasiliانا* and *N. goianensis*, while in *N. froehlichii*, it is long and slender with a narrower apex.

Etymology

Species named in reference of the type locality Chapada Diamantina National Park (or Parque Nacional da Chapada Diamantina – PNCD, in Portuguese), where the individuals of the new species were collected. It is a noun in apposition.

Type material

Holotype

BRAZIL – Bahia • ♂; Palmeiras, Parque Nacional da Chapada Diamantina, near to Cachoeira Conceição dos Gatos; $12^\circ32'19''$ S, $41^\circ30'47.8''$ W; alt. 775 m; 12 May 2021; A.A. Alves leg.; white sheet; [alcohol]; DZSJ TRICHOPTERA9720.

Paratypes

BRAZIL – Bahia • 3 ♂♂; same data as for holotype; DZSJ TRICHOPTERA9721 • 2 ♂♂, 1 ♀; same data as for holotype; [alcohol]; MNRJ-ENT10–268 • 1 ♂; same data as for holotype; [pinned]; MNRJ-ENT10–269 • 1 ♂; Mucugê, Parque Nacional da Chapada Diamantina, Comunidade do Baixão, Rio




Fig. 2. *Notalina* (*Neonotalina*) *diamantina* sp. nov. (paratype, DZRJ TRICHOPTERA9725). **A.** Habitus, lateral view. **B.** Head and thorax, dorsal view. **C.** Living specimen. Scale bars = 1 mm.

Uma; 12°17'36.2" S, 41°14'28.2" W; alt. 741 m; 23 Jul. 2018; J.S. Prando leg.; sweep; [alcohol]; DZRJ TRICHOPTERA9722 • 1 ♂, 2 ♀♀; Mucugê, Chapada Diamantina, Rio Cumbucas, Mar de Espanha; 12°56'52.8" S, 41°21'31.1" W; alt. 858 m; 11 Mar. 2012; D.M. Takiya and A.P.M. Santos leg.; [alcohol]; DZRJ TRICHOPTERA9723 • 1 ♂; same data as for preceding; [pinned]; DZRJ TRICHOPTERA9725 • 1 ♂; same data as for preceding; DZRJ TRICHOPTERA9726 • 1 ♀; same data as for preceding; DZRJ TRICHOPTERA9727.

Description

Adult male (Figs 2–4)

COLOUR (pinned). Body and wings golden brown with off-white spots (Fig. 2). Legs, palps, and antennae golden brown (Fig. 2). Tibial spur formula 2, 2, 4. Forewing with forks I and V present (Fig. 3A); hindwing with forks I, III, and V present (Fig. 3B). Length of forewing = 8.0–9.5 mm. Length of hindwing = 6.0–7.0 mm (n = 11).

MALE GENITALIA (Fig. 4). Abdominal segment IX annular, in lateral view, broadest ventrolaterally, with setae on posterolateral edge, anterior margin slightly produced on median third (Fig. 4A). Preanal appendages digitate, slightly truncate, with long setae at apex, approximately ½ the length of segment X (Fig. 4B). Segment X, in lateral view, is slightly elevated at the base with a concave mesally, bearing two apical processes: a very long, digitate dorsomesal process with a rounded apex and a tuft of very long setae, facing upward dorsally (with an angle >45°, sometimes almost 90°); and a short, truncate apicolateral process with small stout setae at the apex (Fig. 4A). In dorsal view, segment X is broad at the base, with the posterior margin bearing two processes: a pointed lateral process with stout setae and a median process that is digitate and enlarged at the apex, very upturned, bearing a tuft of very long




Fig. 3. *Notalina* (*Neonotalina*) *diamantina* sp. nov., wing venation (holotype, DZRJ TRICHOPTERA9720). **A.** Forewing. **B.** Hind wings. Abbreviations: see Material and methods. Scale bar = 1 mm.

setae (Fig. 4B). The inferior appendage, in ventral view, is broad at the base with an elongate, digitate, and setose apical portion; the basoventral protuberance is short, rounded, and very setose when viewed laterally, and enlarged ventrally (Fig. 4C). The mesoventral process is elongate, with approximately half the length of the apical portion, rounded at the apex (Fig. 4C). In lateral view, it is subtriangular with a median carina, setose on the posterior margin, and has an acuminate apex. The mesodorsal process is long and digitate, flat (like a blade), bearing a long seta on the apical third, postero-dorsally directed with an acute apex when viewed laterally (Fig. 4A). The basodorsal process, in ventral view, is rounded and slightly higher than the basoventral protuberance (Fig. 4C). The phallic apparatus features a well-




Fig. 4. *Notalina (Neonotalina) diamantina* sp. nov., male genitalia (holotype, DZRJ TRICHOPTERA9720). **A.** Lateral view. **B.** Dorsal view. **C.** Ventral view. **D.** Phallic apparatus, ventral view. **E.** Phallic apparatus, lateral view. Abbreviations: see Material and methods. Scale bar = 0.1 mm.

developed phallobase with a pair of laterally directed acuminate flanges (Fig. 4D–E); the phallotremal sclerite is well developed, roughly U-shaped in ventral view (Fig. 4D), and oval and downcurved in lateral view (Fig. 4E).

Adult female (Figs 5–6)

COLOUR. Body golden brown (in alcohol), with some whitish setae on wings (Fig. 5): Tibial spur formula 2, 2, 4. Forewing with forks I and V present; hindwing with forks I, III, and V present (Fig. 5). Length of forewing = 7.0 mm. Length of hindwing = 5.5 mm (n = 4).

FEMALE GENITALIA (Fig. 6). Abdominal segment VIII, in ventral view, has a narrow sclerotized and setose plate, concave medially, presenting a rounded and rugose plate above (Fig. 6A–B). Segment IX sclerotized, annular, in dorsal view, almost straight, in lateral view, mesal portion produced and very setose (Fig. 6A), in ventral view, subquadrate, smooth, and slightly concave medially (Fig. 6B). Segment X forms a dorsal shelf, with a posterior margin slightly curved and bearing some stout setae, in lateral view, the posterior margin truncates (Fig. 6A–B). Preanal appendages short and subtruncate in dorsal



Fig. 5. *Notalina* (*Neonotalina*) *diamantina* sp. nov., female habitus (paratype, DZRJ TRICHOPTERA9723). **A.** Lateral view. **B.** Head and thorax, dorsal view. Scale bars = 1 mm.

view, oval and rounded with apex bearing long setae in lateral view (Fig. 6). Lamellae broad, in lateral view subquadrate, setose, with a posterior margin slightly concave (Fig. 6A). Gonopod plate broad, subtriangular, and sclerotized. Apicomesal process of gonopod plate rounded with small setae (Fig. 6B); Spermathecal sclerite in amphora-shape with central elongate, keyhole-shaped sclerites (Fig. 6B).

Immature stages

Unknown.

Distribution (Fig. 1)

Brazil (Bahia state).




Fig. 6. *Notalina (Neonotalina) diamantina* sp. nov., female genitalia (paratype, DZRJ TRICHOPTERA9723). A. Lateral view. B. Ventral view. C. Dorsal view. Abbreviations: see Material and methods. Scale bars = 0.1 mm.

Notalina (Neonotalina) daniae sp. nov.

urn:lsid:zoobank.org:act:13A4CE3F-DB5D-4519-8048-440DE1C7F385

Figs 7–8

Diagnosis

Notalina (Neonotalina) daniae sp. nov. is identifiable by the distinctive structure of segment X, which features a V-shaped apicomesal incision creating two large, rounded lobes. Additionally, in dorsal view, there are two small protuberances laterally at the mid-length of segment X with two setae in each one. The mesoventral processes of the inferior appendages are sub-quadrate with a pointed projection near the apex, while the mesodorsal processes are long, thin, and flat.

Notalina (Neonotalina) daniae sp. nov. is assigned to the *N. brasiliiana* species group as defined by Holzenthal (1986). The new species shares with *Notalina cipo*, *N. claudiofroehlichii*, *N. morsei*, and *N. ralphi* the shape of segment X with apicomesal incision forming two large lobes and the absence of an apicomesal process. However, *N. cipo* and *N. morsei* have a wider apicomesal incision of segment X and V-shaped with long lobes and straight apices, while in the new species the apicomesal incision is shorter than $\frac{1}{3}$ of tergum X. *Notalina daniae* can be confused with *N. ralphi* and *N. claudiofroehlichii* by showing a similar pattern in the apicolateral processes of segment X that forms two rounded and setose lobes, but in *N. claudiofroehlichii* these lobes are strongly sclerotized apically and the apicomesal incision of segment X is U-shaped, extending anteriorly ca $\frac{1}{2}$ of its length, while segment X of the new species is membranous. Although *Notalina ralphi* possesses the V-shaped incision extending anteriorly ca $\frac{1}{3}$ length of the segment like in *N. daniae*, it differs from *N. daniae* by the absence of lateral protuberances in segment X, and by having a row of 5–7 short, stout setae situated subapically, while *N. daniae* has two small protuberances at the middle-length of segment X in dorsal view. Another difference is observed in the mesoventral process of the inferior appendages in lateral view, where in the new species



Fig. 7. *Notalina (Neonotalina) daniae* sp. nov. (holotype, DZRJ TRICHOPTERA9724). **A.** Habitus, lateral view. Scale bar = 1 mm.

N. daniae the inferior appendage is subquadrate with a small projection near the apex and in *N. ralphi* and *N. claudiofroehlichii* the inferior appendage is indistinct or rounded.

Etymology

The species epithet ‘*daniae*’ is a noun in the genitive case to honor Prof. Dra Daniela ‘Dani’ Takiya, who collected the new species.




Fig. 8. *Notalina* (*Neonotalina*) *daniae* sp. nov., male genitalia (holotype, DZRJ TRICHOPTERA9724). **A.** Lateral view. **B.** Dorsal view. **C.** Ventral view. **D.** Phallic apparatus, ventral view. **E.** Phallic apparatus, lateral view. Abbreviations: see Material and methods. Scale bars = 0.1 mm.

Type material

Holotype

BRAZIL – Bahia • ♂; Mucugê, Igatú, Parque Nacional da Chapada Diamantina, Córrego das Perdizes com Rio Cumbucas; 12°58'29.8" S, 41°21'7.9" W; alt. 889 m; 10 Mar. 2012; A.P.M. Santos and D.M. Takiya leg.; white sheet; [pinned]; DZRJ TRICHOPTERA9724.

Description

Adult male (Figs 7–8)

COLOUR (pinned). Head and thorax brown with off-white spots. Legs, palps, and antennae golden brown (Fig. 7A). Tibial spur formula 2, 2, 4. Length of forewing: 7.00 mm (n = 1).

MALE GENITALIA (Fig. 8). Abdominal segment IX annular, in lateral view, broadest laterally, posterolateral margin with some setae above preanal appendages, posterior margin slightly produced medially (Fig. 8A); in dorsal view, sub-rectangular with posterior margin irregular and protruded laterally; anterior margin almost straight (Fig. 8B). Preanal appendages digitate, longer than half the length of segment X, rounded apically with long setae (Fig. 8A–B). Segment X, in lateral view, almost straight, featuring a small protuberance medially with two setae, and having a rounded apex with sparse stout setae; in dorsal view, broad at the base with two small membranous protuberances laterally at the middle-length of segment X and a V-shaped incision at the apex, dividing it into two rounded lobes (Fig. 8A). Inferior appendages, in ventral view, broad at base with apical portion elongate, digitate and setose; basodorsal process rounded and well developed, basoventral protuberance robust, rounded, and very setose (Fig. 8C); mesoventral process with approximately half-length of apical portion, rounded, protrude with a small point near apex; in lateral view, very large, subquadrate, setose on the posterior margin, with a pointed protuberance near the apex; mesodorsal process digitate long, thin, flat, like a blade with acute apex inward directed (Fig. 8A). Phallic apparatus with phallobase well developed and a pair of laterally directed acuminate flanges (Fig. 8D); phallotremal sclerite well developed, roughly Y-shaped, in ventral view (Fig. 8D).

Female and immature stages

Unknown.

Distribution (Fig. 1)

Brazil (Bahia state).

New records of other Leptoceridae species from Chapada Diamantina, Bahia

Family Leptoceridae Leach, 1815
Subfamily Leptocerinae Ulmer, 1903
Genus *Nectopsyche* Müller, 1879

Nectopsyche bruchi (Navás, 1920)

Fig. 1

Leptocella bruchi Navás, 1920: 66 (Type locality: Argentina, Monte Veloz, Estancia Barreto; MACN, ♂ reported).

Nectopsyche bruchi – Holzenthal & Calor 2017: 322 (catalogue). — Calor *et al.* 2024b (distribution).

Material examined

BRAZIL – Bahia • 10 ♂♂, 2 ♀♀; Palmeiras, PARNA Chapada Diamantina, near to Cachoeira Conceição dos Gatos; 12°32'19" S, 41°30'47.8" W; alt. 775 m; 12 May 2021; A.A. Alves leg.; white sheet; [alcohol and pinned]; DZRJ.

Remarks

In Brazil, this species has a more southeastern distribution recorded in Minas Gerais, Paraná, and Rio de Janeiro states, occurring in areas of the Atlantic Forest and Cerrado biome (Calor *et al.* 2024a). Herein, it is recorded for the first time the state of Bahia, and the Caatinga biome.

Distribution

Argentina, Brazil (states: Bahia (**new record**), Minas Gerais, Paraná and Rio de Janeiro), and Paraguay.

Nectopsyche muhni (Navás 1916)

Fig. 1

Leptocella muhni Navás, 1916: 68 (Type locality: Argentina, Santa Fé, MZBS, ♀ reported).

Nectopsyche muhni – Holzenthal & Calor 2017: 326 (catalogue). — Calor *et al.* 2024b (distribution).

Material examined

BRAZIL – **Bahia** • 1 ♂; Mucugê, PARNA Chapada Diamantina, Cachoeira das Andorinhas; 13°1'2.6" S, 41°20'38.7" W; alt. 961 m; 21–24 Aug. 2018; J.M.S. Rodrigues, J.S. Prando, F.F.F. Moreira, C.C. Gonçalves and R. Carrenho leg.; Malaise trap; DZ RJ • 4 ♂♂; Mucugê, PARNA Chapada Diamantina, road BA-242 near to bridge Dr. Heitor M. Chamusca; 12°59'24" S, 41°21'1.2" W; 20 Aug. 2018; C.C. Gonçalves leg.; white sheet; DZ RJ • 1 ♂; Palmeiras, PARNA Chapada Diamantina, near to Cachoeira Conceição dos Gatos; 12°32'19" S, 41°30'47.8" W; alt. 775 m; 12 May 2021; A.A. Alves leg.; white sheet; DZ RJ.

Distribution

Argentina, Bolivia, Brazil (states: Bahia (**new record**), Espírito Santo, Maranhão, Minas Gerais, Pará, Piauí, Roraima, Rio de Janeiro and São Paulo), Ecuador, Guyana, Paraguay, Peru, Suriname, and Venezuela.

Nectopsyche separata (Banks, 1920)

Fig. 1

Leptocella separata Banks, 1920: 353 (Type locality: Brazil, Santa Catarina, MCZ, ♂ reported).

Nectopsyche separata – Holzenthal & Calor 2017: 328 (catalogue). — Calor *et al.* 2024b. (distribution).

Material examined

BRAZIL – **Bahia** • 1 ♂; Mucugê, PARNA Chapada Diamantina, road BA-242 near to bridge Dr Heitor M. Chamusca; 12°59'24" S, 41°21'1.2" W; 20 Aug. 2018; C.C. Gonçalves leg.; white sheet; DZ RJ.

Remarks

Nectopsyche separata was described from Santa Catarina State, and has a south-southeast distribution pattern in the country, being observed occurring in both the Cerrado and Atlantic Forest biomes (Calor *et al.* 2024a). This is the first northernmost record of the species, and the first for the state of Bahia and the Caatinga biome.

Distribution

Argentina, Brazil (States: Bahia (**new record**), Espírito Santo, Minas Gerais, Rio de Janeiro, São Paulo, Paraná, and Santa Catarina), and Paraguay.

Discussion

The Caatinga is a fragile biome, significantly affected by human activities. It is estimated that more than 28% of the original vegetation of the Caatinga has been altered by human activities, such as slash and burn, agriculture, firewood harvesting, hunting, and grazing (Castelletti *et al.* 2003). Currently, less than 2% of its natural vegetation is surrounded by protected areas. Despite its low diversity and low number of endemic species, recent studies indicate it is an important component of Brazilian biodiversity (Leal *et al.* 2005).

The Caatinga, located between the Atlantic Forest and the Cerrado biomes, shares many biotic components with these domains. Current assessments of Caatinga biodiversity with regard to aquatic insects, mainly caddisflies, are certainly underestimated; despite the increase in recent years, few biological inventories have been concentrated in this region (e.g., Costa *et al.* 2014; Quinteiro *et al.* 2014; Takiya *et al.* 2016; Rafael *et al.* 2017; Calor *et al.* 2024b). Calor *et al.* (2024b) provided the latest inventory of caddisflies in the northeast region, where 138 species were recorded for Bahia. Here, although the collections were conducted at only a few sites in PNCD, we observed the existence of two new species of *Notalina*, as well as records of three species of *Nectopsyche*: *N. bruchi*, *N. munhi*, and *N. separata*.

This study contributes to the understanding of aquatic insect diversity, specifically caddisflies, in the region. The documentation of two new *Notalina* species (*Notalina* (*Neonotalina*) *diamantina* sp. nov. and *Notalina* (*Neonotalina*) *daniae* sp. nov.) and the records of the three additional *Nectopsyche* species from Chapada Diamantina highlights the importance of biodiversity assessments for conservation efforts in this unique biome.

Acknowledgements

We are grateful to the Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio) for issuing collection permits. We thank the director of the Chapada Diamantina National Park for all the assistance provided during the collection expeditions. We also thank the project coordinator, Daniela M. Takiya. This research was funded by the Conselho Nacional de Desenvolvimento Científico e Tecnológico - CNPq (Process. 421.413/2017-4). A.A. Alves was funded by CNPq, Process. 165746/2021-1. F.F.F. Moreira was funded by CNPq Process. 307115/2023-2, and the Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro FAPERJ, Process. E-26/201.362/2021 and E-26/203.152/2023. A.L. Henriques-Oliveira thanks the Conselho Nacional de Desenvolvimento Científico e Tecnológico for the post-doctoral financial support (CNPq Process.101308/2022-1).

References

- Banks N. 1920. New neuropteroid insects. *Bulletin of the Museum of Comparative Zoology* 64 (3): 297–362. <https://doi.org/10.5962/bhl.title.28705>
- Blahnik R.J., Holzenthal R.W. & Prather A. 2007. The lactic acid method for clearing Trichoptera genitalia. In: Bueno-Soria J., Barba-Alvarez R. & Armitage B. (eds) *Proceedings of the 12th International Symposium on Trichoptera*: 9–14. The Caddis Press, Columbus.
- Calor A.R. 2008. A new species of *Notalina* Mosely, 1936 (Trichoptera: Leptoceridae) from Chapada dos Veadeiros National Park, Goiás state, Brazil. *Biota Neotropica* 8 (3): 175–178. <https://doi.org/10.1590/S1676-06032008000300016>
- Calor A.R., Holzenthal R.W. & Amorim D.S. 2006. Phylogenetic analysis of *Notalina* (*Neonotalina*) Holzenthal (Trichoptera: Leptoceridae), with the description of two new species from southeastern Brazil. *Zootaxa* 1131 (1): 33–48. <https://doi.org/10.11646/zootaxa.1131.1.2>

- Calor A.R., Henriques-Oliveira A.L. & Quinteiro F.B. 2024a. Leptoceridae Leach, 1815. In: *Catálogo Taxonômico da Fauna do Brasil*. PNUD. Available from <http://fauna.jbrj.gov.br/fauna/faunadobrasil/278> [accessed 31 Jul. 2024].
- Calor A.R., Pereira R., Queiroz L.L., Vilarino A., Azevedo C.C.D., Queiroz A., Burgos-Miranda M.J., Cavalcante-Silva A., Oliveira-Silva M.V., Lucca G., Quinteiro F.B., Dias E.S., Gomes V.A., França D., Costa A.M., Desidério G.R., Santos A.P.M., Dumas L.L. & Bispo P.B. 2024b. LEAq – Laboratório de Entomologia Aquática “Prof. Claudio Gilberto Froehlich” and the task of facing the biodiversity knowledge deficits on Caddisflies (Trichoptera), Bahia, Brazil. *Revista Brasileira de Entomologia* 67 (spe): e20230065. <https://doi.org/10.1590/1806-9665-RBENT-2023-0065>
- Castelletti C.H.M., Santos A.M.M., Tabarelli M. & da Silva J. 2003. Quanto ainda resta da Caatinga? Uma estimativa preliminar. In: Leal I.R., Tabarelli M. & da Silva J.M.C. (eds) *Ecologia e Conservação da Caatinga*: 91–100. Universidade Federal de Pernambuco, Recife.
- Costa A.M., Quinteiro F.B. & Calor A.R. 2014. Trichoptera do Semiárido I: Annulipalpia. In: Bravo F. & Calor A.R. (eds) *Artrópodes do Semiárido: biodiversidade e conservação*: 225–228. Printmídia, Feira de Santana, BA.
- Gressitt J.L. & Gressitt M.K. 1962. An improved Malaise trap. *Pacific Insects* 4: 87–90.
- Henriques-Oliveira A.L., Spies M.R. & Dumas L.L. 2012. A new species of *Notalina* Mosely, 1936 (Trichoptera: Leptoceridae) from Southeastern Brazil. *Biota Neotropica* 12 (4): 130–134. <https://doi.org/10.1590/S1676-06032012000400014>
- Henriques-Oliveira A.L. Rocha I.C. & Nessimian J.L. 2018. Leptoceridae (Insecta, Trichoptera) from Serra da Canastra Mountain Range, Southeast Brazil: diversity, distribution, and description of two new species. *Neotropical Entomology* 48: 277–289. <https://doi.org/10.1007/s13744-018-0633-4>
- Henriques-Oliveira A.L., Silva A.L.R., Nessimian J.L. & Takiya D.M. 2021. New long-horned caddisfly genus and species of Leptoceridae (Insecta: Trichoptera) from Brazil. *Zootaxa* 5057 (4): 530–544. <https://doi.org/10.11646/zootaxa.5057.4.4>
- Holzenthal R.W. 1984. Studies in Neotropical Leptoceridae (Trichoptera) I: *Achoropsyche*, a new genus. In: Morse J.C. (ed.), *Proceedings of the 4th International Symposium on Trichoptera*: 181–184. Dr W. Junk, The Hague.
- Holzenthal R.W. 1986. The Neotropical species of *Notalina*, a southern group of long-horned caddisflies (Trichoptera: Leptoceridae). *Systematic Entomology* 11: 61–73. <https://doi.org/10.1111/j.1365-3113.1986.tb00165.x>
- Holzenthal R.W. 1989. Studies in Neotropical Leptoceridae (Trichoptera), IX: A new genus and species from southeastern Brazil. *Aquatic Insects* 11 (1): 29–32. <https://doi.org/10.1080/01650428909361344>
- Holzenthal R.W. & Calor R.A. 2017. Catalog of the Neotropical Trichoptera (Caddisflies). *ZooKeys* 654: 1–566. <https://doi.org/10.3897/zookeys.654.9516>
- Holzenthal R.W. & Pes A.M.O. 2004. A new genus of long-horned caddisfly from the Amazon basin (Trichoptera: Leptoceridae: Grumichellini). *Zootaxa* 621 (1): 1–16. <https://doi.org/10.11646/zootaxa.621.1.1>
- IBGE – Fundação Instituto Brasileiro de Geografia e Estatística 1977. *Geografia do Brasil. Região Nordeste. Vol. 2*. Centro de Serviços Gráficos do IBGE, Rio de Janeiro, RJ.
- Ito T., Kuhara N. & Katsuma N. 2013. The genus *Adicella* McLachlan (Trichoptera, Leptoceridae) in Japan. *Zootaxa* 3635 (1): 27–39. <https://doi.org/10.11646/zootaxa.3635.1.3>

- Kirby W. 1813. Strepsiptera, a new order of insects proposed; and the characters of the order, with those of its genera, laid down. *Transactions of the Linnean Society, London* 11 (1): 86–122.
<https://doi.org/10.1111/j.1096-3642.1813.tb00040.x>
- Kolenati F.A. 1859. Genera et species Trichopterorum, Pars Altera. *Nouveaux Mémoires de la Société Impériale des Naturalistes de Moscou* 11: 141–296.
- Leach W.E. 1815. Entomology. In: Brewster D (ed.) *Brewster's Edinburgh Encyclopedia* 9: 57–172. Blackwood, Edinburgh. <https://doi.org/10.5962/bhl.title.30911>
- Leal I.R., Da Silva J.M.C., Tabarelli M. & Lacher T.E. Jr. 2005. Changing the course of biodiversity conservation in the Caatinga of Northeastern Brazil. *Conservation Biology* 19 (3): 701–706.
<https://doi.org/10.1111/j.1523-1739.2005.00703.x>
- Linnaeus C. 1758. *Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Tomus I. Editio decima, reformata*. Laurentii Salvii, Holmiae [Stockholm]. <https://doi.org/10.5962/bhl.title.542>
- Malm T. & Johanson K.A. 2011. A new classification of the long-horned caddisflies (Trichoptera: Leptoceridae) based on molecular data. *BMC Evolutionary Biology* 11: Art. 10.
<https://doi.org/10.1186/1471-2148-11-10>
- Martynov A.V. 1924. Rucheiniki (caddisflies [Trichoptera]). In: Bogdanov-Kat'kov N.N. (ed.) *Prakticheskaya entomologiya, Volume 5: I–IV, 1–384*. Gosudarstvennoe izdatel'stvo, Leningrad. [In Russian.]
- McLachlan R. 1877. Fam. Leptoceridae. Part VI. In: *A Monographic Revision and Synopsis of the Trichoptera of the European Fauna. Part 6*: 281–348. John van Voorst, London.
- MMA – Instituto Chico Mendes 2007. *Plano de Manejo – Parque Nacional da Chapada Diamantina*. Versão preliminar, Documento de Trabalho – parte 1. Brasília, DF.
- Morse J.C. 1981. A phylogeny and classification of a family-group taxa of Leptoceridae (Trichoptera). *Proceedings of the 3rd International Symposium on Trichoptera, Series Entomologica* 20: 257–264.
- Morse J.C., Frandsen P.B., Graf W. & Thomas J.A. 2019. Diversity and ecosystem services of Trichoptera. *Insects* 10: 125. <https://doi.org/10.3390/insects10050125>
- Mosely M.E. 1936. A revision of the Triplectidinae, a subfamily of the Leptoceridae (Trichoptera). *Transactions of the Royal Entomological Society of London* 85 (3): 91–130.
<https://doi.org/10.1111/j.1365-2311.1936.tb00241.x>
- Mosely M.E. & Kimmins D.E. 1953. *The Trichoptera of Australia and New Zealand*. British Museum (Natural History), London. <https://doi.org/10.5962/bhl.title.118696>
- Müller F. 1879. Über Phryganiden (letters to his brother). *Zoologischer Anzeiger* 2: 38–40 + 180–182 + 283–284 + 404–407.
- Navás L. 1916. Neuroptera Nova Americana. Series I, II. *Memorie della Pontificia Accademia Romana dei Nuovi Lincei* 2: 59–80.
- Navás L. 1920. Insectos sudamericanos – 1^a, 2^a e 3^a séries. *Anales de la Sociedad Científica Argentina* 90: 33–72.
- Nessimian J.L., Santos A.P.M., Sampaio B.H.L., Dumas L.L., Pes A.M.O. & Ferreira Jr. N. 2024. The Collapsible Light Trap: a portable Pennsylvania Light Trap for collecting aquatic insects. *Anais da Academia Brasileira de Ciências* 96 (3): e20230784. <https://doi.org/10.1590/0001-3765202420230784>

Prado D.E. 2003. As Caatingas da América do Sul. In: Leal I.R., Tabarelli M. & da Silva J.M.C. (eds) *Ecologia e Conservação da Caatinga*: 3–75. Ed. Universitária da UFPE, Recife.

Quinteiro F.B., Costa A.M. & Calor A.R. 2014. Trichoptera do Semiárido II: Integripalpia. In: Bravo F. & Calor A.R. (eds) *Artrópodes do Semiárido: biodiversidade e conservação*: 229–243. Printmídia, Feira de Santana.

QGIS Development Team 2020. *QGIS Geographic Information System*. Open Source Geospatial Foundation. Available from <https://www.qgis.org/> [accessed 28 Mar. 2024].

Rafael J.A., Oliveira M.L., Souza U.R.M., Galileo M.H.M., Silva A.S., Mello F.V., Leivas F.W.T., Arriagada G., Grossi P.C., Ale-Rocha R., Takiya D.M., Câmara J., Marques D.W.A., Silva R.A.P.F., Souza E.A., Menezes I.S., Sousa A.A.T., Silva T.T.A., Santos L.L.M. ... & Limeira-de-Oliveira F. 2017. Capítulo 12: Insetos dos Parques Nacionais de Sete Cidades e Ubajara. In: Mantovani W., Monteiro R.F., Anjos L. & Cariello M.O. (Org.). *Pesquisas em unidades de Conservação no Domínio da Caatinga: Subsídios à gestão*: 301–313. Edições UFC, Fortaleza.

Silva-Pereira E., Oliveira I., Desidério G.R., Calor A. & Hamada N. 2022. *Notalina* (Neonotalina) *ralphi* sp. nov. (Trichoptera, Leptoceridae), a new long-horned caddisfly from the Cerrado biome of Brazil, with new records for *N. (Neonotalina) brasiliiana* Holzenthal, 1986 and an identification key. In: Pauls S.U., Thomson R. & Rázuri-Gonzales E. (eds) Special issue in honor of Ralph W. Holzenthal for a lifelong contribution to Trichoptera systematics. *ZooKeys* 1111: 413–424. <https://doi.org/10.3897/zookeys.1111.77581>

Takiya D.M., Santos A.P.M., Pinto A.P., Henriques-Oliveira A.L., Carvalho A., Sampaio B.H.L., Clarkson B., Moreira F.F.F., Avelino-Capistrano F., Gonçalves I.C., Cordeiro I., Câmara J., Barbosa J.F., de Souza W.R.M. & Rafael J.A. 2016. Aquatic Insects from the Caatinga: check lists and diversity assessments of Ubajara (Ceará State) and Sete Cidades (Piauí State) National Parks, Northeastern Brazil. *Biodiversity Data Journal* 14: e8354. <https://doi.org/10.3897/BDJ.4.e8354>

Ulmer G. 1903. Über die Metamorphose der Trichopteren. *Abhandlungen des Naturwissenschaftlichen Vereins in Hamburg* 18: 1–154

Ulmer G. 1906. Neuer Beitrag zur Kenntnis aussereuropäischer Trichopteren. *Notes from the Leyden Museum* 28: 1–116.

Ulmer G. 1908. Trichoptera und Ephemeridae. In: Michaelsen W. & Hartmeyer R. (eds) *Fauna Südwest-Australiens* 2: 25–40. Jena.

Printed versions of all papers are deposited in the libraries of four of the institutes that are members of the *EJT* consortium: Muséum national d'Histoire naturelle, Paris, France; Meise Botanic Garden, Belgium; Royal Museum for Central Africa, Tervuren, Belgium; Royal Belgian Institute of Natural Sciences, Brussels, Belgium. The other members of the consortium are: Natural History Museum of Denmark, Copenhagen, Denmark; Naturalis Biodiversity Center, Leiden, the Netherlands; Museo Nacional de Ciencias Naturales-CSIC, Madrid, Spain; Leibniz Institute for the Analysis of Biodiversity Change, Bonn – Hamburg, Germany; National Museum of the Czech Republic, Prague, Czech Republic; The Steinhardt Museum of Natural History, Tel Aviv, Israël.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [European Journal of Taxonomy](#)

Jahr/Year: 2025

Band/Volume: [1004](#)

Autor(en)/Author(s): Henriques-Oliveira Ana L., Alves Andre Almeida, Moreira Felipe Ferraz Figueiredo

Artikel/Article: [New species of Notalina Mosely, 1939 \(Trichoptera: Leptoceridae\) from Chapada Diamantina National Park, Northeast Brazil, with new records of other leptocerids from the Caatinga biome 1-18](#)