



Entomofauna

ZEITSCHRIFT FÜR ENTOMOLOGIE

Band 30, Heft 24: 437-444

ISSN 0250-4413

Ansfelden, 16. Oktober 2009

**Faunistic Note on the Braconidae
(Hymenoptera: Ichneumonoidea)
in Iranian Alfalfa Fields and Surrounding Grasslands**

**H. GHAHARI¹, M. FISCHER, O.C. ERDOGAN, A. BEYARSLAN,
K.J. HEDQVIST & H. OSTOVAN**

Abstract

A total of six braconid species (Hymenoptera: Braconidae) of five genera (*Agathis* LATREILLE, *Apanteles* FÖRSTER, *Bassus* FABRICIUS, *Bracon* FABRICIUS and *Chorebus* HALIDAY) and four subfamilies (Agathidinae, Braconinae, Microgastrinae and Alysiinae) were collected from Iranian alfalfa fields and surrounding grasslands.

Key words: Braconidae, Alfalfa fields, Fauna, New record, Iran

Zusammenfassung

Sechs Braconidenarten (Hymenoptera: Braconidae) aus fünf Gattungen (*Agathis* LATREILLE, *Apanteles* FÖRSTER, *Bassus* FABRICIUS, *Bracon* FABRICIUS und *Chorebus*

¹ The first author Hassan Ghahari is responsible for the contents of this article.

HALIDAY) und vier Unterfamilien (Agathidinae, Braconinae, Microgastrinae und Alysiinae) wurden auf iranischen Lutzernefeldern und umliegendem Grünland nachgewiesen.

Introduction

The Braconidae constitute the second largest family of Hymenoptera. This family is of cosmopolitan distribution and diverse in all areas, with no strong preference for tropical or temperate regions or for wet or dry habitats (SHARKEY & WAHL, 1992; GHAHARI et al., 2006). The most common hosts of these beneficial insect are Lepidoptera larvae, followed by Coleoptera larvae and Diptera larvae. Other hosts include Hemiptera nymphs and adults, Orthoptera and Psocoptera nymphs, adult Hymenoptera, Coleoptera, and Chrysopidae (Neuroptera), and Symphyta larvae (TOBIAS, 1967; MATTHEWS, 1974; WHARTON, 1993).

Alfalfa (*Medicago sativa*) is one of the most important legumes used in agriculture. Alfalfa is widely grown throughout the world as forage for cattle, and is most often harvested as hay, but can also be made into silage, grazed, or fed. Alfalfa has the highest feeding value of all common hay crops, being used less frequently as pasture. When grown on soils where it is well-adapted, alfalfa is the highest yielding forage plant (HANSON et al., 1988; PARKER & PARKER, 2003).

There are several insect pests in alfalfa fields which damage to this crop in different seasons. These pests are attacked by many natural enemies including predators and parasitoids. The braconid wasps are one of these beneficial insects which play an effective role in biological control programs. Despite of the importance of these efficient parasitoids in almost all agroecosystems, their diversity was not perfectly studied in alfalfa fields of the world so far. In this paper, we deal with the faunistic diversity of Braconidae in some Iranian alfalfa fields. The goal of this work is determining the efficient parasitoids in alfalfa fields for decreasing the pesticides' application.

Materials and Methods

Specimens were collected by sweep netting and malaise traps from different regions of Iran which contain alfalfa fields. These sampled regions were Ardabil (Ardabil province), Bojnord (Khorasan province), Gonbad (Golestan province), Hamadan (Hamadan province), Maragheh (East Azarbayan province) and Ourmiah (West Azarbayan province). The samplings were conducted between 2004 and 2007, and the collected specimens were killed with ethyl acetate and mounted on triangular labels and were examined with a stereoscopic binocular microscope.

Results

Up on the conducted faunistic survey on the Braconidae of alfalfa fields, totally 6 species from 5 genera were identified as the new records for fauna of Iran. The list of species is given below together with the general distribution data.

Subfamily Agathidinae

Genus *Agathis* LATREILLE, 1804

Agathis fuscipennis (ZETTERSTEDT, 1838)

Material examined: West Azarbayjan province: Ourmieh (2♂♂), 18.IX.2006.

General Distribution: Germany, Finland, Sweden (FISCHER, 1957, 1968; Early indications of distribution of *Agathis* in Palearctic region; SHENEFELT, 1970); Austria, Bulgaria, England, Ireland (NIXON, 1986); Switzerland (SIMBOLOTTI & ACHTERBERG, 1999); Turkey (ZETTEL and BEYARSLAN, 1992; CETIN & BEYARSLAN, 2001).

Genus *Bassus* FABRICIUS, 1804

Bassus armeniacus (TELENKA, 1955)

Material examined: East Azarbayjan province: Maragheh (1♂), 9.VI.2006.

General Distribution: Austria, Russia, Turkey (NIXON, 1986; CETIN & BEYARSLAN, 2001).

Subfamily Alysiinae

Tribe *Dacnusini*

Genus *Chorebus* HALIDAY, 1833

Chorebus (Phaenolexis) posticus (HALIDAY, 1839)

Material examined: Hamadan province: Hamadan (1♀), 13.VII.2006.

General Distribution: Widely distributed in the Palaearctic region (PAPP, 2005).

Subfamily Braconinae

Genus *Bracon* FABRICIUS, 1804

Bracon (Bracon) leptus MARSHALL, 1897

Material examined: Khorasan province: Bojnord (1♀), 9.X.2007.

General Distribution: Austria, Caucasus, Far East, Hungary, Kazakhstan, Siberia, Spain, Tajikistan, Turkmenistan, Ukraine, Uzbekistan (PAPP, 1968; SHENEFELT, 1978; TOBIAS, 1995), Turkey (BEYARSLAN et al., 2005; GÜLER & ÇAĞATAY, 2007).

***Bracon (Bracon) pectoralis* WESMAEL, 1838**

Material examined: Golestan province: Gonbad (2♀♀), 7.IV.2005.

General Distribution: Albania, Austria, Azerbaijan, Belgium, Bulgaria, Caucasia, England, France, Germany, Hungary, Italy, Kazakhstan, Russia, Spain, Tunisia, Turkmenistan, Ukraine, Yugoslavia (PAPP, 1968; SHENEFELT, 1978; TOBIAS, 1995), Turkey (BEYARSLAN et al., 2005; GÜLER & ÇAĞATAY, 2007).

Subfamily Microgastrinae

***Apanteles* FÖRSTER, 1862**

***Apanteles (Apanteles) emarginatus* (NEES, 1834)**

Material examined: Ardabil province: Ardabil (1♀), 22.VIII.2004.

General Distribution: Europe, Armenia, Azerbaidzhan (PAPP, 1978), Turkey (INANC & ERDOGAN, 2004).

Discussion

Alfalfa is a vigorous and productive crop- Like all farm crops; however, alfalfa is subject to stand injury and yield loss from disease, insect injury and nutrient deficiencies. Prompt and accurate diagnosis of a problem can allow early treatment to modify or correct the situation before yields are seriously affected or the situation stands are lost (WILLIS, 1983; JENNINGS, 1999). Although several natural enemies were determined from Iranian alfalfa fields, but pesticides are applied severely in almost fields. The main goal of pesticides' application in alfalfa fields is control of alfalfa weevil, *Hypera postica* (GYLLENHAL), which is a key pest of alfalfa in almost regions of the world. In addition to *H. postica*, there are many other important pests including, Alfalfa Weevil, *Hypera postica* (GYLLENHAL), Clover leaf weevil, *Hypera punctata* (FABRICIUS), Clover root curculio, *Sitona hispidulus* (FABR.), Alfalfa snout beetle, *Brachyrhinus ugustici* (L.), Potato leafhopper, *Empoasca fabae* (HARRIS), Pea aphid, *Acyrthosiphon pisum* (HARRIS), Spotted Alfalfa aphid, *Theroaphis maculata* (BUCKTON), Meadow Spittle bug, *Philaenus spumarius* (L.), Variegated cutworm, *Peridroma saucia* (HÜBNER), Differential grasshopper, *Melanopus differentialis* (THOMAS) and Plant bugs, *Lygus lineolaris* (PALISOT de BEAUVOIS) (DAY, 1999; PARKER & PARKER, 2003).

Insect pest management in an organic system depends on several factors including climate, beneficial organisms already present in the area, and hay-cutting schemes. Many types of insects and mites inhabit alfalfa plantings, yet only a few species threaten yields. Proper identification of alfalfa pests as well as their natural enemies is the first step in successful management of pests. Some local Extension service specialists are familiar with pests common to specific areas and can help with proper identification. After

determining of all the pests and natural enemies in an agroecosystem, start to prepare a proper IPM program and continuing it will be resulted the successful of pest control (FLINT & DREISTADT, 1998; GHAHARI et al. 2008).

In this research, although the braconids' fauna of Iranian alfalfa fields was preliminary studied, but there are several other alfalfa fields in different regions of Iran which were not sampled in this project. Continuing this research in other Iranian alfalfa fields surely will be resulted to many other new records and probably new species in the family Braconidae. Therefore, we suggest strongly that the Iranian researchers especially the top researchers of different branches of Islamic Azad University carry on these invaluable research topics for solving the crop loss problem step by step.

Acknowledgements

The authors are indebted to Dr. T. Finlayson of Canada and Dr. Kees van Achterberg of The Netherlands for invaluable helps in progress of the project. The research was supported by Shahre Rey Islamic Azad University and Naturhistorisches Museum Vienna of Austria.

References

- BEYARSLAN A., ÇETİN Ö. & M. AYDOĞDU (2005): A Survey Braconinae (Hymenoptera, Braconidae) of Turkish Western Black Sea Region. — Linzer biol. Beitr. **37** (1): 195-213.
- DAY W.H. (1999): Host preferences of introduced and native parasites (Hymenoptera: Braconidae) of phytophagous plant bugs (Hemiptera: Miridae) in alfalfa-grass fields in the northeastern U.S.A. — BioControl **44**: 249-261.
- CETİN O. & A. BEYARSLAN (2001): The Agathidinae (Hymenoptera: Braconidae) Fauna of the Marmara Region. — Turkish J. Zoology **25**: 257-268.
- FISCHER M. (1957): Beiträge zur Kenntnis der paläarktischen Braconiden (Hymenoptera) — Mitteilungen der Münchener Entomologischen Gesellschaft **47**: 1-21, table XVIII (with *Agathis*).
- FISCHER M. (1968): Über gezüchtete Raupenwespen (Hymenoptera, Braconidae) — Pflanzenschutz-Berichte **37** (7, 8, 9): 97-140 (key for *Agathis*, palearctic species, as far as Mongolia, China).
- FLINT M.L. & S.H. DREISTADT (1998): Natural Enemies Handbook; The Illustrated Guide to Biological Pest Control. — University of California Statewide IPM Project. Publication **3386**. 63 pp.
- GHAHARI H., YU, D.S. & C. van ACHTERBERG (2006): World Bibliography of the Family Baraconidae (Hymenoptera: Ichneumonoidea) (1964-2003). — NNM Technical Bulletin **8**: 293 pp.
- GHAHARI H., HAYAT R., TABARI M., OSTOVAN H. & S. IMANI (2008): A contribution to the predator and parasitoid fauna of rice pests in Iran, and a discussion on the biodiversity and IPM in rice fields. — Linzer biol. Beitr. **40** (1): 735-764.

- GÜLER Y. & N. ÇAĞATAY (2007): A study on the genus *Bracon* FABRICIUS 1804 (Hymenoptera, Braconidae) in Ankara province in Turkey — Mun. Entomol. Zool. **2**(2): 517-524.
- HANSON A.A., BARNES D.K. & R.R. Jr. HILL (1988): Alfalfa and Alfalfa Improvement, — American Society Agronomy Monograph **29**. Madison, WI. 322 pp.
- INANC F. & C.O. ERDOGAN (2004): Contribution to the Microgastrinae (Hymenoptera, Braconidae) fauna of Turkey, with description of a new species of *Dolichogenidea*. — Biologia, Bratislava **59** (5): 547-551.
- JENNINGS J. (1999): Understanding autotoxicity of alfalfa. 2001. — Proceedings of WI Forage Council 25th Forage Production and Use Symposium, pp. 110-116.
- MATTHEWS R.W. (1974): Biology of Braconidae. — Ann. Rev. Entomol. **19**: 15-32.
- NIXON G.E.J. (1986): A revision of the European Agathidinae (Hymenoptera: Braconidae), — Bull. Brit. Mus. nat. Hist. (Ent). Ser. **52** (3): 183-242.
- PAPP J. (1968): A Synopsis of the Bracon Fabr. Species of the Carpathian Basin, Central Europe (Hymenoptera, Braconidae), II. Subgenus *Bracon* FABRICIUS. — Annales Historico-Naturales Musei Nationalis Hungarici **60**: 195- 211.
- PAPP J. (1978): A survey of the European species of *Apanteles* FOERSTER (Hym.: Braconidae: Microgastrinae) II. The leavigatus group 1.s — Ann. His. Natur. Mus. Nat. Hung. **70**: 265-300.
- PAPP J. (2005): Braconidae (Hymenoptera) from Mongolia XVI. subfamilies Gnaphodontinae, Brachitinae, Euphorinae, Alysiinae. — Acta Zoologica Academiae Scientiarum Hungaricae **51** (3): 221-251.
- PARKER J.N. & P.M. PARKER (2003): Alfalfa - A Medical Dictionary, Bibliography, and Annotated Research Guide to Internet References. ICON Group International Inc., 192 pp. Sharkey M.J. & D.B. Wahl 1992. Cladistics of the Ichneumonoidea (Hymenoptera). Journal of Hymenoptera Research **1** (1): 15-24.
- SHENEFELT R.D. (1970): Hymenopterum Catalogus (nov. ed.), Part 6. Braconidae **3**: 307-428.
- SHENEFELT R.D. (1978): Hymenopterum Catalogus (nov. ed.). Pars 15- Braconidae 10, Junk, S-Gravenhage **10**: 1459-1638.
- SIMBOLOTTI G. & C. van ACHTERBERG (1999): Revision of the West Palearctic species of the genus *Agathis* LATREILLE (Hymenoptera: Braconidae: Agathidinae). — Zool. Verh. Leiden **325**: 1-167.
- TOBIAS V.I. (1967): A review of the classification, phylogeny and evolution of the family Braconidae (Hymenoptera). — Entomol. Rev. **46**: 387-399.
- TOBIAS V.I. (1995): Keys of the Insects of the European Part of the USSR. Volume 3, Hymenoptera, Part 4, — Science Publishers, Lebanon, New Hampshire, USA, xvi+883 pages.
- WHARTON R.A. (1993): Bionomics of the Braconidae. — Ann. Rev. Entomol. **38**: 121-143.
- WILLIS H. (1983): How to Grow Great Alfalfa. — Harold L. Willis, Publisher. 44 p.
- ZETTEL H. & A. BEYARSLAN (1992): Über Agathidinae aus der Türkei (Hymenoptera: Braconidae). — Entomofauna, Zeitschrift für Entomologie **13** (5): 121-132.

Addresses of the authors:

Hassan GHAHARI
Department of Agriculture
Islamic Azad University
Shahre Rey Branch, Tehran, Iran
E-mail: h_ghahhari@yahoo.com

Maximilian FISCHER
Hofrat i.R. Univ.-Doz.,
Naturhistorisches Museum,
2. Zoologische Abteilung,
A-1010 Wien, Burgring 7, Austria;
E-mail: maximilian.fischer@chello.at

Ozlem Cetin ERDOGAN
Ahmet BEYARSLAN
Trakya University, Faculty of Arts and Science,
Department of Biology, 22030 Edirne, Turkey;
E-mail: abeyars@trakya.edu.tr;
cetinozlem@hotmail.com

Karl Johan HEDQVIST
Höstvägen 1,
SE-186 31 Vallentuna, Sweden

Hadi OSTOVAN
Department of Entomology
Islamic Azad University, Fars Science & Research Branch,
Marvdasht, Iran
E-mail: ostovan2001@yahoo.com

Druck, Eigentümer, Herausgeber, Verleger und für den Inhalt verantwortlich:
Maximilian SCHWARZ, Konsulent f. Wissenschaft der Oberösterreichischen Landesregierung, Eibenweg 6,
A-4052 Ansfelden, E-Mail: maximilian.schwarz@liwest.at.

Redaktion: Erich DILLER, ZSM, Münchhausenstraße 21, D-81247 München;
Fritz GUSENLEITNER, Lungitzerstr. 51, A-4222 St. Georgen/Gusen;
Wolfgang SCHACHT, Scherrerstraße 8, D-82296 Schöngeising;
Wolfgang SPEIDEL, MWM, Tengstraße 33, D-80796 München;
Thomas WITT, Tengstraße 33, D-80796 München.

Adresse: Entomofauna, Redaktion und Schrifttausch c/o Museum Witt, Tengstr. 33, 80796 München,
Deutschland, E-Mail: thomas@witt-thomas.com; Entomofauna, Redaktion c/o Fritz Gusenleitner,
Lungitzerstr. 51, 4222 St. Georgen/Gusen, Austria, E-Mail: f.gusenleitner@landesmuseum.at

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Entomofauna](#)

Jahr/Year: 2009

Band/Volume: [0030](#)

Autor(en)/Author(s): Ghahari Hassan, Fischer Maximilian (Max), Erdogan Özlem Çetin, Beyarslan Ahmet, Hedqvist Karl Johan, Ostovan Hadi

Artikel/Article: [Faunistic Note on the Braconidae \(Hymenoptera: Ichneumonoidea\) in Iranian Alfalfa Fields and Surrounding Grasslands. 437-444](#)