



# Entomofauna

ZEITSCHRIFT FÜR ENTOMOLOGIE

---

Band 37, Heft 12: 217-224

ISSN 0250-4413

Ansfelden, 4. Januar 2016

---

## Coleoptera fauna of poultry litter in Izmir province of Turkey (Coleoptera)

Utku SANVER & Serdar TEZCAN

### Abstract

This study was carried out to determine the species of Coleoptera existing in poultry litter of Izmir province of western Turkey during July-December 2013. At the end of this study, five different species belonging to three families of Coleoptera were determined. The species reported from Izmir for the first time were *Carcinops pumilio* (ERICHSON 1834), *Saprinus caerulescens* (HOFFMANN 1803) (Histeridae) and *Harpalus fuscicornis* MÉNÉTRIÉS (Carabidae).

### Zusammenfassung

In dieser Studie werden die Käferarten untersucht, die in einer Geflügelfarm in der Izmir-Provinz (West-Türkei) zwischen Juli und Dezember 2013 gefunden wurden. Abschließend konnten fünf verschiedene Arten aus drei Familien der Coleoptera determiniert werden. Für die Provinz Izmir werden zum ersten Mal die Arten *Carcinops*

*pumilio* (ERICHSON 1834), *Saprinus caerulescens* (HOFFMANN 1803) (Histeridae) und *Harpalus fuscicornis* MÉNÉTRIÉS (Carabidae) gemeldet.

## Introduction

Poultry husbandry occupy an important place in the growing Turkish economy. Chicken's meal production has been reported about 1.6 million tons in Turkey (GHANIZADEH 2014). Being an important sector in the country's economy, it has been widely practiced in all provinces of Turkey. However production from Izmir province has a significant contribution to the overall production of Turkey (AKBAY 1990). A huge studies have been documented by many researchers regarding pests and the beneficial species occurring in poultry litter and poultry houses in many countries across the world (PFEIFFER & AXTELL 1980, STAFFORD et al. 1988, HALD et al. 1998, PINTO et al. 2009). However, no study on the insect fauna of poultry litter in Turkey has been conducted so far. Therefore, the present study was executed with aim of monitoring the presence of Coleoptera fauna in poultry litter of Izmir province of western Turkey.

## Material and Methods

The study on Coleoptera fauna of poultry litter in Izmir province of Turkey was conducted in 120 poultry houses with area ranging between  $1182.6 \pm 68.2 \text{ m}^2$  (1114.4-1250.8) ( $n=120$ ) of 17 different districts (Aliaga, Bayindir, Bergama, Buca, Cesme, Dikili, Foca, Guzelbahce, Kemalpasa, Kinik, Menderes, Menemen, Odemis, Seferihisar, Tire, Torbali, Urla) of Izmir, Turkey. The samples were collected from different poultry litter and brought to the laboratory during the months of July-December 2013. At three points, an area of about  $0.1 \text{ m}^2$  was chosen randomly in each poultry house and litter samples were picked up carefully by hand trowel. It is important to mention that to avoid contamination risk and to ensure health safety, each time sterile gloves and mask were used for sample collection. The samples were carefully examined in the laboratory and insect specimens were collected from the samples litter. The collected specimens were killed in 70% ethyl-alcohol, mounted and identified using identification keys. The districts and the number of poultry houses which were sampled were given in Table 1 and 2.

## Result and Discussion

The results of the present study revealed a total of five species belonging to three families of Coleoptera. These species were *Alphitobius diaperinus* (PANZER 1797), *Tribolium castaneum* (HERBST 1797) (Tenebrionidae), *Carcinops pumilio* (ERICHSON 1834), *Saprinus caerulescens* (HOFFMANN 1803) (Histeridae) and *Harpalus fuscicornis* MÉNÉTRIÉS 1832 (Carabidae).

Among these, *Alphitobius diaperinus* was found the most abundant one which was sampled from the 93.33% of the poultry houses in Izmir province (Table 1). It was followed by *Carcinops pumilio* (29 specimens), *Tribolium castaneum* (3 specimens), *Saprinus caerulescens* (1 specimen) and *Harpalus fuscicornis* (1 specimen) (Table 2).

The *Alphitobius diaperinus* is a polyphagous species that has also been found very common in poultry litter in many countries (ROWLAND et al., 2007). The main mode of damage of this pest is to bring destruction to the poultry structures as reported by ROWLAND et al. (2007). The same has also been reported to play important role in transmission of many poultry diseases as a vector and to reduce weight grain of chicks (CALIBEO 2002; CRIPPEN et al. 2012).

Among other pests, *Tribolium castaneum* is also a pest of stored products (OZER 1957; JBILOU et al. 2006; BAKHTAWAR et al. 2013; MAGDA et al. 2014).

*Carcinops pumilio* is an effective predator of *Musca domestica* LINNAEUS 1758 (GEDEN & AXTELL 1988; FLETCHER et al. 1991; ANONYMOUS 2007) whereas *Saprinus caerulescens* feeds on carcasses of animals (OZDEMIR & SERT 2008; KARAPAZARLIOĞLU & OZMAN-SULLIVAN 2011).

*Harpalus fuscicornis* is known to exhibit an omnivorous feeding behavior (KOCATEPE & MERGEN 2004).

In the present study, three species *Carcinops pumilio*, *Saprinus caerulescens* and *Harpalus fuscicorni* were reported for the first time from Izmir province which is one of the main findings of this research work. However, further similar studies in the other parts of Turkey will be very fruitful to increase the sources of information in this field.

### Acknowledgement

We sincerely would like to thank to Assoc. Prof. Dr. Bekir Keskin (Ege University, Izmir, Turkey), Dr. Giovanni Ratto (Genova, Italy) and Dr. Memiş Kesdek ((Muğla University, Fethiye, Turkey) for their great support regarding the identification and confirmation of the collected material of Tenebrionidae, Histeridae and Carabidae, respectively. We wish to give special thanks to Muhammad Hamayoon Khan for proofreading of manuscript.

### Literature

- AKBAY R. (1990): Poultry Production in Turkey. – <http://om.ciheam.org/om/pdf/a07/CI901605.pdf>. – (Access date: 20 May 2015).
- ANONYMOUS (2007): Insect guide: *Carcinops*. [http://www.pestcontrolexpert.com/bayer/cropscience/bespestcontrol.nsf/id/8502501CEF762DB8C1257984005605BC/\\$file/Carcinop\\_08.pdf](http://www.pestcontrolexpert.com/bayer/cropscience/bespestcontrol.nsf/id/8502501CEF762DB8C1257984005605BC/$file/Carcinop_08.pdf). – (Access date: 4 February 2014).
- BAKHTAWAR M., SAEED Q., ZAKA S.M., SAEED S., ANSARI T., IDREES N., JALEEL W., NAQQASH M.N. & M.R. AKRAM (2013): Role of storage facilities for insect pest control strategies in Punjab, Pakistan. – Journal of Stored Products and Postharvest Research 4 (4): 55-60.

- CALIBEO D.R. (2002): Role and Mitigation of Two Vectors of Turkey Coronavirus, *Musca domestica* L. and *Alphitobius diaperinus* PANZER. – MSc Thesis, North Carolina State University, 127 pp.
- CRIPPEN T.L., ZENG L., SHEFFIELD C.L., TOMBERLIN J. K., BEIER R.C. & Z. YU (2012): Transient gut retention and persistence of *Salmonella* through metamorphosis in the lesser mealworm, *Alphitobius diaperinus* (Coleoptera: Tenebrionidae). – Journal of Applied Microbiology **112** (1): 920-926.
- FLETCHER M.G., AXTELL R.C., STINNER R.E. & L.R. WILHOIT (1991): Temperature-dependent development of immature *Carcinops pumilio* (Coleoptera: Histeridae), a predator of *Musca domestica* (Diptera: Muscidae). – Journal of Entomological Science **26** (1): 99-108.
- GEDEN C.J. & R.C. AXTELL (1988): Predation by *Carcinops pumilio* (Coleoptera: Histeridae) and *Macrocheles muscaedomesticae* (Acarina: Macrochelidae) on the house fly (Diptera: Muscidae): functional response, effects of temperature, and availability of alternative prey. – Journal of Entomology Society of America **17** (4): 739-744.
- GHANIZADEH L. (2014): Adana ve Urmia örneğinde bazı etlik piliç kümelerinde yapı elemanlarının uygunluğunun değerlendirilmesi. – Yüksek Lisans Tezi, Çukurova Üniversitesi Fen Bilimleri Enstitüsü, 94. (unpublished).
- JBILOU R., ENNABILI A. & F. SAYAH (2006): Insecticidal activity of four medicinal plant extracts against *Tribolium castaneum* (HERBST) (Coleoptera: Tenebrionidae). – African Journal of Biotechnology **5** (10): 936-940.
- HALD B., OLSEN A. & M. MADSEN (1998): *Typhaea stercorea* (Coleoptera: Mycetophagidae) a carrier of *Salmonella enterica* serovar *infantis* in a Danish broiler house. – Journal of Economic Entomology **91** (1): 660-664.
- KARAPAZARLIOĞLU E. & S.K. OZMAN-SULLIVAN (2011): Kapalı ortamda domuz karkasları üzerinde gelen böcek türlerinin ve süksesyonlarının belirlenmesi ve bir örnek vaka çalışması. – Türkiye IV. Bitki Koruma Kongresi Bildirileri Kitapçığı, Kahramanmaraş, 39pp.
- KOCATEPE N. & O. MERGEN (2004): Ankara ili Carabidae (Coleoptera) familyası türleri üzerinde faunistik araştırmalar. – Türkiye Entomoloji Dergisi **28** (4): 295-309.
- MAGDA S., FODA M.S. & M.E. MOHARAM (2014): Evaluations of *Bacillus* species against red flour beetle *Tribolium castaneum* and confused flour beetle, *Tribolium confusum* (Coleoptera: Tenebrionidae) under laboratory and store conditions. – European Journal of Academic Essays **1** (10): 36-41.
- OZDEMIR S. & O. SERT (2008): Systematic studies on male genitalia of Coleoptera species found on decomposing pig (*Sus scrofa* L.) carcasses at Ankara province. – Hacettepe Journal of Biology and Chemistry: 137-161.
- OZER M. (1957): Türkiye'de Depo, Ambar, Fabrika ve Silolarda Muhtelif Hububat Taneleri, Un ve Mamulleri ile Kuru Meyvalar ve Tütünlerde Önemli Zarar Yapan Böcek Türlerinin Morfolojileri, Kısa Biyolojileri ve Yayılışları Üzerine Araştırmalar. – Ankara Üniversitesi Basimevi, Ankara, 136pp.
- PFEIFFER D.G. & R.C. AXTELL (1980): Coleoptera of poultry manure in cagedlayer houses in North Carolina. – Environmental Entomology **9** (21): 21-28.
- PINTO D.M., DUARTE J.L.P., RIBEIRO P.B. & P.S. JUNIOR (2009): Collection of Coleoptera from a poultry in Pelotas. – Rio Grande do Sul, Brazil, Ciência Rural **39** (2): 319-324.

ROWLAND R., MACKLIN K., SIMPSON G., DONALD J. & A. CAMPBELL (2007): Understanding and Controlling Litter Beetles. <http://www.aces.edu/poultryventilation/documents/Nwsltr50DarklingBeetles.pdf>. – (Access date: 31 July 2013).

STAFFORD K.C., COLLISON C.H., BURG J.G. & J.A. CLOUD (1988): Distribution and monitoring lesser mealworms, hide beetle and other fauna in high-rise caged-layer poultry house. – Journal of Agricultural Entomology **5** (2): 89-101.

**Table 1:** Related information about *Alphitobius diaperinus* in poultry houses of different districts of Izmir province in 2013.

Districts	Total number of sampled poultry	Number of poultry houses occurring	Rate of infection (%)	T					Total	Rate (%)
				Adult	t o	t a	l n	p u m		
Aliağa	5	5	100	50	848	1	899	1.60		
Bayındır	1	1	100	23	290	0	313	0.56		
Bergama	5	2	40	0	13	0	13	0.02		
Buca	1	1	100	242	466	15	723	1.30		
Çeşme	3	3	100	63	5849	14	6126	10.92		
Dikili	3	3	100	86	157	0	243	0.43		
Foça	7	7	100	425	3541	4	3970	7.07		
Güzelbahçe	1	1	100	26	428	7	461	0.82		
Kemalpaşa	35	35	100	2388	12535	142	15065	26.86		
Kınık	7	7	100	244	1011	3	1258	2.24		
Menderes	1	1	100	80	1448	28	1556	2.77		
Menemen	2	2	100	40	1307	41	1388	2.48		
Ödemiş	9	9	100	142	35	0	177	0.31		
Seferihisar	1	1	100	140	1333	5	1478	2.64		
Tire	10	5	50	59	127	0	186	0.33		
Torbalı	28	28	100	4791	17316	111	22218	39.60		
Urla	1	1	100	0	4	0	4	0.01		
<b>TOTAL</b>	<b>120</b>	<b>112</b>		<b>8999</b>	<b>46708</b>	<b>371</b>	<b>56078</b>	<b>100.00</b>		
<b>Rate (%)</b>				<b>16.04</b>	<b>83.29</b>	<b>0.60</b>				

Table 2: Related information about other species in poultry houses of different districts of Izmir province in 2013.

Districts	Total number of sampled poultry houses	<i>Tribolium castaneum</i>		<i>Carcinops pumilio</i>		<i>Saprinus caeruleascens</i>		<i>Harpalus fuscicornis</i>	
		Number of poultry houses occurring insect specimens	Number of adults	Number of poultry houses occurring insect specimens	Number of adults	Number of poultry houses occurring insect specimens	Number of adults	Number of poultry houses occurring insect specimens	Number of adults
Aliaga	5	0	0	0	0	0	0	0	0
Bayındır	1	0	0	0	0	0	0	0	0
Bergama	5	0	0	0	0	0	0	0	0
Büca	1	0	0	1	1	0	0	0	0
Çeşme	3	0	0	0	0	0	0	0	0
Dikili	3	0	0	0	0	0	0	0	0
Foca	7	0	0	0	0	0	0	0	0
Güzelbahçe	1	0	0	0	0	0	0	1	1
Kemalpaşa	35	2	2	4	1	1	0	0	0
Kimlik	7	0	0	0	0	0	0	0	0
Menderes	1	0	0	0	0	0	0	0	0
Menemen	2	0	0	0	0	0	0	0	0
Ödemiş	9	0	0	0	0	0	0	0	0
Seferihisar	1	0	1	23	0	0	0	0	0
Tire	10	0	0	0	0	0	0	0	0
Torbalı	28	1	1	1	1	0	0	0	0
Urla	1	0	1	0	0	0	0	0	0
<b>TOTAL</b>	<b>120</b>	<b>3</b>	<b>3</b>	<b>5</b>	<b>29</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>

Authors' addresses:

Utku SANVER  
Ege University, Faculty of Agriculture, Department of Plant Protection  
TR-Bornova-Izmir, Turkey  
E-mail: [utkussanver@hotmail.com](mailto:utkussanver@hotmail.com)

Prof. Dr. Serdar TEZCAN  
Ege University, Faculty of Agriculture, Department of Plant Protection  
TR-Bornova-Izmir, Turkey  
E-mail: [serdar.tezcan@gmail.com](mailto:serdar.tezcan@gmail.com)

---

Druck, Eigentümer, Herausgeber, Verleger und für den Inhalt verantwortlich:  
Maximilian SCHWARZ, Konsulent f. Wissenschaft der Oberösterreichischen Landesregierung, Eibenweg 6, A-  
4052 Ansfelden, Austria; [maximilian.schwarz@liwest.at](mailto:maximilian.schwarz@liwest.at).

Redaktion: Fritz GUSENLEITNER, Biologiezentrum Linz, [f.gusenleitner@landesmuseum.at](mailto:f.gusenleitner@landesmuseum.at);  
Roland GERSTMAYER, Lehrstuhl f. Zoologie, TU München, [gerstmayer@wzw.tum.de](mailto:gerstmayer@wzw.tum.de);  
Thomas WITT, Tengstraße 33, D-80796 München, [thomas@witt-thomas.com](mailto:thomas@witt-thomas.com);  
Berthold CLEWING, Akademischer Verlag München, [avm@druckmedien.de](mailto:avm@druckmedien.de);  
Harald SULAK, Museum Witt München, [h.sulak@atelier-sulak.de](mailto:h.sulak@atelier-sulak.de).

Mitarbeiter: Karin TRAXLER, Biologiezentrum Linz, [bio.redaktion@landesmuseum.at](mailto:bio.redaktion@landesmuseum.at);  
Heike REICHERT, Museum Witt München, [heike\\_reichert66@web.de](mailto:heike_reichert66@web.de);  
Erich DILLER, Zool. Staatssammlung München, [erich.diller@zsm.mwn.de](mailto:erich.diller@zsm.mwn.de).

Adresse: Entomofauna, Redaktion und Schrifttausch Thomas WITT, c/o Museum Witt München,  
Tengstr. 33, 80796 München, Deutschland, [thomas@witt-thomas.com](mailto:thomas@witt-thomas.com);  
Entomofauna, Redaktion c/o Fritz GUSENLEITNER, Lungitzerstr. 51, 4222 St. Georgen/Gusen,  
Austria, [f.gusenleitner@landesmuseum.at](mailto:f.gusenleitner@landesmuseum.at).

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Entomofauna](#)

Jahr/Year: 2016

Band/Volume: [0037](#)

Autor(en)/Author(s): Sanver Utku, Tezcan Serdar

Artikel/Article: [Coleoptera fauna of poultry litter in Izmir province of Turkey](#)  
[\(Coleoptera\) 217-224](#)